Just below the clavicle, about its middle point, a very strong, but not heaving, pulsation was felt over a very small extent. A short, dull sound was heard with the pulse. The pulse was the same at both wrists.

Saturnine lotions were applied to the part, and the pulsation gradually diminished. He was kept under observation for a long time without any fresh symptoms. On the last day of July, it was noted that there was much less pulsation than on his admission, but still more than on the other side. He was anxious to leave the hospital, but was persuaded to stay, more especially as he was suffering from a slight cough. Next day (August 1st) he had an attack described as difficulty of breathing; the pulse failed in a remarkable manner, the lips became white, and skin cold. He was ordered a saline draught, with antimony and digitalis. Auscultation detected nothing except bronchial rale at the top of the left lung, and increased pulsation at the apex of the heart, both sounds of which were distant and low.

He had similar attacks to the above two or three times in the next three days. On the 5th of Aug. he seemed extremely depressed; the pulse was small and quick; face very pale, with clammy perspiration; he complained occasionally of pain between the shoulders and in the right armpit, and seemed sinking. During the night he had another attack of syncope, in which he died.

At the *post mortem* examination the lungs were found congested; the bronchial tubes contained much thick mucus; their lining membrane was very vascular. There was a pouched condition of the upper part of the arch of the aorta, about as large as a walnut, between the origin of the innominate artery and the left carotid, whose origin was somewhat involved. This dilated part contained some rather firm coagulum. Besides this there was a large aneurismal pouch at the posterior and upper part of the thoracic aorta, communicating by an oval even edged orifice, of the size of a florin, with the aorta. This pouch was firmly adherent to the bodies of the upper five dorsal vertebrae, and bulged obliquely into the pleural sacs, on either side of the vertebral column. It contained much loose blood and fibrinous coagulum; and on opening it the anterior surfaces of the vertebrae were found to be much corroded. The whole length of the aorta was very rough with atheroma.

The large vessels given off from the aorta were quite natural, but the fasciae and other tissues about the root of the neck on the left side, and in the neighbourhood of the subclavian artery, were thickened and hardened, and the lymphatic glands enlarged.

ROYAL FREE HOSPITAL.

IRREDUCIBLE HERNIA OF THE STOMACH (?).

By WILLIAM BRINTON, M.D., F.R.C.P., Physician to the Hospital.

THE pathology of any given organ receives few better illustrations than from the rare instances which afford us the means of contrasting the results of injury with those of disease. In the case of the nervous system, the delicacy of its texture renders these two classes of lesion far more closely parallel to each other, than is the case with organs the structure of which better enables them to resist external violence. But even in those parts of the body in which the mechanical effects of injury are least definite and calculable, and in which the peculiar connexion of the sympathetic system of nerves with the cerebro-spinal centre adds a further element of obscurity to the diagnosis, by depriving us of that information which is elsewhere afforded by the phenomena of sensation and movement:—even here, careful comparison will often throw an invaluable light on the symptomatology of the particular organ or structure. Like the rude sketches of an artist of genius, the symp-

toms are all the more impressive from the fewness and boldness of the outlines by which they are indicated. To speak less metaphorically, the previous health of the patient often renders these indications of lesion unusually valuable, by excluding all those secondary phenomena—those refractions, as we may call them, of the primary effects of the mischief—which often colour or obscure the results of idiopathic disease. Practically, indeed, there can be little objection to our summing up the chief peculiarities of these cases of mechanical injury, in a diagnostic point of view, by the statement—that they represent the minimum of the various symptoms that betray the true diseases of the corresponding organ; and hence that their careful study may sometimes materially aid us in penetrating the obscurity of these visceral maladies.

Such observations especially apply to injuries of the various parts of the intestinal canal. And they may therefore suitably introduce a brief description of an interesting case which lately came under my notice; and which, at once diagnosed by me as being probably a hernia of the stomach, offers some interesting analogies to those cases of gastric ulcer I have already reported in this JOURNAL.

CASE. G. G., a tall robust coal-heaver, aged 45 years, was engaged in his usual avocation, about four hours after a tolerably hearty dinner, on the 29th of December, 1853. He was lifting a heavy sack of coals from the ground on to the back of his work-mate; and while thus stooping forward, some one startled him extremely by touching his back. The shock made him throw back his body by a sudden and violent extension of the spine; and at this very moment, he felt something snap in the pit of his stomach. At the same time there came on a pain in the epigastrium, and this pain soon acquired a sharp stabbing character. It was attended with nausea, but not with vomiting. Half an hour afterwards he had a stool, not, to his knowledge, an unnatural one.

The next morning, after taking what seems to have been an ordinary aperient medicine, prescribed for him by his medical attendant, he began to pass a considerable quantity of blood in his stools; and for about seven or eight days, none of the evacuations were altogether free from this admixture. On the second day after the injury, he twice passed more than a pint of nearly pure blood; but at the end of the week, the quantity of blood was not more than enough to give the stools a purplish black colour.

This hæmorrhage subsequently recurred twice: once six months, once nine months, after the injury. On both occasions it lasted about a week.

The pain and nausea quite unfitting him for work, he has at various times sought for medical relief. With this view he has been largely bled; and has also been cupped five times, and blistered five times, in the epigastrium. These depletions have, he believes, weakened him extremely. But they have in no degree alleviated his symptoms.

In December 1854, he became an out-patient of the Royal Free Hospital. On examining the belly, I found a small tumour, of the size of a cherry, just on the left side of the median line, about midway between the navel and the ensiform cartilage. The convexity of the tumour could be detected by the eye, but its size and shape could only be appreciated properly by careful manipulation. It was tough, rather elastic, quite unchangeable in size by any pressure; and its deepest portion or contracted neck was tightly and immovably embraced by the sheet of tendon that here covers in the *rectus abdominis* muscle, through an aperture on the inner side of which muscle it evidently protruded. As far as one could judge, it had none of the usual soft doughy feel of omentum.

He generally suffers from more or less pain in the neighbourhood of this tumour. The exact situation of this pain usually corresponds to an arched line (P, P, Fig.), the convex centre of which is at the ensiform cartilage (X), while its bases or extremities, below (**), tolerably coincide with the meeting point of the umbilical, hypogastric, lumbar, and iliac regions of the belly. The pain is referred to this situ-

ation even when it is immediately excited (as it readily is) by pressure or manipulation of the tumour. It is relieved by the recumbent posture; and is rendered much more severe by stooping, and by muscular exertion. Its worst paroxysms are accompanied by nausea and loss of appetite. During the twelve months it has been present, the patient has lost flesh, strength, and colour: indeed, his general health has seriously suffered; while the severe pain to which even moderate exercise gives rise quite precludes him from earning his living.

The accompanying diagram will perhaps assist the reader in appreciating the situation and relations of the tumour (T). The dotted lines indicate the relative position of the liver (L) and stomach (S), as mapped out by careful and repeated examinations. Such inquiries confirm the inference obviously deducible from the site of the tumour, by tracing towards it (almost into it) the relaxed and tympanitic stomach, in passing gradually from the left extremity towards its pyloric pouch.

The further history of the case is unsatisfactory enough. Supposing my diagnosis to be correct, it followed that the man was suffering from a small but irreducible hernia of some part of the stomach, completely disabling him for work, and inflicting on him continual suffering. Such circumstances seemed to me to suggest an operation for the relief and return of the impacted viscus. On explaining this to the patient, he declared himself willing to submit to any remedial measures of the kind that it might be thought advisable to adopt.

With that view, I sent him to Mr. Fergusson, my friend and former colleague, who, after examining the tumour, came to the same conclusion that I had done respecting its nature, and decided to operate. But, just before the time fixed for the operation, the patient's resolution failed him; and he absented himself altogether. A few weeks ago, I saw him again; according to his own statement, decidedly worse, and suffering from more pain than before.

REMARKS. Without venturing to regard the above diagnosis as more than a probability, we may at least assert, not only that it explains a larger number of the circumstances connected with the case than any second opinion, but also that these circumstances are scarcely compatible with any other diagnosis. The pain, the nausea, the aperture in the spongerosities, and the characters of the tumour,

decidedly indicate the protrusion of some mobile, flexible, and important portion of the contents of the belly:—some part, in short, of the digestive canal or its immediate attachments. But, to say nothing of the situation of the tumour, and its consistence under manipulation, the date and amount of the subsequent hemorrhage was such as any mere impaction of omentum could scarcely explain. The injury and the constriction might therefore be regarded as involving the tube itself.

That the portion of tube thus impacted was stomach, and not intestine, seemed deducible, not only from the situation of the tumour, and its relation to the traceable outline of the flaccid and mobile stomach, but also from the complete absence of all symptoms of obstruction of the canal. Had either small or large intestine been impacted in the form of such a tumour, even by but a portion of its calibre, or by its free edge, still the mechanical and physiological results of so much constriction would almost to a certainty have occluded the cavity of the tube; at any rate would pretty certainly have prevented the normal transmission of its contents. The pathology of hernia does indeed abundantly show, how easily the worst results of strangulation are brought about by no greater occlusion than such a partial impaction of the bowel would imply.

It is hardly necessary to add that neither depletion, counterirritation, nor mercury, could reasonably be expected to relieve the symptoms produced by such an accident as that above described. It is not, however, at all impossible that, at the date of its first protrusion, the tumour might have been reduced by proper manipulation: especially with the aid of chloroform, or the local application of ice. At the time I first saw it (nearly a year after the accident), any such proceeding was quite out of the question.

I regret that I cannot refer the readers of this JOURNAL to the three or four cases of this kind which I know to have been recorded by various observers. It is my impression that none of them quite resemble the above in the precision of their history and symptoms: and that one or two of them have been mere instances of painful epigastric protrusions, reducible by pressure, and relieved by proper bandaging. But all efforts to discover the references to these cases in my manuscript notes are, for the present moment, fruitless.

Brook Street, Grosvenor Square, Sept. 4th, 1886.

Hospital Gleanings.

VALUE OF THE OPHTHALMOSCOPE IN EYE DISEASES:
USE OF UPPER OR LOWER SECTION IN KERATONYXIS:
CURIOSITIES OF OPHTHALMIC PRACTICE.

THE recent introduction of the ophthalmoscope into the surgery of diseases of the organ of vision begins to be followed every year by better and better results. Too much at first was expected from it in eye diseases, and some amount of disappointment speedily followed its first introduction—the certain forerunner, very often, in science, of a better state of things. Every season now, since the labours of Jäger especially, the ophthalmoscope gives us at least negative results of very great value. A very long experience indeed and comparison of cases is necessary before the student or surgeon begins to see anything at all, but then it is very remarkable how easily various inflammatory or other morbid deposits are discovered in the retina, “Jacob's membrane”, lens, vitreous humour, etc. Chalk-like deposits in the “membrane of Jacob”, are, for instance, common in old people; inflammatory thickenings of retina, etc., are at once detected.

An interesting case of double cataract was recently operated on by Mr. BOWMAN, and is here appended, where the examination beforehand with the ophthalmoscope at least