ASSOCIATION MEDICAL JOURNAL:

BEING

THE JOURNAL OF THE BRITISH MEDICAL ASSOCIATION.

No. CXCIV.

LONDON: SATURDAY, SEPTEMBER 27, 1856.

Hospital Reports.

ST. GEORGE'S HOSPITAL.

RETENTION OF URINE FROM ENLARGED PROSTATE: PUNCTURE OF THE BLADDER FROM THE RECTUM.

By T. HOLMES, Esq., Surgical-Registrar to the Hospital.

The following case is the third of puncture of the bladder from the rectum which has occurred in St. George's Hospital during late years. The other two may be found reported by me in the *ASSOCIATION JOURNAL* for February 16th, p. 133, with some remarks on the controversy to which the frequent adoption of this operation at Guy's Hospital led. The case before us derives additional interest from having been performed for the relief of retention of urine arising from enlarged prostate. This condition is very generally said to contraindicate—in fact, to render impossible—the introduction of the trocar from the rectum. No doubt it does greatly increase the difficulty of the operation—especially when, as in this case, the patient is fat and the enlargement considerable; but even in the present case (I should think one of the most unfavourable in these respects for the operation), it was far more easy than either of the other methods of reaching the bladder would have been. The puncture above the pubes would have been an extremely formidable operation, from the vast depth of fat which intervened; and that from the perinæum hardly possible. It was necessary, however, that the man's bladder should be emptied somehow, as he was rapidly sinking from typhoid symptoms, and instruments had been tried over and over again without success. The relief which followed the operation was most marked; and the patient's condition while he retained the canula in the bladder, contrasted strikingly with the state from which he had been rescued, and into which he subsequently fell. This latter was a consequence apparently of effusion of urine into the cellular tissue between the bladder and rectum—"an accident", as Sir B. Brodie says, "too surely followed by the death of the patient." There is no reason, however, for supposing that this was a consequence of the operation; in fact, it will be seen by the notes of the case, that so long as the catheter could be passed easily, the patient's state continued to improve. Unfortunately, one day in drawing off the water, a false passage was made, or, which is perhaps more likely, an old one reopened. After this, it was impossible to pass a catheter without inducing fresh irritation; and he could not bear an instrument left in the bladder. His condition continued to grow worse; and, during the last few days of his life, he sank more rapidly, exhibiting those symptoms which Sir B. Brodie has described in cases where diffuse suppuration is found in the subperitoneal cellular tissue. It will be seen by the sketch of the *post mortem* examination, which I subjoin, that besides this subperitoneal inflammation, the peritoneum itself was perforated by ulceration.

CASE. K. K.—, an old man, probably about 70 years of age, a foreigner, was admitted into St. George's Hospital on Sept. 1st, under the care of Mr. Pollock. He had had difficulty in making water for a very long time; and for

the last nine days, the urine had been passed only in drops: for two days he had been incapable of passing any, but a small quantity had been dribbling away. The scrotum had been oedematous for about the same time. Mr. Pollock had seen him before his admission, as had another surgeon, and both had made unsuccessful attempts to introduce an instrument. This attempt was repeated, with the same result, on his arrival at the hospital. At that time, the bladder was much distended; there was great distress. The pulse was very weak; the tongue dry and brown. The whole cellular tissue of the penis and scrotum was very oedematous, and the prepuce so swollen that the orifice of the urethra could hardly be discovered; but there were no signs of extravasation of urine. It was determined to puncture the bladder without delay. The finger, introduced into the rectum, came upon a large hard tumour—the enlarged prostate; and beyond this, just within reach of the tip of the forefinger, when pressed in as far as possible, was a spot which felt less hard, and which was thought to be the projection of the distended bladder. The canula was thrust in towards this spot, but at first failed to reach the bladder. From the sensation communicated to the trocar, it was supposed that it had pierced the prostate. The next attempt was more successful; and fifty-six ounces of offensive urine were drawn off. Next day, he had rallied considerably—the pulse being stronger and the tongue moist and clean. The canula had become blocked up, and no water would flow through it: a gum catheter was therefore introduced through the tube of the canula, and passed readily into the bladder. Eighty ounces of more healthy urine were drawn off, to his very great relief.

From this date, he continued to improve steadily. The canula was left in the rectum for seven days. On the 7th, a full sized catheter was passed without any difficulty into the bladder. The tube was now withdrawn from the rectum, and the opening appeared to close at once, no more urine being observed to flow *per anum*. He could not, however, bear to retain an instrument in the bladder; and therefore it was necessary to pass one twice a day. This was done for two or three days without any untoward symptoms—when a gentleman was summoned to pass the catheter who had not previously had charge of the case. The instrument did not reach the bladder, but appeared to have reopened (for the force used was not thought sufficient to have made it) a false passage near the prostatic part of the urethra. From this time, it was difficult to pass the catheter without getting into this false passage, and a good deal of bleeding used to follow the introduction of the instrument. The urine became again very offensive, alkaline, and loaded with blood; he became gradually weaker, and was evidently sinking. Four days before his death he was found very much worse—with brown, dry tongue; pinched face; a tendency to low delirium; loss of appetite, etc.

He died nineteen days after the operation.

The contents of the pelvis were examined after death. There was found a good deal of suppuration in the subperitoneal cellular tissue, and a circumscribed abscess behind the upper end of the bladder. The coats of the latter presented the usual appearance of long continued inflammation. The opening in the rectum was nearly healed, but a probe could be forced through it. It led into the bladder exactly behind the projection of the prostate, which was greatly enlarged. This "middle lobe" formed a tumour in the cavity of the bladder, larger than a walnut.

The whole substance of the prostate gland was riddled by abscesses. Besides the opening made by the trocar, there were several other holes (with ulcerated margins) leading from the bladder into the cellular tissue between it and the rectum. There was also a false passage in the floor of the urethra, leading into the same space; and as the opening made by the trocar was pretty exactly in the middle line, an instrument passed with great readiness from the false passage through that opening into the bladder. Besides this, there was an ulcerated opening leading into the recto-vesical pouch of the peritoneum; and bloody fluid had been extravasated into the latter cavity. On the opposite side of the pouch, the gut was adhering to the bladder, near the ulcerated opening, and its coats showed traces of commencing ulceration. The whole of the body was enormously loaded with fat. No other organs were examined.

Hospital Gleanings.

BEST TREATMENT OF VARICOSE ULCERS AND VARICOSE VEINS: MODE OF FORMING CAUSTIC ISSUES: SUPERIORITY OF THE VIENNA PASTE, ETC.

THE treatment of varicose veins is a practical every day subject, inferior to none in interest for the hospital surgeon. Indeed, allied as this affection is, externally, to questions of practical surgery on the one hand, and depending for its causes on relations of the general venous system and general internal health on the other, a wide field of speculation is afforded as to treatment. In hospital practice, accordingly, as the practitioner may be more of a physiologist and anatomist, or more or less influenced by the ordinary routine of simple surgical treatment, ligature of veins, constitutional treatment, or means of blocking up these vessels, will be trusted to, previous to healing the varicose ulcer. A good deal of the danger of ligature of veins is found to originate in the fact that the vein is enlarged; and the ligature, when it should go deeper than the vein, will be found to have wounded or transixed the vein, and caused phlebitis.

Mr. Paget's experience every year assures him more and more of the superior value of the treatment of varicose veins by the caustic issue, or the extemporised mixture of lime and potash. There is no mode of treatment yet discovered, according to Mr. Paget, which is entirely free from risk; but, from pretty extensive inquiries in the hospital practice of such surgeons as Sir B. Brodie, Mr. Lawrence, Mr. Skey, various hospital surgeons in the provinces, and in France and Germany, etc., Mr. Paget is inclined to the impression that the treatment of varicose ulcers and varicose veins is most safely and most effectually conducted by means of local applications of caustic issues on the surface, which thicken the coats of the venous trunk.

A very well marked case of varicose ulcer of the leg and foot, which incapacitated the poor young man from employment and deprived his family of support, has been for some weeks in St. Bartholomew's, as well as a second case, where the varicose vessel gave way, but has again healed up. The former case is that of A. D—, a young and apparently otherwise active and healthy man aged 28. He has been all the present month (September) under treatment for a varicose condition of the veins of the lower extremity: his disease consists rather of a very troublesome ulcerated condition of the ankle and foot, which he cannot himself conceive to be dependent on the state of the veins. The history he gives of the case is not very defined, or is but a recapitulation of a set of abortive attempts to cure his ankle by ointments, lotions, salves, etc., each in

its turn recommended to him as the most specific thing in the world for an inward disorder of the constitution, but still only making his disease worse.

The form of caustic used by Mr. Paget consisted of potassa fusa $\mathfrak{D}$ ij, quick-lime $\mathfrak{z}$ j, separately in powders in small phials, and subsequently mixed with a glass or asbestos rod at the bedside, with spirits of wine, when used. A dozen or two of small pieces of common adhesive plaster are next obtained, each the size of a penny, a small circular hole being cut in each, of the size of a pea, or a silver threepence. One of these perforated pieces of plaster is placed over each projection or varicose enlargement of the vein, a very small quantity of the caustic paste is next applied with a bone spatula to the skin over the vein, in the circular hole left in the plaster: ten to twelve or fifteen minutes will be sufficient time for the issue to produce its effect, at the expiration of which period, the whole limb is carefully sponged with warm water, and all the plasters and caustic washed away. Little else was necessary in this case, as in the general class of these cases, the issue merely requiring simple dressing, while the ulcers in the ankle went on healing *pari passu*.

As a local application to the ulcer during the action on the varicose vein of the issue, any simple ointment or black wash is all that is required.

TREATMENT OF PURULENT AND OTHER BAD FORMS OF OPHTHALMIA: NEW FORM OF LOTION.

The interest attaching to eye cases is so great that, even at the risk of being deemed empirical, so that one may save an eye from destruction, we may here give a form of lotion in extensive use in hospitals, due to the practice, we believe, of Mr. WHARTON JONES—a lotion which is almost specific in various forms of purulent ophthalmia and chronic conjunctivitis. Mercury, perhaps it may be said, will rarely, if ever, be required internally in purulent ophthalmia: the utmost decision, cleanliness, and care, are, however, necessary in treating such cases with lotions, as well as sedulous attention to the condition of the *primæ viæ*. Some practitioners prescribe small doses of mercury so as to affect the system slightly; but as observed to his class recently by Mr. Wharton Jones, inflammations of mucous passages are not those that are ever much benefited by mercury. This active agent or leeches may restrain the submucous cellular effusion, and prevent swelling; but we should take care how we affect the bowels and various chylopoietic viscera, by pushing the use of the mineral to pyalism.

When chemosis is slight, and ulceration has already set in, mercury can do very little good. Tyrrell's plan of scarification, though not very strongly recommended by Mr. Dixon, will be found useful, especially in the lighter and more chronic forms of ophthalmia; and where we require a lotion in place of that usually made of nitrate of silver, the following, in an ounce bottle, is best, with or without leeches:

$\mathfrak{R}$ Cupri sulphatis gr. ij; vini opii $\mathfrak{z}$ j; aquæ dest. 3 vij. Fiat lotio.

It is applied freely with soft camel's hair brush three times a day.

ENLARGED BURSA OF THE KNEE INJECTED WITH IODINE.

There is a very practical case at present under the care of Mr. PAGET—a case of enlarged bursa of the knee, treated by injecting tincture of iodine into the bursal cavity. The lad is aged 18, and was admitted a month since (August 26th). Mr. Paget believes the iodine a better plan of treatment than issue, or anything more formidable. The swelling, which existed for seven months, has now disappeared.

PHAGEDÆNIC ULCERS AND THEIR TREATMENT: INUTILITY OF CHLORATE OF POTASH: VALUE OF OPIUM.

A form of disease, which is often met, in various aggravated shapes, in hospital practice, is the spreading phagedænic sore, not specifically related to syphilis, but