Published in final edited form as:

Phytochemistry. 2008 April ; 69(6): 1359–1371.

Fractionation and Characterization of Biologically-active Polysaccharides from *Artemisia tripartita*

Gang Xie, Igor A. Schepetkin, Daniel W. Siemsen, Liliya N. Kirpotina, James A. Wiley, and Mark T. Quinn*

Department of Veterinary Molecular Biology, Montana State University, Bozeman, MT 59717, USA

Abstract

The leaves of *Artemisia* species have been traditionally used for prevention and treatment of a number of diseases. In this study, five polysaccharide fractions (designated A-I to A-V) were isolated from the leaves of *Artemisia tripartita* Rydb. by the sequential use of hot-water extraction, ethanol precipitation, ultra-filtration, and chromatography. The homogeneity and average molecular weight of each fraction were determined by high performance size-exclusion chromatography. Sugar composition analysis revealed that *Artemisia* polysaccharides consisted primarily of xylose, glucose, arabinose, galactose, and galactosamine. Moreover, all fractions contained at least 3.4% sulfate, and fractions A-II through A-V contained an arabinogalactan type II structure. All fractions exhibited macrophage-activating activity, enhancing production of intracellular reactive oxygen species and release of nitric oxide, interleukin 6, interleukin 10, tumor necrosis factor α , and monocyte chemotactic protein-1. In addition, all fractions exhibited scavenging activity for reactive oxygen species generated enzymatically or produced extracellularly by human neutrophils. Finally, fractions A-I and A-V exhibited complement-fixing activity. Taken together, our results provide a molecular basis to explain at least part of the beneficial therapeutic effects of *Artemisia* extracts, and suggest the possibility of using *Artemisia* polysaccharides as an immunotherapeutic adjuvant.

Keywords

Artemisia tripartita Rydb.; Asteraceae; immunomodulator; polysaccharide; macrophage; neutrophil; complement-fixing activity; reactive oxygen species; cytokines

1. Introduction

Processes that remove invading microbes and harmful foreign or endogenous substances are essential to maintain normal development and homeostasis in multicellular organisms (Beutler, 2004). Phagocytic leukocytes play multiple roles in these immune processes, serving as a link between the innate and acquired immune systems and contributing to the inflammatory response, angiogenesis, and the promotion of wound healing (Beutler, 2004; Hoebe et al., 2004). Importantly, phagocytes can directly kill invading microorganisms and tumor cells, using both oxidative and non-oxidative mechanisms (Tosi, 2005). Thus, the development of

*Address for Correspondence: Dr. Mark T. Quinn, Veterinary Molecular Biology, Montana State University, Bozeman, MT 59717, Phone: 406-994-5721; Fax 406-994-4303, E-mail: mquinn@montana.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

novel therapeutics to nonspecifically augment macrophage immune responses represents an ideal strategy for enhancing defense against microbial infection (Finlay and Hancock, 2004).

A wide range of bioactive polysaccharides have been isolated from various medicinal plants, and these polysaccharides have been shown to possess immunomodulatory activity through their ability to modulate macrophage function [reviewed in (Schepetkin and Quinn, 2006)]. Indeed, botanical polysaccharides have been reported to increase macrophage cytotoxicity against tumor cells and microorganisms, activate phagocytosis, increase reactive oxygen species (ROS) and nitric oxide (NO) production, and enhance secretion of a variety of cytokines [reviewed in (Schepetkin and Quinn, 2006)]. Appropriate enhancement of these innate immune functions by bioactive compounds can then lead to improved host defense responsiveness (Finlay and Hancock, 2004). Moreover, most plant-derived polysaccharides are relatively non-toxic and do not cause severe side effects, which is a major problem associated with immunomodulatory bacterial polysaccharides and synthetic compounds (Finlay and Hancock, 2004). Thus, plant polysaccharides represent ideal candidates for therapeutics with immunomodulatory, anti-tumor, and wound-healing action.

The genus *Artemisia* belongs to the family Compositae (Asteraceae) and has been used in folk medicine to treat a variety of diseases, such as hepatitis (Hong et al., 2004), fever and malaria (Klayman, 1985), rheumatoid arthritis (Wang et al., 2005), and asthma (Kim et al., 2006). In the western United States, *Artemisia tripartita* (three-tip sagebrush) is a native species that has been used in the treatment of colds, sore throats, tonsillitis, headaches and wounds by Native Americans (Moerman, 1998); however, the biologically active components in *A. tripartita* extracts are unknown. Low-molecular-weight compounds (e.g., lignans and sesquiterpene lactones) have been identified and isolated from *A. tripartita* (Kelsey and Shafizadeh, 1979), and similar compounds from other *Artemisia* species have been shown to have anti-fungal properties (Tan et al., 1998). Likewise, polysaccharides from various *Artemisia* species have been reported to exhibit therapeutic activities (Hayakawa et al., 1995; Hwang et al., 2003; Lee et al., 2003); however, potential immunomodulatory properties of *A. tripartita* polysaccharides have not been evaluated.

In the present study, we isolated five polysaccharide fractions from aqueous extracts of the leaves of *A. tripartita*. Analysis of these polysaccharide fractions showed that they had potent phagocyte immunomodulatory activity, as well as ROS scavenging and complement-fixing activity. Thus, the immunomodulatory activities of *Artemisia* polysaccharides likely contribute to the known therapeutic effects of *Artemisia* extracts.

2. Results and Discussion

2.1. Preparation and characterization of *Artemisia* polysaccharides

Acidic polysaccharides obtained from *Artemisia* extract were fractionated by preparative size-exclusion chromatography to obtain five main fractions, which were selected based on total carbohydrate elution profile (designated as A-I, A-II, A-III, A-IV and A-V) (Fig. 1A). Each of the five fractions contained <2.4% protein (Table 1). Analysis of the fractions using the Yariv test showed that fraction A-I lacked arabinogalactan; whereas, all other fractions tested positive for the presence of type II arabinogalactan (Table 1). Type II arabinogalactans have a β -(1,3)-linked galactan backbone with side chains containing arabinose and galactose residues and have been reported to possess a variety of biological activities (Paulsen, 2001). All five fractions also contained at least 3.4% sulfate (Table 1), suggesting the possibility that *Artemisia* polysaccharides are also sulfated; however, further purification and analysis of the individual polysaccharides will be necessary to verify this issue. Note that the presence of sulfate groups has been reported to play a role in polysaccharide biological activity (Melo et al., 2002; Ghosh et al., 2004). For example, Schaeffer and Krylov (Schaeffer and Krylov, 2000) reported that

the presence of sulfate groups was necessary for anti-HIV activity of extracts from algae, and that potency increased with an increased degree of sulfation. Thus, it will be important in future studies to determine the role of sulfation in *Artemisia* polysaccharide biological activity.

Analytical high performance size-exclusion chromatography (HP-SEC) of the individual fractions showed that each fraction was represented by a broad, symmetrical peak on the chromatograms (Fig. 1B). Average molecular weights of the fractions were determined to be 355, 251, 126, 78, and 49 kDa for fractions A-I through A-V, respectively (Table 1).

Analysis of sugar composition of the *Artemisia* polysaccharide fractions showed that these polysaccharides consisted primarily of xylose, glucose, arabinose, galactose, and galactosamine, which accounted for the majority of monosaccharides present (Table 2). Small amounts of galacturonic acid, glucuronic acid, mannose, lyxose, and glucosamine were also detected in the fractions (Table 2). In polysaccharide fractions A-II through A-V, xylose was the dominant monosaccharide and represented >50 mol% of the total sugars in these fractions. In contrast, fraction A-I had a much higher glucose content (29.5 mol%), making it the predominant sugar in fraction A-I (Table 2). In comparison to *A. tripartita* polysaccharides, mannose was the predominant sugar in polysaccharides from *Artemisia sphaerocephala* (Zhang et al., 2006)], suggesting there are differences in polysaccharide structure between *Artemisia* species, which may confer distinct biological properties.

Very-high-field (600 MHz) ^1H NMR was used to characterize structure of the native *Artemisia* polysaccharides (Fig. 2). The spectra of fractions A-II, A-III, A-IV, and A-V were similar to each other, suggesting a common backbone structure, and resembled the spectra of native arabinogalactans isolated from other plant sources, which is consistent with the results of the Yariv test (Gane et al., 1995; Dong and Fang, 2001; Schepetkin et al., 2005). The chemical shifts of anomeric protons were evaluated according to data previously reported for the sugar composition of the arabino-3,6-galactans (type II) (Gane et al., 1995; Dong and Fang, 2001). The signals present at 3.40–4.38 and 4.95–5.17 ppm were consistent with the presence of β -D-galactopyranose (β -D-Gal p) and α -arabinofuranose (L-Ara f) residues, respectively (Gane et al., 1995). In comparison, a very weak signal was detected at these chemical shifts in the spectrum of fraction A-I, confirming our Yariv test that also showed the absence of arabinogalactan in this fraction. Fractions A-II to A-V also contained much stronger signals for *N*- and *O*-acetyl (1.97–2.08 ppm) and methyl groups (1.13–1.20 ppm). A paired doublet at 2.43–2.54 ppm was present in the spectra of all 5 fractions; however, we subsequently found this signal was due to the presence of sodium citrate bound to the polysaccharides. In all *Artemisia* fractions, signals corresponding to aromatic protons (5.45–6.25 ppm) were absent or very weak, suggesting the lack of aromatic groups or their presence in very low molar concentrations in the large polysaccharide structures.

To directly evaluate possible lipopolysaccharide (LPS) contamination in our samples, we analyzed the *Artemisia* fractions for binding to rhodamine-labeled S3 peptide (Rh-S3), as described previously (Schepetkin et al., 2005). Analysis of the samples by HP-SEC showed free Rh-S3 was eluted as a single fluorescent peak with retention time of 25 min; whereas, addition of 5 ng LPS to Rh-S3 resulted in the appearance of a new fluorescent peak at 14 min, which represents Rh-S3 bound to LPS (Supplemental Fig. S1). In contrast, no new fluorescence peaks were observed after addition of the individual *Artemisia* polysaccharide fractions, even at 1000-fold higher concentrations of the fractions (Supplemental Fig. S1). Thus, these data directly demonstrate that LPS contamination was not present in the *Artemisia* fractions and, therefore, was not responsible for their immunomodulatory activity.

2.2. Effect of *Artemisia* polysaccharides on macrophage NO and ROS production

A minimum amount of NO was produced when murine J774.A1 macrophages were incubated with medium alone, whereas treatment of these cells with polysaccharide fractions resulted in a dose-dependent increase in NO production (Fig. 3A). Fractions A-I and A-V were the most potent inducers of macrophage NO production, enhancing the response >16-fold at concentrations of 800 µg/ml compared to the negative control (medium alone). In comparison, NO production induced by fractions A-I and A-V was a little over 50% of that induced by 50 ng/ml lipopolysaccharide (LPS).

In the absence of any treatment, murine macrophages generated very low levels of ROS; whereas, a dose-dependent enhancement of ROS production was observed in macrophages treated with 100–800 µg/ml doses of the polysaccharide fractions (Fig. 3B and Supplemental Fig. S2), with fractions A-I and A-V being the most active fractions. Thus, the individual polysaccharide fractions showed a similar pattern with respect to their ability to induce NO and ROS production by murine J774.A1 macrophages. If we consider activity in terms of polysaccharide molar concentration, fraction A-I was still the most potent inducer of NO and ROS production (relative potency was A-I>A-II>A-III>A-IV≈A-V); whereas, fractions A-IV and A-V were the least active.

Macrophages play critical roles in host defense, including phagocytosis of pathogens and apoptotic cells, production of cytokines, and proteolytic processing and presentation of foreign antigens [reviewed in (Hume, 2006)]. Thus, the identification of agents that can modulate macrophages is of significant interest. Indeed, a variety of plant polysaccharides have been reported to exhibit beneficial pharmacological effects via their ability to modulate macrophage function [reviewed in (Schepetkin and Quinn, 2006)]. In this study, *Artemisia* polysaccharide fractions exhibited macrophage immunomodulatory properties, inducing NO and ROS production. These reactive oxidants play key roles host defense and various physiological processes and are also involved in the regulation of apoptosis and immune homeostasis [reviewed in (Serhan and Savill, 2005;Quinn et al., 2006)].

Although these functional assays suggested *Artemisia* polysaccharides were relatively non-toxic, we evaluated the potential cytotoxic effect of *Artemisia* polysaccharides to determine if the results might be influenced by background toxicity. Using an assay that measures ATP from viable cells, we found that none of the fractions significantly affected proliferation/viability of J774.A1 cells over the entire concentration range of *Artemisia* polysaccharides (100–800 µg/ml), confirming that these polysaccharides were not toxic to these cells (data not shown).

2.3. Effect of *Artemisia* polysaccharides on macrophage cytokine production

Many immunomodulatory compounds modulate cytokine and/or chemokine production. Thus, we analyzed the effects of *Artemisia* polysaccharide fractions on this response. As shown in Fig. 4, each of the five *Artemisia* polysaccharide fractions stimulated macrophage production of both pro-inflammatory and anti-inflammatory cytokines. Among the pro-inflammatory cytokines induced, interleukin 6 (IL-6) is one of the most important mediators of fever and the acute-phase response (Gabay, 2006). Tumor necrosis factor α (TNF- α) also plays an important role in inflammation and can act on monocytes and macrophages in an autocrine manner to enhance various functional responses and induce the expression of additional immunoregulatory and inflammatory mediators (Baugh and Bucala, 2001). Production of the inflammatory chemokine, monocyte chemoattractant protein 1 (MCP-1), was also induced by *Artemisia* polysaccharides at least 1.4-fold, as compared to untreated controls. Indeed, the enhancement of MCP-1 production was comparable to that obtained with LPS, which is well known as a strong cytokine inducer (Fig. 4). This chemokine is involved in the recruitment of

monocytes during acute and chronic inflammation (Conti and DiGioacchino, 2001). Along with these inflammatory mediators, the anti-inflammatory cytokine, IL-10, was also induced by *Artemisia* polysaccharides. IL-10 has been reported to inhibit cytokine synthesis and deactivate macrophages, thereby preventing or reducing potential detrimental effects from excessive macrophage activation during inflammation (Williams et al., 2004). Thus, the modulation of macrophage cytokine and chemokine production likely contributes to part of the therapeutic effects of *Artemisia* extracts. Note that previous studies have shown many plant-derived polysaccharides can induce the production of a range of cytokines [reviewed in (Schepetkin and Quinn, 2006)]; however, only polysaccharides from *Juniperus scopolorum* (Schepetkin et al., 2005) and *Artemisia*, as shown here, were found to induce IL-10 and MCP-1 production.

2.4. Effect of *Artemisia* polysaccharides on the complement system

Many plant-derived polysaccharides, including some arabinogalactans, have complement fixing activity (Nergard et al., 2004). Therefore, *Artemisia* polysaccharide fractions were tested for their ability to fix complement, as compared to heparin, a known complement-fixing agent. As shown in Fig. 5, addition of each of the 5 fractions resulted in a dose-dependent inhibition of erythrocyte hemolysis, indicating that the isolated *Artemisia* polysaccharides exhibited complement fixing activity. Fractions A-I and A-V showed the highest complement-fixing activity, whereas fraction A-IV was the least active (Fig. 5). All fractions inhibited hemolysis with $IC_{50} \geq 800 \mu\text{g/ml}$, which is similar to the activity of polysaccharides isolated from *Brassica oleracea* (Samuelsen et al., 2007). However, it should be noted that polysaccharides isolated from other plants (e.g., *Biophytum petersianum* Klotzsch and *Veronia kotschyana*) exhibit higher complement fixing activity with $IC_{50} < 100 \mu\text{g/ml}$ (Nergard et al., 2006; Inngjerdigen et al., 2006). Thus, there appears to be quite a bit of variability in level of complement-fixing activity between polysaccharides extracted from different species. Nevertheless, our results suggest the possibility that complement-fixing activity could contribute to the immunomodulatory properties of *Artemisia* polysaccharides. Indeed, removal of complement by polysaccharide fixation has been proposed to be a potential therapeutic strategy for treating inflammatory diseases (Nergard et al., 2004), and various reports have shown that polysaccharides from different medicinal plants have anti-complement activity [reviewed in (Schepetkin and Quinn, 2006)].

2.5. Effect of *Artemisia* polysaccharides on neutrophil ROS production

To determine whether neutrophils also responded to *Artemisia* polysaccharides, we evaluated their effects on murine bone marrow neutrophils. In the absence of any treatment, murine neutrophils spontaneously generated significant levels of ROS (Fig. 6A). Surprisingly, all five polysaccharide fractions appeared to dose-dependently inhibit spontaneous ROS production by murine neutrophils, with fraction A-I being the most inhibitory (Fig. 6B). To evaluate whether this response was specific or not to murine neutrophils, we performed similar studies using human neutrophils and found that *Artemisia* polysaccharides also seemed to inhibit spontaneous ROS production by unstimulated human neutrophils (Fig. 6C,6D). Treatment of human neutrophils with fractions A-I, A-II, A-IV and A-V resulted in dose-dependent loss of the ROS signal. Although fraction A-III was much less potent, it still inhibited ROS produced by human neutrophils at concentrations $> 200 \mu\text{g/ml}$.

Since the inhibition of ROS by neutrophils treated with *Artemisia* polysaccharides was completely opposite to the activation observed in murine macrophages (see Fig. 3B), we considered the possibility that *Artemisia* polysaccharides might have ROS scavenging activity and that these cell types could be generating ROS in different cellular locations (i.e., intracellular vs. extracellular). We first analyzed the effect of *Artemisia* polysaccharides on intracellular ROS production in human neutrophils phagocytosing serum-opsonized zymosan.

Phagocytosis of zymosan particles triggers intracellular ROS production, which can be specifically detected by the inclusion of superoxide dismutase (SOD) in the assay buffer to remove any extracellular ROS. In this system, Artemisia polysaccharides did not affect intracellular ROS production (Fig. 6E,6F), suggesting the polysaccharides only had access to extracellular ROS.

To evaluate ROS scavenging activity of the polysaccharide fractions in the J774.A1 macrophage system, cells were pretreated with 50 ng/ml LPS for 20 hr to elicit maximal ROS production, and chemiluminescence was measured immediately following treatment of the cells with 200 nM phorbol-12-myristate-13-acetate (PMA) with and without the indicated concentrations of polysaccharides. As shown in Fig. 7, none of the fractions reduced the ROS signal, suggesting ROS were being generated intracellularly and were, therefore, inaccessible to the polysaccharides located outside of the cell. To evaluate this possibility, we utilized cell-permeable and cell-impermeable ROS scavengers. Mannitol, a cell-permeable $\text{OH}^\bullet$ scavenger (Li et al., 1996), significantly reduced the ROS response in this system; whereas, SOD, a cell-impermeable $\text{O}_2^{\bullet-}$ scavenger (Dahlgren and Karlsson, 1999), and dextran, a cell-impermeable $\text{OH}^\bullet$ scavenger (Li et al., 1996), had no ROS scavenging effect (Fig. 7). Thus, we conclude that J774.A1 macrophages primarily produce intracellular ROS, which would be inaccessible to scavenging by cell-impermeable Artemisia polysaccharides.

To directly confirm ROS scavenging activity of Artemisia polysaccharides, we evaluated their ability to scavenge enzymatically-generated $\text{O}_2^{\bullet-}/\text{H}_2\text{O}_2$ (xanthine/xanthine oxidase system) and $\text{OH}^\bullet$ [H_2O_2 /horseradish peroxidase (HRP) system]. Artemisia polysaccharide fractions A-II to A-V exhibited ROS scavenging activity in the xanthine/xanthine oxidase system (Fig. 8A). As a positive control, SOD also inhibited most of the chemiluminescence signal. Addition of Artemisia polysaccharide fractions to an $\text{OH}^\bullet$ -generating system (H_2O_2 /HRP) also resulted in dose-dependent scavenging of $\text{OH}^\bullet$, which was similar to that of the positive controls, mannitol and dextran (Fig. 8B). Thus, these data support the conclusion that the apparent inhibition of neutrophil ROS production by Artemisia polysaccharides was actually due, at least in part, to direct scavenging of neutrophil-generated ROS. Indeed, when neutrophils treated with Artemisia polysaccharides were washed to remove the polysaccharides prior to measuring ROS, we found that fractions A-II and A-III actually enhanced human neutrophil ROS production >6-fold and >8-fold, respectively; whereas, fractions A-I, A-IV, and A-V had little effect on this response (Fig. 9A). In comparison, none of the polysaccharide fractions stimulated murine neutrophil ROS production (Fig. 9B). The reason for the differences in responses between human and murine neutrophils is not clear; however, this may simply reflect differences between bone marrow and peripheral blood neutrophils.

The ability of Artemisia polysaccharides to scavenge ROS may play an important role in the therapeutic properties of Artemisia extracts. Various other botanical polysaccharides have also been reported to possess ROS scavenging activity. For example, protein-bound polysaccharide from the mycelium of *Coriolus versicolor* (Sakagami et al., 1992) and polysaccharide extracts from other mushroom species (Liu et al., 1997) were found to possess ROS-scavenging activity. Moreover, dextran was reported to function as an $\text{OH}^\bullet$ scavenger to protect proteins from metal-catalyzed oxidation (Li et al., 1996). Thus, our data are consistent with the putative role of various botanical polysaccharides as ROS scavengers. It should be noted, however, that the large Artemisia polysaccharides were only effective scavengers of extracellular ROS, and ROS generated intracellularly were not accessible.

Neutrophils are principal effectors of the initial host response to injury or infection and constitute a significant threat to invading bacterial pathogens (Tosi, 2005). Thus, the ability to regulate neutrophil function represents an important immunomodulatory property of Artemisia polysaccharides. A number of other plant-derived polysaccharides have been reported to

stimulate neutrophil functional activity. For example, chitosan, a polysaccharide of D-glucosamine, accelerates the wound healing process by enhancing neutrophil phagocytic activity as well as ROS production (Morimoto et al., 2001). Likewise, polysaccharides from *Acacia cyanophylla* stimulated ROS production by murine neutrophils (El Abbouyi et al., 2004), and polysaccharides purified from *Ganoderma lucidum* enhanced human neutrophil phagocytosis and chemotaxis (Hsu et al., 2003).

3. Conclusion

Although Artemisia has been widely used in traditional remedies, little is known regarding the active component(s) responsible for its therapeutic properties. While previous studies indicated that some of the biological properties of Artemisia were due to the presence of low-molecular weight compounds, such as lignans and sesquiterpene lactones, we have demonstrated here that polysaccharides of Artemisia have potent immunomodulatory properties. We isolated five sulfated polysaccharide fractions from *A. tripartita* and demonstrated that these polysaccharides exhibited immunomodulatory and anti-inflammatory activities, including modulation of macrophage and neutrophil functions, scavenging of extracellular ROS, and complement-fixation. Collectively, these activities could lead to enhanced phagocyte microbicidal responses, while protecting host tissues from excessive oxidant injury and enhancing wound-healing.

4. Experimental

4.1. Reagents

β -glucosyl Yariv reagent [1,3,5-tri-(4- β -D-glucosopyranosyloxyphenyl-azo)-2,4,6-trihydroxybenzene] was purchased from Biosupplies Australia (Parkville, Australia). Gum arabic was purchased from Fluka BioChemica (Buchs, Switzerland). 8-amino-5-chloro-7-phenylpyridol[3,4-d]pyridazine-1,4-(2H,3H)dione (L-012) was purchased from Wako Chemicals (Richmond, VA). The cytometric bead array kit was purchased from BD Biosciences Pharmingen (San Diego, CA). DEAE-cellulose, Sepharose-6B, dextran from *Leuconostoc mesenteroides*, N-(1-naphthyl)ethylenediamine, sulfanilamide, horseradish peroxidase (HRP), Histopaque 1077, zymosan A from *Saccharomyces cerevisiae*, LPS from *Escherichia coli* K-235, xanthine oxidase, xanthine, SOD from bovine erythrocytes, PMA, antibody sensitized sheep erythrocytes, sodium nitrite (NaNO_2), and gelatin veronal buffer (GVB) were purchased from Sigma Chemical Co. (St. Louis, MO). Pullulan standards were purchased from Phenomenex (Torrance, CA).

4.2. Fractionation of polysaccharides

Leaves of *A. tripartita* were collected in the mountain areas of Montana and were taxonomically confirmed. Dried and ground leaves (1.2 kg) were extracted with 10 L boiling distilled H_2O for 1 hr, and the aqueous extract was centrifuged at $2,000 \times g$ for 15 min. A four-fold volume of ethanol was added to the supernatant to precipitate the polysaccharides overnight at 4°C . The precipitate was pelleted by centrifugation, re-dissolved in distilled H_2O , sonicated for 10 min, and centrifuged at $80,000 \times g$ for 1 hr. The supernatant was then filtered through a 0.22 μm filter and concentrated in an Amicon concentrator with a 5 kDa PM5 membrane to obtain a crude extract (yield of 1.2% by weight).

The crude polysaccharide extract was further purified using ion-exchange chromatography on a DEAE-cellulose column equilibrated with 0.05 M Tris-HCl buffer (pH 8.0). The column was washed with equilibration buffer, and bound material was eluted with equilibration buffer containing 2 M NaCl. The eluate was concentrated to obtain the crude acid polysaccharide extract, which was then fractionated by size-exclusion chromatography on a Sepharose-6B

column (2.5×95 cm) equilibrated with 0.01 M sodium citrate buffer (pH 7.0) containing 0.15 M NaCl and eluted with the same buffer at a flow rate of 21 ml/hr. The carbohydrate elution profile was determined by the phenol-H₂SO₄ method (Dubois et al., 1956), modified to a microplate format, and absorbance was measured at 488 nm using a SpectraMax Plus microplate reader (Molecular Devices, Palo Alto, CA). The relevant fractions were pooled and concentrated. For analysis of biological activity, the fractions were diluted in Hanks balanced salt solution (HBSS) to a concentration of 5 mg/ml and filtered through sterile 0.22 µm filters.

4.3. Characterization of polysaccharide fractions

Samples were analyzed for protein content using a modified Lowry assay (Sigma) with bovine serum albumin as a standard. Sulfate determination was carried out by the sodium rhodizonate method (Terho and Hartiala, 1971). For nuclear magnetic resonance (NMR) analysis, samples (5 mg) were dissolved in deuterium oxide (0.5 ml), and ¹H NMR spectra were recorded on a Bruker DRX-600 spectrometer (Bruker BioSpin, Billerica, MA) at 20°C using 3-(trimethylsilyl)-propionic 2,2,3,3,-d₄ acid sodium salt as an internal reference.

The presence of arabinogalactan in the samples was detected by single radial gel diffusion in a 1% agarose gel containing 100 µg/ml β-glucosyl Yariv reagent, which selectively interacts with and precipitates compounds containing type II arabinogalactan structures (van Holst and Clarke, 1985). Four µl of polysaccharide samples (5 mg/ml) were loaded into the wells, and the samples were incubated at room temperature for 24 hr in a humid atmosphere. A positive reaction was identified by a reddish circle around the well, and arabic gum (4 mg/ml) served as a positive control.

The homogeneity and average molecular weight of the polysaccharide fractions were determined by HP-SEC using a Shimadzu Class VP HPLC and Shodex OHpak SB-804 HQ column (8 mm × 300 mm) eluted with 50 mM sodium citrate buffer, pH 7.5, containing 0.15 M NaCl and 0.01% NaN₃ at a flow rate of 0.3 ml/min. Peaks were detected using a refractive index detector (RID-10A; Shimadzu, Torrance, CA). The molecular weights of the polysaccharide fractions were estimated by comparison with retention times of pullulan standards (P-800, 400, 200, 100, 50, 20, and 10).

Samples were tested for possible LPS contamination using a rhodamine-labeled peptide (EndoDtec-F, BioDtech, Inc.) derived from the sushi 3 domain of horseshoe crab Factor C (Tan et al., 2000). This peptide binds to LPS with high affinity and specificity (Tan et al., 2000), and we showed previously that EndoDtec-F binding provided a sensitive measurement of LPS in plant-derived polysaccharide samples (Schepetkin et al., 2005). EndoDtec-F (250 ng) was mixed with 5 ng LPS or 5 µg polysaccharide fractions, and 10 µl samples were analyzed by HP-SEC as described above, but at a flow rate of 0.5 ml/min. The elution was monitored using tandem refractive index (RID-10 A) and fluorescence detectors (RF-10 A, Shimadzu). Fluorescence peaks were detected using excitation and emission wavelengths of 550 and 580 nm, respectively.

4.4. Monosaccharide composition analysis

Buffer salts were removed from the polysaccharide samples by repeated (6×) concentration in an Amicon concentrator (5 kDa cut-off PM5 membrane) and dilution with a 10- fold volume of distilled H₂O. The final concentrates and background blanks (distilled H₂O) were lyophilized and submitted for analysis. Monosaccharide analysis was performed by the Oklahoma Center for Glycobiology Analytical Core Lab (Oklahoma City, OK). Briefly, polysaccharide samples or background blanks were subjected to methanolysis (methanolic 2 M HCl, 16 hr, 80°C), followed by acid hydrolysis (2 M trifluoroacetic acid, 4 hr, 100°C), and the resulting monosaccharide mixtures were analyzed by high-performance anion-exchange

chromatography with pulsed amperometric detection (HPAEC-PAD) on a Dionex DX-600 HPAEC system equipped with an ED50 detector (Dionex Corporation, Sunnyvale, CA). The samples were separated on a Dionex CarboPac PA20 column eluted isocratically with 12 mM NaOH at a flow rate of 1 ml/min at 22°C. For analysis of uronic acids, the column was eluted with 10 mM NaOH for 20 min, followed by a gradient of 100 mM NaOH/150 mM sodium acetate (0–100% in 45 min). Background signals were subtracted from all samples, and individual components were quantified based on electrochemical detection relative to known standards (De Ruiter et al., 1992; Hardy and Townsend, 1994).

4.5. Cell culture

Murine macrophage J774.A1 cells were cultured in Dulbecco's Modified Eagle's Medium (Invitrogen) supplemented with 10% (v/v) heat-inactivated fetal bovine serum, 100 µg/ml streptomycin, and 100 U/ml penicillin at 37°C in a humidified atmosphere containing 5% CO₂. Cells were grown to confluence in sterile tissue culture flasks and gently detached by scraping. Cell number and viability were assessed microscopically using trypan blue exclusion.

4.6 Isolation of human and murine neutrophils

Human neutrophils were purified in accordance with a protocol approved by the Institutional Review Board at Montana State University. Briefly, these cells were isolated from peripheral blood of healthy donors using dextran sedimentation followed by Histopaque 1077 gradient separation and hypotonic lysis of erythrocytes, as described previously (Gauss et al., 2006). Isolated neutrophils were washed and resuspended in HBSS without Ca²⁺ and Mg²⁺. Cell viability was determined by using the trypan blue exclusion test (>98%), and the purity of neutrophils was determined by light microscopy (>95%).

Murine neutrophils were isolated and purified from bone marrow following the method described previously (Schepetkin et al., 2007). All animal use was conducted in accordance with a protocol approved by the Institutional Animal Care and Use Committee at Montana State University. Briefly, leukocytes were flushed from tibias and femurs of BALB/c mice with HBSS containing 0.1% bovine serum albumin and 1% glucose, filtered through a 70 µm nylon cell strainer (Becton Dickson, Franklin lakes, NJ) to remove cell clumps and bone particles, and pelleted by centrifugation. The bone marrow leukocyte pellets were resuspended in 3 ml of 45% Percoll solution and layered on top of a Percoll gradient consisting of 2 ml each of 50, 55, 62, and 81% Percoll solutions in a conical 15-ml polypropylene tube, and the gradient was centrifuged at 1,600×g for 30 min at 10°C. The neutrophil band located between 61 and 81% Percoll was collected, washed, layered on top of Histopaque 1119, and centrifuged at 1,600×g for 30 min at 10°C to remove contaminating erythrocytes. The purified neutrophils were collected, washed, and resuspended in HBSS without Ca²⁺ and Mg²⁺.

4.7. Measurement of NO production

J774.A1 cells were plated at a density of 1.5×10^5 cells/well in a final volume of 200 µl in 96-well flat-bottomed tissue culture plates and incubated in medium alone or medium containing various concentrations of polysaccharide fractions or *E. coli* LPS as a positive control. Cells were incubated at 37°C in the presence of 5% CO₂ for 24 hr., and 100 µl of the cell culture supernatants were removed and analyzed for NO using a colorimetric method with NaNO₂ as a standard, as described previously (Schepetkin et al., 2005). Briefly, supernatants were mixed with an equal volume of Griess reagent, which was prepared by mixing one part of 0.1% (w/v) N-(1-naphthyl)ethylenediamine with one part of 1% (w/v) sulfanilamide in 5% phosphoric acid. After 20 min, absorbance was measured at 540 nm using a SpectraMax Plus microplate reader.

4.8. Analysis of phagocyte ROS production

Macrophage ROS production was analyzed using the chemiluminescent probe, L-012, which is highly sensitive for ROS generated in biologically complex systems (Imada et al., 1999; Daiber et al., 2004). Murine J774.A1 macrophages (1.5×10^5 cells/well) were incubated with various concentrations of polysaccharide fractions or positive control LPS for 24 hr. After incubation, 100 μ l of culture supernatant (taken for NO or cytokine analysis) was replaced by an equal volume of HBSS supplemented with 25 μ M L-012 and 5 μ g/ml horseradish peroxidase, as described previously (Schepetkin et al., 2007). The reaction was monitored on a Fluoroskan Ascent FL microtiter plate reader (ThermoElectron, Milford, MA) at 37°C. Chemiluminescence was measured every 2 min for 2 hr and is expressed as the integrated response over this time (arbitrary units).

To evaluate *Artemisia* polysaccharide effects on neutrophil ROS production, human neutrophils (10^5 cells/well) or murine neutrophils (5×10^4 cells/well) were added to microplate wells containing the indicated concentrations of polysaccharide fractions plus 25 μ M L-012, and chemiluminescence was monitored at 37°C for 1 hr, as described above. To evaluate effects of polysaccharides on intracellular ROS production, human neutrophils were added to microplate wells containing serum-opsonized zymosan particles (100 μ g/ml), 25 μ M L-012, and 50 U/ml SOD, and chemiluminescence was monitored at 37°C for 1 hr (Schepetkin et al., 2006). Serumopsonized zymosan was prepared by incubating zymosan A with human serum for 45 min at 25°C. After centrifugation (800 $\times$ g for 10 min), the zymosan particles were washed and resuspended in HBSS (without Ca^{2+} and Mg^{2+}).

To evaluate polysaccharide effects on ROS production after polysaccharide removal, human or murine neutrophils were pre-incubated with polysaccharides fractions for 15 min at 25°C. After centrifugation of the samples at 82 $\times$ g for 5 min to pellet the cells, the media was aspirated, and the cells were washed with HBSS. The washed cells were placed in L-012 assay buffer, and chemiluminescence was monitored at 37°C for 1 hr.

4.9. Cytokine determination

J774.A1 cells were incubated at a density of 1.5×10^5 cells/well in final volume of 200 μ l for 24 h with or without *Artemisia* polysaccharide fractions or LPS. A cytokine cytometric bead array was used to simultaneously detect interleukins (IL)-6 and -10, tumor necrosis factor (TNF)- α , and monocyte chemoattractant protein (MCP)-1. Standard curves (ranging from 0 to 5000 pg/ml) for the assay system were obtained from dilutions of the mixed pool of all four cytokines that was provided with the kit. The concentration for each cytokine in the cell supernatants was determined by extrapolation from the corresponding standard curve using the CBA Software (BD Biosciences Pharmingen).

4.10. Complement fixation assay

The complement fixation assay was performed as previously described by Diallo et al. (Diallo et al., 2001). Antibody sensitized sheep erythrocytes were washed three times with GVB containing 0.5 mM Mg^{2+} and 0.15 mM Ca^{2+} (GVB^{2+}) before use. The erythrocytes were resuspended in GVB^{2+} at a concentration of 2×10^8 cells/ml, and human serum was diluted with GVB^{2+} to a concentration giving about 50% hemolysis. Triplicate samples containing 50 μ l of each serially-diluted polysaccharide fraction were mixed with 50 μ l diluted serum and added to microplate wells and incubated at 37°C. After 30 min, the sensitized sheep erythrocytes (50 μ l) were added to each well, and the samples were incubated for an additional 30 min at 37°C. After centrifugation (900 $\times$ g for 5 min), 50 μ l of the supernatants was mixed with 200 μ l distilled H_2O in flat-bottom microplates, and absorbance was measured at 405 nm. 100% lysis was obtained by adding distilled H_2O to sensitized sheep erythrocytes. Samples containing GVB^{2+} , serum, and sensitized sheep erythrocytes were used as background controls.

(A_{control}), while heparin served as a positive control. Inhibition of hemolysis induced by the test sample was calculated by the formula: $[(A_{\text{control}} - A_{\text{sample}}) / A_{\text{control}}] \times 100\%$ (Diallo et al., 2001).

4.11. Evaluation of ROS scavenging activity

ROS ($\text{O}_2^-/\text{H}_2\text{O}_2$) were generated enzymatically using a xanthine/xanthine oxidase system consisting of 500 μM xanthine, 18 mU/ml xanthine oxidase, and 0.1 M phosphate buffer (pH 7.5). This system produces O_2^- , which can spontaneously dismutate to H_2O_2 , but does not generate $\text{OH}^\bullet$ (Britigan et al., 1990). L-012 (25 μM) was used as a probe to measure ROS in this system. ROS-scavenging activity was evaluated in the presence or absence of polysaccharide fractions at 25°C. SOD (25 U/ml) was used as a positive control for O_2^- -scavenging activity.

$\text{OH}^\bullet$ was generated enzymatically using an H_2O_2 /HRP system consisting of 7 μM H_2O_2 , 0.25 $\mu\text{g/ml}$ HRP, 25 μM L-012, and 0.1 M phosphate buffer (pH 7.5), and $\text{OH}^\bullet$ production was determined by monitoring L-012 chemiluminescence. Scavenging activity was evaluated by monitoring $\text{OH}^\bullet$ in the presence or absence polysaccharide fractions at 25°C. Mannitol (200 mM) and dextran (5 mM) were used as positive controls for $\text{OH}^\bullet$ -scavenging activity.

4.12. Cytotoxicity assay

Cytotoxicity was analyzed with a CellTiter-Glo Luminescent Cell Viability Assay Kit (Promega, Inc., Madison, WI) according to the manufacturer's protocol. Briefly, J774.A1 cells were cultured at a density of 3×10^4 cells/well with the polysaccharide fractions for 24 hr. at 37°C and 5% CO_2 . Following treatment, the cells were allowed to equilibrate to room temperature for 30 min, substrate was added, and the samples were analyzed with a Fluoroskan Ascent FL.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

We would like to thank Dr. Scott Busse, Montana State University, Bozeman, MT for help in running NMR samples and Dr. Christopher West and Wendy Ives, Oklahoma Center for Glycobiology Analytical Core Lab, Oklahoma City, OK for expert monosaccharide analysis. This work was supported in part by Department of Defense grant W9113M-04-1-0001, National Institutes of Health grant RR020185, and equipment grant from the M.J. Murdock Charitable Trust, and the Montana State University Agricultural Experimental Station. The U.S. Army Space and Missile Defense Command, 64 Thomas Drive, Frederick, MD 21702 is the awarding and administering acquisition office. The content of this report does not necessarily reflect the position or policy of the U.S. Government.

References

- Baugh JA, Bucala R. Mechanisms for modulating TNF α in immune and inflammatory disease. *Curr. Opin. Drug Discov. Devel* 2001;4:635–650.
- Beutler B. Innate immunity: an overview. *Mol. Immunol* 2004;40:845–859. [PubMed: 14698223]
- Britigan BE, Pou S, Rosen GM, Lilleg DM, Buettner GR. Hydroxyl radical is not a product of the reaction of xanthine oxidase and xanthine. The confounding problem of adventitious iron bound to xanthine oxidase. *J. Biol. Chem* 1990;265:17533–17538. [PubMed: 2170383]
- Conti P, DiGioacchino M. MCP-1 and RANTES are mediators of acute and chronic inflammation. *Allergy Asthma Proc* 2001;22:133–137. [PubMed: 11424873]
- Dahlgren C, Karlsson A. Respiratory burst in human neutrophils. *J. Immunol. Methods* 1999;232:3–14. [PubMed: 10618505]

- Daiber A, August M, Baldus S, Wendt M, Oelze M, Sydow K, Kleschyov AL, Munzel T. Measurement of NAD(P)H oxidase-derived superoxide with the luminol analogue L-012. *Free Radic. Biol. Med* 2004;36:101–111. [PubMed: 14732294]
- De Ruiter GA, Schols HA, Voragen AG, Rombouts FM. Carbohydrate analysis of water-soluble uronic acid-containing polysaccharides with high-performance anion-exchange chromatography using methanolysis combined with TFA hydrolysis is superior to four other methods. *Anal. Biochem* 1992;207:176–185. [PubMed: 1489092]
- Diallo D, Paulsen BS, Liljeback TH, Michaelsen TE. Polysaccharides from the roots of *Entada africana* Guill. et Perr., Mimosaceae, with complement fixing activity. *J. Ethnopharmacol* 2001;74:159–171. [PubMed: 11167034]
- Dong Q, Fang JN. Structural elucidation of a new arabinogalactan from the leaves of *Nerium indicum*. *Carbohydr. Res* 2001;332:109–114. [PubMed: 11403084]
- Dubois M, Gilles KA, Hamilton JK, Rebers PA, Smith F. Colorimetric method for determination of sugars and related substances. *Anal. Chem* 1956;28:350–356.
- El Abbouyi A, Toumi M, El Hachimi Y, Jossang A. In vitro effects of aqueous seeds extract of *Acacia cyanophylla* on the opsonized zymosan-induced superoxide anions production by rat polymorphonuclear leukocytes. *J. Ethnopharmacol* 2004;91:159–165. [PubMed: 15036483]
- Finlay BB, Hancock REW. Can innate immunity be enhanced to treat microbial infections? *Nat. Rev. Microbiol* 2004;2:497–504. [PubMed: 15152205]
- Gabay C. Interleukin-6 and chronic inflammation. *Arthritis Res. Ther* 2006;8:S3. [PubMed: 16899107]
- Gane AM, Craik D, Munro SL, Howlett GJ, Clarke AE, Bacic A. Structural analysis of the carbohydrate moiety of arabinogalactan-proteins from stigmas and styles of *Nicotiana glauca*. *Carbohydr. Res* 1995;277:67–85. [PubMed: 8548791]
- Gauss KA, Bunger PL, Crawford MA, McDermott BE, Swearingen R, Nelson-Overton LK, Siemsen DW, Kobayashi SD, DeLeo FR, Quinn MT. Variants of the 5'-untranslated region of human *NCF2*: Expression and translational efficiency. *Gene* 2006;366:169–179. [PubMed: 16310324]
- Ghosh P, Adhikari U, Ghosal PK, Pujol CA, Carlucci MJ, Damonte EB, Ray B. In vitro anti-herpetic activity of sulfated polysaccharide fractions from *Caulerpa racemosa*. *Phytochemistry* 2004;65:3151–3157. [PubMed: 15541745]
- Hardy MR, Townsend RR. High-pH anion-exchange chromatography of glycoprotein-derived carbohydrates. *Methods Enzymol* 1994;230:208–225. [PubMed: 8139497]
- Hayakawa Y, Hayashi T, Hayashi T, Niiya K, Sakuragawa N. Selective activation of heparin cofactor II by a sulfated polysaccharide isolated from the leaves of *Artemisia princeps*. *Blood Coagul. Fibrinolysis* 1995;6:643–649. [PubMed: 8562835]
- Hoebe K, Janssen E, Beutler B. The interface between innate and adaptive immunity. *Nat. Immunol* 2004;5:971–974. [PubMed: 15454919]
- Hong SH, Seo SH, Lee JH, Choi BT. The aqueous extract from *Artemisia capillaris* Thunb. inhibits lipopolysaccharide-induced inflammatory response through preventing NF- κ B activation in human hepatoma cell line and rat liver. *Int. J. Mol. Med* 2004;13:717–720. [PubMed: 15067376]
- Hsu MJ, Lee SS, Lee ST, Lin WW. Signaling mechanisms of enhanced neutrophil phagocytosis and chemotaxis by the polysaccharide purified from *Ganoderma lucidum*. *Br. J Pharmacol* 2003;139:289–298. [PubMed: 12770934]
- Hume DA. The mononuclear phagocyte system. *Curr. Opin. Immunol* 2006;18:49–53. [PubMed: 16338128]
- Hwang JS, Chung HK, Bae EK, Lee AY, Ji HJ, Park DW, Jung HJ, Cho CW, Choi HJ, Lee DS, Lee KR, Youn HJ. The polysaccharide fraction AIP1 from *Artemisia iwayomogi* suppresses apoptotic death of the mouse spleen cells in culture. *Arch. Pharm. Res* 2003;26:294–300. [PubMed: 12735687]
- Imada I, Sato EF, Miyamoto M, Ichimori Y, Minamiyama Y, Konaka R, Inoue M. Analysis of reactive oxygen species generated by neutrophils using a chemiluminescence probe L-012. *Anal. Biochem* 1999;271:53–58. [PubMed: 10361004]
- Inngjerdingen KT, Coulibaly A, Diallo D, Michaelsen TE, Paulsen BS. A complement fixing polysaccharide from *Biophytum petersianum* Klotzsch, a medicinal plant from Mali, West Africa. *Biomacromolecules* 2006;7:48–53. [PubMed: 16398497]

- Kelsey RG, Shafizadeh F. Sesquiterpene lactones and systematics of the Genus *Artemisia*. *Phytochemistry* 1979;18:1591–1611.
- Kim JY, Kim DY, Lee YS, Lee BK, Lee KH, Ro JY. DA-9601, *Artemisia asiatica* herbal extract, ameliorates airway inflammation of allergic asthma in mice. *Mol. Cells* 2006;22:104–112. [PubMed: 16951557]
- Klayman DL. Qinghaosu (artemisinin): an antimalarial drug from China. *Science* 1985;228:1049–1055. [PubMed: 3887571]
- Lee SH, Lee MY, Kang HM, Han DC, Son KH, Yang DC, Sung ND, Lee CW, Kim HM, Kwon BM. Anti-tumor activity of the farnesyl-protein transferase inhibitors arteminolides, isolated from *Artemisia*. *Bioorg. Med. Chem* 2003;11:4545–4549. [PubMed: 14527550]
- Li S, Patapoff TW, Nguyen TH, Borchardt RT. Inhibitory effect of sugars and polyols on the metal-catalyzed oxidation of human relaxin. *J. Pharm. Sci* 1996;85:868–872. [PubMed: 8863279]
- Liu F, Ooi VE, Chang ST. Free radical scavenging activities of mushroom polysaccharide extracts. *Life Sci* 1997;60:763–771. [PubMed: 9064481]
- Melo MRS, Freitas JPA, Freitas ALP, de Paula RCM. Isolation and characterization of soluble sulfated polysaccharide from the red seaweed *Gracilaria cornea*. *Carbohydrate Polymers* 2002;49:491–498.
- Moerman, D. Native American Ethnobotany. Portland: Timber Press, Inc.; 1998.
- Morimoto M, Saimoto H, Usui H, Okamoto Y, Minami S, Shigemasa Y. Biological activities of carbohydrate-branched chitosan derivatives. *Biomacromolecules* 2001;2:1133–1136. [PubMed: 11777384]
- Nergard CS, Diallo D, Michaelsen TE, Malterud KE, Kiyohara H, Matsumoto T, Yamada H, Paulsen BS. Isolation, partial characterisation and immunomodulating activities of polysaccharides from *Vernonia kotschyana* Sch. Bip. ex Walp. *J. Ethnopharmacol* 2004;91:141–152. [PubMed: 15036481]
- Nergard CS, Kiyohara H, Reynolds JC, Thomas-Oates JE, Matsumoto T, Yamada H, Patel T, Petersen D, Michaelsen TE, Diallo D, Paulsen BS. Structures and structure-activity relationships of three mitogenic and complement fixing pectic arabinogalactans from the malian antiulcer plants *Cochlospermum tinctorium* A. Rich and *Vernonia kotschyana* Sch. Bip. ex Walp. *Biomacromolecules* 2006;7:71–79. [PubMed: 16398500]
- Paulsen BS. Plant polysaccharides with immunostimulatory activities. *Curr. Organic Chem* 2001;5:939–950.
- Quinn MT, Ammons MC, DeLeo FR. The expanding role of NADPH oxidases in health and disease: no longer just agents of death and destruction. *Clin. Sci* 2006;111:1–20. [PubMed: 16764554]
- Sakagami H, Kohno S, Takeda M, Nakamura K, Nomoto K, Ueno I, Kanegasaki S, Naoe T, Kawazoe Y. O₂^{•−} scavenging activity of lignins, tannins and PSK. *Anticancer Res* 1992;12:1995–2000. [PubMed: 1338273]
- Samuelsen AB, Westereng B, Yousif O, Holtekjolen AK, Michaelsen TE, Knutsen SH. Structural features and complement-fixing activity of pectin from three *Brassica oleracea* varieties: white cabbage, kale, and red kale. *Biomacromolecules* 2007;8:644–649. [PubMed: 17253763]
- Schaeffer DJ, Krylov VS. Anti-HIV activity of extracts and compounds from algae and cyanobacteria. *Ecotoxicol. Environ. Safety* 2000;45:208–227. [PubMed: 10702339]
- Schepetkin I, Potapov A, Khlebnikov A, Korotkova E, Lukina A, Malovichko G, Kirpotina L, Quinn MT. Decomposition of reactive oxygen species by copper(II) bis(1-pyrazolyl)methane complexes. *J. Biol. Inorg. Chem* 2006;11:499–513. [PubMed: 16680452]
- Schepetkin IA, Faulkner CL, Nelson-Overton LK, Wiley JA, Quinn MT. Macrophage immunomodulatory activity of polysaccharides isolated from *Juniperus scopolorum*. *Int. Immunopharmacol* 2005;5:1783–1799. [PubMed: 16275615]
- Schepetkin IA, Kirpotina LN, Khlebnikov AI, Quinn MT. High-throughput screening for small-molecule activators of neutrophils: Identification of novel N-formyl peptide receptor agonists. *Mol. Pharm* 2007;71:1061–1074.
- Schepetkin IA, Quinn MT. Botanical polysaccharides: macrophage immunomodulation and therapeutic potential. *Int. Immunopharmacol* 2006;6:317–333. [PubMed: 16428067]
- Serhan CN, Savill J. Resolution of inflammation: the beginning programs the end. *Nat. Immunol* 2005;6:1191–1197. [PubMed: 16369558]

- Tan NS, Ng ML, Yau YH, Chong PK, Ho B, Ding JL. Definition of endotoxin binding sites in horseshoe crab factor C recombinant sushi proteins and neutralization of endotoxin by sushi peptides. *FASEB J* 2000;14:1801–1813. [PubMed: 10973930]
- Tan RX, Tang HQ, Hu J, Shuai B. Lignans and sesquiterpene lactones from *Artemisia sieversiana* and *Inula racemosa*. *Phytochemistry* 1998;49:157–161. [PubMed: 9745767]
- Terho TT, Hartiala K. Method for determination of the sulfate content of glycosaminoglycans. *Anal. Biochem* 1971;41:471–476. [PubMed: 4253906]
- Tosi MF. Innate immune responses to infection. *J. Allerg. Clin. Immunol* 2005;116:241–249.
- van Holst GJ, Clarke AE. Quantification of arabinogalactan-protein in plant extracts by single radial gel diffusion. *Anal. Biochem* 1985;148:446–450. [PubMed: 3933380]
- Wang J, Sun Y, Li Y, Xu Q. Aqueous extract from aerial parts of *Artemisia vestita*, a traditional Tibetan medicine, reduces contact sensitivity in mice by down-regulating the activation, adhesion and metalloproteinase production of T lymphocytes. *Int. Immunopharmacol* 2005;5:407–415. [PubMed: 15652769]
- Williams LM, Ricchetti G, Sarma U, Smallie T, Foxwell BM. Interleukin-10 suppression of myeloid cell activation--a continuing puzzle. *Immunology* 2004;113:281–292. [PubMed: 15500614]
- Zhang J, Huang Y, Hou T, Wang Y. Hypoglycaemic effect of *Artemisia sphaerocephala* Krasch seed polysaccharide in alloxan-induced diabetic rats. *Swiss. Med. Wkly* 2006;136:529–532. [PubMed: 16983594]

Fig. 1. Fractionation of Artemisia polysaccharides by size-exclusion chromatography. Panel A: Polysaccharides from crude Artemisia extract were isolated by DEAE-cellulose chromatography, and the eluate was further fractionated using Sepharose 6B column chromatography. Total carbohydrate content of the fractions was determined by a phenol-sulfuric acid method (detected at 488 nm). Five fractions (designated A-I through A-V) were selected for further analyses. Panel B: Fractions A-I to A-V were analyzed by HP-SEC and monitored with a refractive index detector, as described. The arrows show peak retention times of the indicated standards used for calibration.

Fig. 2.

^1H NMR spectra of native Artemisia polysaccharide fractions. The fractions were dissolved in D_2O , and spectra were recorded at 20°C , as described. The paired doublet at 2.48–2.55 ppm in the spectra is due to the presence of sodium citrate bound to the polysaccharides.

Fig. 3.

Effects of Artemisia polysaccharide fractions on murine macrophage NO and ROS production. J774.A1 cells were cultured for 24 hr with the indicated concentrations of polysaccharide fractions or 50 ng/ml LPS (positive control). NO production was quantified by measuring nitrite in the cell-free supernatants (Panel A), and chemiluminescence was monitored using an L-012 detection system, as described (Panel B). The data are presented as the mean $\pm$ SEM of quadruplicate samples from one experiment that is representative of four independent experiments. Statistically significant differences between control (0 μ g/ml) and the indicated concentrations of Fraction A-I (a,b: $P < 0.001$; f: $P < 0.05$), Fraction A-II (a: $P < 0.001$; c: $P < 0.01$),

Fraction A-III (a: $P<0.001$; d: $P<0.01$), Fraction A-IV (a: $P<0.001$; e: $P<0.05$), and Fraction A-V (a,b: $P<0.001$; f: $P<0.05$) are indicated.

Fig. 4.

Effects of Artemisia polysaccharide fractions on macrophage cytokine production. J774.A1 cells were cultured for 24 hr with the indicated polysaccharide fractions (800 μ g/ml) or LPS (50 ng/ml). The concentration of cytokines in cell-free supernatants was determined by a cytometric bead array analysis (IL-6, MCP-1 - left axis; TNF- α , IL-10 - right axis). The data are from one experiment that is representative of two independent experiments.

Fig. 5.

Complement fixation activity of Artemisia polysaccharide fractions. The indicated concentrations of each polysaccharide fraction were analyzed for their ability to fix complement, as described. Heparin (150 µg/ml) served as positive control. The data are presented as the mean $\pm$ SEM of triplicate samples from one experiment that is representative of four independent experiments. Statistically significant differences between control (0 µg/ml) and the indicated concentrations of Fraction A-I (a: $P < 0.001$; b: $P < 0.01$), Fraction A-II (a: $P < 0.001$), Fraction A-III (a: $P < 0.001$), Fraction A-IV (a: $P < 0.001$), and Fraction A-V (a: $P < 0.001$; b: $P < 0.01$) are indicated.

Fig. 6.

Effects of Artemisia polysaccharides on ROS production by murine and human neutrophils. Murine neutrophils (5 × 10⁴ cells/well) (Panels A and B) or human neutrophils (10⁵ cells/well) (Panels C and D) were treated with 800 μg/ml polysaccharide fraction A-I (Panels A and C) or the indicated concentrations of polysaccharide fractions, and chemiluminescence was monitored immediately for 1 hr in the presence of 25 μM L-012 (symbol legend is shown in Panel F). Panels E and F: Human neutrophils (10⁵ cells/well) were treated with 800 μg/ml polysaccharide fraction A-I (Panel E) or the indicated concentrations of polysaccharide fractions plus serum-opsonized zymosan particles (100 μg/ml), and chemiluminescence was monitored immediately for 1 hr with 25 μM L-012 and 50 U/ml SOD. The data are presented

as the mean $\pm$ SEM of triplicate samples from one experiment that is representative of three independent experiments. Statistically significant differences between control (0 $\mu\text{g/ml}$) and the indicated concentrations of Fraction A-I (a: $P<0.001$), Fraction A-II (a: $P<0.001$), Fraction A-III (a: $P<0.001$), Fraction A-IV (a: $P<0.001$), and Fraction A-V (a: $P<0.001$) are indicated.

Fig. 7.

Analysis of ROS scavenging by Artemisia polysaccharides in J774.A1 macrophages. J774.A1 cells were pre-incubated for 20 hr with 50 ng/ml LPS. Following the culture, the cells were treated with 200 nM PMA plus Artemisia polysaccharide fractions (800 μ g/ml), mannitol (200 mM), SOD (25 U/ml), or dextran (5 mM). Chemiluminescence was monitored immediately after addition of PMA, and the results are presented as % inhibition of chemiluminescence compared to the response of control PMA-stimulated cells. The data are presented as the mean $\pm$ SEM of triplicate samples from one experiment that is representative of three independent experiments. Statistically significant differences (a: $P < 0.001$) compared to control samples treated with PMA alone are indicated.

Fig. 8.

Scavenging of ROS by Artemisia polysaccharide fractions in cell-free systems. The indicated concentrations of polysaccharide fractions [A-I (■), A-II (△), A-III (◇), A-IV (◆), A-V (○)] were added to the xanthine/xanthine oxidase system (Panel A) or H₂O₂/HRP system (Panel B) containing 25 μM L-012, and chemiluminescence was monitored for 20 min. SOD (25 U/ml) was used as a positive control in the O₂⁻/H₂O₂-generating system, while mannitol (200 mM) and dextran (5 mM) were used as positive controls in the OH[•]-generating system. The data are presented as the mean ± SEM of triplicate samples from one experiment that is representative of two independent experiments. Statistically significant differences in inhibition between

control (0 $\mu\text{g/ml}$) and the indicated concentrations of Fraction A-I (c: $P<0.001$), Fraction A-II (b: $P<0.001$), Fraction A-III (a: $P<0.001$), Fraction A-IV (a: $P<0.001$), and Fraction A-V (a: $P<0.001$) are indicated.

Fig. 9.

Analysis of ROS production by polysaccharide-treated neutrophils after polysaccharide removal. Human neutrophils (10^5 cells/well) (Panel A) or murine neutrophils (5×10^4 cells/well) (Panel B) were pre-incubated with polysaccharides (800 $\mu\text{g/ml}$) for 15 min at 25°C . The cells were pelleted by centrifugation, the media was aspirated, and the cells were washed again. The washed cells were placed in assay buffer, and L-012 chemiluminescence was monitored for 1 hr, as described. The data are presented as the mean $\pm$ SEM of five replicate samples from one experiment that is representative of four independent experiments. Statistically significant differences (a: $P < 0.001$) compared to control samples are indicated.

Table 1

Chemical properties and molecular weights of polysaccharide fractions

Polysaccharide Fraction	Protein Content (%)	Arabinogalactan	Sulfate Content (%)	Molecular Weight (kDa)
A-I	0.85	Negative	5.49	355
A-II	1.04	Positive	4.29	251
A-III	1.54	Positive	3.53	126
A-IV	2.03	Positive	4.49	78
A-V	2.39	Positive	3.43	49

Table 2

Sugar composition of polysaccharide fractions

	Polysaccharide Fraction				
	A-I	A-II	A-III	A-IV	A-V
Arabinose	13.64	10.91	14.32	10.02	12.07
Fucose	0.14	0.00	0.00	0.00	0.00
Galactosamine	16.56	6.19	12.07	9.79	18.60
Galactose	7.56	8.93	18.15	15.68	12.41
Galacturonic Acid	1.57	1.79	1.46	0.74	0.74
Glucosamine	0.86	1.17	0.16	0.02	0.11
Glucose	29.47	10.01	0.43	4.73	1.32
Glucuronic Acid	0.04	0.24	0.22	0.14	0.16
Lyxose	3.39	1.86	1.66	1.14	0.98
Mannose	3.05	1.67	1.49	1.03	0.88
Xylose	23.71	57.22	50.04	56.71	52.73

The data are presented as mol% for each sugar. Individual components were identified and quantified based on elution of known standards.