cellular pathology, but with now an additional type—"matrical"—for diseases peculiar to the supporting or "matrix" tissues. Classed under the last-named heading would be such diseases as arthritis, fibrositis, neuritis, etc., about which little is known except that their pathology is unlike that of the other group diseases. But since each of the three groups A, B, and C has its own characteristic make-up (as shown above) it is not surprising that manifestations of abnormality should also be different.

The table shown below summarizes the findings discussed in the text.

TABLE II.—Summary

	A Body Fluids	B Nuclear Tissues	C Anuclear Tissues
Water percentage ...	90 and over	76 to 85	Below 75
Rate of metabolism...	Nil	High	Low
Concentration of chloride	High	Low	Midway between Groups A and B
Relationship of chloride to water	Increases with increasing water percentage		Decreases with higher water percentage
Relative reaction ...	More alkaline	More acid	Midway between Groups A and B
Na	High	Low	"
K	Low	High	"
Protein	Low	Midway between Groups A and C	High

Elsewhere I have acknowledged with gratitude the help received from many friends and from the Medical Research Council. I am also indebted to an anonymous donor for providing the greater part of the expenses of this investigation.

REFERENCES

- ¹ Close, H. G.: *Biochem. Journ.*, 1933, xxvii, No. 4, 967.
- ² Rous, P., and Beattie, W. W.: *Journ. Exper. Med.*, 1926, lxiv, 836.
- ³ Warburg, O., quoted by Bayliss, W. M.: *Principles of General Physiology*, London.
- ⁴ Warburg, O., quoted by Rous, P.: *Journ. Exper. Med.*, 1925, xli, 739.
- ⁵ Green, A. A.: *Journ. Biol. Chem.*, 1932, xcv, 47.

SYMPATHETIC GANGLIONECTOMY FOR GANGRENE DUE TO THROMBO- ANGIITIS OBLITERANS

BY

H. HAMILTON STEWART, M.A., M.B.,
B.CHIR.CANTAB., F.R.C.S.

HONORARY ASSISTANT SURGEON, BRADFORD ROYAL INFIRMARY;
HONORARY SURGEON, BRADFORD CHILDREN'S HOSPITAL

The value of sympathetic ganglionectomy for the treatment of Raynaud's disease is well known, but doubt is still expressed as to its value in the treatment of thrombo-angiitis obliterans, or, as it is often called, juvenile gangrene or Buerger's disease. It is the purpose of this paper to give a brief description of this rather rare disease and to record a case treated successfully by sympathetic ganglionectomy.

THE DISEASE

The cause of thrombo-angiitis obliterans is unknown. It usually occurs in young males between the ages of 17 and 35, and was originally considered to be almost limited to the Hebrew race: this has now been shown not to be the case. Over-indulgence in tobacco has been said to be a determining factor, but there is little evidence to support this theory. Buerger¹ originally suggested an inflammatory or toxic origin, and workers at the Mayo Clinic² have performed experiments which tend to support this view. In this disease thrombosis occurs in the larger arteries of the limbs, and, if the arterial system is extensively involved, gangrene is liable to occur. The capillaries, arterioles, and finest arteries appear to escape. The thrombosis may affect the whole or only a portion

of an artery at one time, but recurrences of the thrombosis will occur, now in this, now in that artery, until the signs of deficient blood supply to the limb become evident. The thrombus is eventually organized, and may be recanalized, and with this process some degree of periarteritis is prone to occur. The veins of the limbs may be affected in the same way.

It is interesting to note that the actual thrombosis would appear to pass unnoticed by the patient, and to be free from constitutional reactions. This was so in the case I am about to describe, even though the patient had suffered from the disease in three of his limbs and was warned that the last one might be affected in the same way. It is not until the obliterative process has extended so widely as to interfere materially with the blood supply of the limb that the characteristic symptoms of the disease become manifest. The patient suffers from the result of the disease rather than from the disease itself. He (or she) complains in the first instance of a sensation of numbness and coldness in the affected limb in the cold weather, and indefinite pains, which he often considers to be rheumatic in origin. These affect particularly the arch of the foot, calf of the leg, and the wrist. The symptoms of intermittent claudication are complained of quite early in the history of the disease, and when the affected limb is allowed to hang down the extremity acquires a purplish blush, which gives place to excessive pallor on elevation of the limb. Degenerative changes occur in the skin and nails, the skin becoming thickened and cornified and the nails brittle and hard. Ulceration is prone to occur under the nails and in any site of local injury. Later, the patient suffers from persistent pain, and sits holding the affected limb. Gangrene eventually supervenes. On examination the affected limb is cold, and if the progress of the disease has been slow some wasting is evident. It may not be possible to feel pulsation in such arteries as the dorsalis pedis, posterior tibial (at the ankle), popliteal, radial, and ulnar arteries.

It is difficult at first to understand how sympathetic ganglionectomy can improve the blood supply to a limb when the larger arteries are blocked with organizing thrombi. It would appear to act by allowing the smaller arteries to dilate and thus improving the collateral circulation. The value of periarthral sympathectomy is known to be transitory. Now if satisfactory results are to be obtained the appropriate sympathetic ganglia corresponding to the limbs must be removed. In the case here recorded the inferior cervical and first thoracic ganglia were removed on both sides by an exposure obtained by costo-transversectomy at the level of the second rib. In this operation the transverse process of the second thoracic vertebra is removed along with about three inches of the second rib. The pleura is then depressed and a good exposure of the sympathetic chain with its branches is obtained, rather better on the left than on the right side. The desired amount of sympathetic tissue can be removed exactly, and the risk of leaving a portion of the first thoracic ganglion behind is very small. The anterior approach is difficult, the exposure often inadequate, and an incomplete operation is liable to result.

The case described below is that of a young man, aged 25, a gentile, who had lost both lower limbs from gangrene due to Buerger's disease, and who was in imminent danger of losing both upper limbs from the same disease. By a timely sympathetic ganglionectomy on both sides it was possible to save these limbs.

CASE RECORD

The patient comes of a healthy family, and with no history of syphilis: on two occasions his Wassermann reaction was negative. At school he was the fastest runner, and was in perfect health until February, 1925, when at the age of 18 a box fell on his right foot. A phlebitis resulted, which

prevented him from working for ten weeks. During the next three years he limped slightly, and experienced "tightening feelings" in the right calf, with stiffness on occasions. Eventually pain occurred in the calf on exercise, and he was unable to walk far without resting. His foot was cold in the winter months. In October, 1928, he complained of persistent burning pain in the right big toe, and he was admitted to hospital with commencing gangrene of this toe. In spite of periarterial sympathectomy and removal of the toe, amputation through the right thigh was necessary in May, 1929. In April, 1930, a similar series of symptoms occurred in the left lower limb, and in November, 1930, amputation through the left thigh was performed.

November, 1931, marked the commencement of trouble in the left upper limb. The hand began to perspire excessively and itch, and the fingers became cold and dead when exposed to the cold air. The condition of the hand became gradually worse, and by February, 1932, there was spreading moist gangrene of the little and ring fingers, and the pulp of the terminal phalanx of the thumb. The patient was in persistent pain, and sat holding his hand. The pulse could not be felt in either radial or ulnar arteries.

The inferior cervical and first thoracic sympathetic ganglia on the left side were removed under general anaesthesia by the route described. While the ganglia were being dissected away the pulse rate became rapid (135), but settled again after interference with the sympathetic had ceased. Recovery was uneventful. The typical signs of sympathetic loss to the head and upper limb on the left side were evident. There was a pin-point pupil, enophthalmos, increase of surface temperature, and absence of sweating to a sharp line of demarcation at the level of the second thoracic segment. For some few days after the operation near vision was misty, and, on reading, the words split into syllables. Distant vision was normal. The effect of the operation upon the hand was excellent. The pain disappeared immediately and the hand became warm. Lines of demarcation occurred, and the gangrenous portions were removed. The gangrenous part of the pulp of the thumb separated and the terminal phalanx was left healthy.

The patient makes the astounding statement that in the following summer he worked his wheel chair to Morecambe and back, a distance of 120 miles, during a week-end! He experienced no trouble with his upper limb, in spite of the hard work entailed. In December, 1932, he was examined, and the condition of the left upper limb was found to be very satisfactory. A faint pulse could now be felt in the radial artery. In the beginning of 1933 the circulation in the right upper limb became defective and gradually deteriorated, and in April, 1933, after knocking his index finger, septic lesions occurred which failed to heal. The hand became painful and the fingers purplish in colour. The pulse could not be felt in the radial or ulnar arteries.

In June, 1933, the inferior cervical and first thoracic ganglia were removed on the right side by the posterior approach, as before. Recovery was uneventful, and the result of the operation excellent, as on the other side. He was discharged from hospital three weeks after operation, and is at the present time (November) carrying out his work as an artificial flower maker.

SUMMARY

1. A brief description of thrombo-angiitis obliterans, or Buerger's disease, is given.
2. A case is described in which both upper limbs were saved from gangrene by sympathetic ganglionectomy. Both lower limbs had been lost previously from the disease.
3. The posterior route through the chest, for the removal of the inferior cervical and first thoracic ganglia, is considered to be better than the anterior approach. The exposure is superior, and incomplete removal of the first thoracic ganglion, liable to occur by the anterior route, is by this means unlikely.
4. Removal of these ganglia on both sides was not accompanied by any permanent ill effects on the brain, eyes, or heart.

I am grateful to Mr. Goyder for permission to publish this case, which was under his care.

REFERENCES

- ¹ *Amer. Journ. Med. Sci.*, 1908, cxxxvi, 567.
- ² *Proc. Mayo Clinic*, November 19th, 1930, p. 337.

RELATION OF THROMBOPHLEBITIS MIGRANS TO THROMBO-ANGIITIS OBLITERANS

BY

A. L. D'ABREU, M.B., CH.B., F.R.C.S.

ASSISTANT IN THE SURGICAL UNIT OF THE WELSH NATIONAL SCHOOL OF MEDICINE, CARDIFF; LATE SURGICAL REGISTRAR, GENERAL HOSPITAL, BIRMINGHAM

In this country thrombophlebitis migrans is usually regarded as a malady quite separate from the general vascular disease known as thrombo-angiitis obliterans. Buerger¹ insists that phlebitis migrans is part of a widespread disease affecting both veins and arteries. He states: "The association of thrombosis of superficial veins of the upper and lower extremities with other evidence of obliteration of the larger arteries occurs in a sufficiently large number of cases to make the affection of the veins almost pathognomonic." After a full description of a series of these cases some of his conclusions are as follows:

1. The disease thrombo-angiitis obliterans is often associated with thrombophlebitis of the superficial veins of the arms and legs.
2. The clinical disease of the superficial veins may be subsiding or it may dominate the clinical picture. . . .
3. Migrating thrombophlebitis may give no symptoms, the signs referable to the deep vessels being of most importance.
4. Patients may suffer at one time from migrating thrombophlebitis, at another from the progress of the occlusive changes in the deep vessels.

He found that in all cases of thrombophlebitis migrans which he studied, the pulses eventually became imperceptible. He based these conclusions upon exhaustive clinical and pathological examinations of limbs, many of which required amputation, and found that both superficial and deep veins, in addition to the arteries, exhibited those curious inflammatory and thrombotic changes so typically seen in thrombo-angiitis obliterans.

In this country we have been slow, or perhaps reluctant, to recognize the apparent unity of the disease processes. Moorhead and Abrahamson² in 1928 described phlebitis migrans as a new disease complex, because in their four cases visceral as well as superficial veins were involved. They write: "It is interesting, though perhaps inapposite, to draw attention to cases quoted by Buerger in which recurrent phlebitis of superficial veins was the precursor of thrombo-angiitis obliterans. Such a sequel has not been encountered by us, and it is likely that Buerger's cases did not belong to the group with which we are dealing." The case described below is reported because the patient for many years clearly belonged to the group described by Moorhead and Abrahamson, with migrating thromboses that involved visceral and superficial veins, and yet ultimately he presented all the features of thrombo-angiitis obliterans, from which he died.

A. B. Walker³ in 1932 published a paper on thrombophlebitis migrans, and collected fifteen cases in which, up to that time, apparently no arterial disease coexisted; he quotes, however, a case recorded by Campbell and Morgan⁴ in which apparently a thrombosis of the posterior cerebral arteries accompanied multiple venous thromboses elsewhere. N. Kletz,⁵ in recording cases of migratory phlebitis, mentions the association with thrombo-angiitis obliterans; he utters the important warning that "though the prognosis has been regarded by most writers as uniformly favourable, this is not so." He records a fatal case, and to this may be added the one described below. F. Parkes Weber⁶ refers to a case he recorded with Schwarz in 1908 under the heading of "Arteritis Obliterans of the Lower Extremity." The patient subsequently developed the typical symptoms of thrombophlebitis