To avoid inconvenience from the diplopia the child subconsciously seeks relief in one of two ways. If his refraction (hypermetropia, and especially anisometropia) favours it, he develops a convergent squint and suppresses the offending image. This, I believe, is what most frequently occurs. I have found a considerable number of

FIG. 1.—Primary position.

my case notes, both private and hospital, in which, in addition to the description of the squint, I have noted that, on looking up and to one side, the opposite eye "shoots up." I was not at the time fully alive to the importance and significance of the observation, and merely recorded it. The other way in which the discomfort of

FIG. 2.—Eyes looking to left.

diplopia is subconsciously avoided is by tilting the head to one shoulder, the so-called ocular or pseudo-torticollis. This expedient will occur, as I imagine, where the refraction is normal, and where there is no tendency to lack of fusion, and therefore to the ready suppression of one image. The appearance is quite typical. The boy (I have only met with it in boys, which lends force to the suggestion of birth injury, since boys have bigger heads than girls) carries his head on one side, rather like a bird, and it is this peculiarity which attracts the notice of the parents.

FIG. 3.—a-b. Line of incision.

REMEDY

Since the inferior oblique is inserted at the back of the lower surface of the eyeball it can only be reached at its origin. This is from a small shallow depression on the orbital floor, just within the inferior orbital margin, close to the opening of the naso-lachrymal canal. A skin incision an inch and a half long is made, commencing at the naso-orbital angle exactly over the inferior orbital margin (Fig. 3). A lachrymal speculum (Müller's) is now inserted, and the incision deepened through fascia and muscle on to the periosteum of the orbital rim, which is felt for with the gloved finger. This is defined in its whole length as far as the naso-orbital angle. Any bleeding points are clamped with forceps, and the deep fascia

lightly incised just within the orbital margin. Some orbital fat may appear, but it may be ignored. A squint-hook is now passed just within the orbit towards the naso-orbital angle, and the muscle will be withdrawn upon it, and will resemble a flat, grey earthworm. It is muscular throughout, and is also hypertrophied. A quarter of an inch is excised and the wound closed with two or three deep sutures. On two occasions I have encountered some difficulty in defining the muscle, but usually no trouble is met with if the deep fascia is carefully incised. The appearance of the muscle on the hook is quite spectacular, owing to its unexpected bulk.

The result in the head-tilting cases is very satisfactory. The head becomes straight, and diplopia is not complained of. I have performed this operation, apart from three head-tilters, on many old squints which had been rectified but in which this factor of an up-shooting eye remained. In future, in such cases I propose to tenotomize the oblique muscle first and see what effect this will have on the convergence, which was probably first induced by this delinquent factor; but this, of course, will depend on the duration of the squint.

CONSTRUCTION OF A VAGINA FROM A LOOP OF SIGMOID

BY

E. ROCK CARLING, F.R.C.S.

SURGEON TO WESTMINSTER HOSPITAL

In February, 1933, a young woman, aged 26, who was contemplating matrimony, sought advice because she had never menstruated. Her figure and general conformation were those of a well-developed young woman; the breasts were full and well formed; the anterior part of the vulva with the clitoris was normal, but there was no vagina. At the vaginal site it was possible, by pressure, to make a very slight dimple. Per rectum no uterine parts could be felt, though the pelvis appeared to be of normal dimensions. The girl stated that she had sexual feeling.

There was a large reducible left inguinal hernia, the contents of which were tender and seemed solid. The hernia was operated upon, and the opportunity was taken to make a thorough examination of the parts. Both ovaries were normal; the tubes with their fimbriated extremities were normal, but each ended in a long, thin, conical body, which appeared to be the rudimentary half of a uterus. In the peritoneal fold behind the bladder each tailed away into nothing. The left tube and uterine remnant entered the hernia, and possibly the ovary also, but this was not verified.

Inquiry elicited a hesitating admission that monthly periods had been recognized, in spite of there being no loss; it is very doubtful if that was really the case.

The operation for formation of a vagina was thus conducted. The whole sigmoid loop, freed on its left aspect, was examined to determine exactly the arrangement of the blood supply. By dividing the vessels so as to leave the springs of successive arcades intact, a length of seven inches of bowel was so isolated as to receive its blood supply from the lowest branch of the inferior mesenteric. Pulsation at the anatomically proximal—but henceforth the distal—end of the isolated loop was full and strong. Both ends were closed by inversion—the anatomically distal end with great care and thoroughness. The divided parts of the large bowel were then united end to end. An incision was now made in the skin, at the posterior part of the vulval site, and a passage enlarged upwards between the bladder and rectum. As soon as it was large enough a full-size tubular speculum was passed up to its full length.

Returning to the abdomen, it was found a little difficult to locate the upper end of the speculum, so as to make the safest possible aperture for the isolated loop of gut; but when well placed, the peritoneum over it was incised and the open end of the speculum cut down upon.

A pair of sponge holders was passed up from below, and a ligature on the closed end of the freed sigmoid loop seized and pulled down to the perineum. The peritoneum and such subjacent parts as could be gathered, together with the lower ends of the rudimentary uterine halves, were attached to and round the upper closed end of the new vagina. The lower end was now opened and the mucosa and muscular wall sutured to the skin margin.

At first there was some tendency of the new tube to diminish in diameter, but an obturator of stent, and later one of vulcanite, was provided, so that the lumen might be kept patent. After a few months an interesting observation was made: about a couple of inches within the orifice a band of circular muscular contraction, about one-half to three-quarters of an inch in extent, was encountered by the examining finger. The first stimulus to the mucosa excited a very strong contraction, which gripped the finger, but if gentle pressure were maintained the muscle relaxed completely, and the whole of the rest of the passage could be reached. Eventually the new vagina admitted a full-sized tubular speculum.

At first there was a considerable discharge of mucus, but with simple douching that was controlled, and generally has been very much less, though it has not entirely stopped. Moreover, on two or three occasions there has been a slight loss of blood, which appeared to come from a little ulcer in the mucosa high up in the passage. The mucosa itself is slowly losing its distinctly intestinal appearance, so that, with the speculum in place, it might pass for a normal vaginal lining.

Marriage, which was postponed for nine months, has been followed by sexual intercourse, believed to be normal. The patient experiences normal sensations, and states that she is able to relax and contract the sphincter-like muscular tract, if not quite at will, certainly with conscious effort. It is an interesting question whether this very definite sphincter-like section of the sigmoid loop represents the sphincter responsible for Hirschsprung's megacolon and allied conditions.

In this case the mesentery of the small intestine was very short, and to have carried a double loop successfully to the perineum would have been a difficult task; indeed, it appeared impossible without putting such tension on the vessels as would have made the life of the loop extremely precarious.

Upon a review of all the circumstances it seems to be wiser to use colon rather than small gut; at any rate, this case indicates that, as an alternative, the colon is available.

The prospective husband was informed of the circumstances, the findings, and the implications. The psychological and other aspects of the whole undertaking were carefully considered.

This is not the first use of sigmoid for the construction of the vagina. At least five cases are recorded—those of Albrecht (three), Boldt and Rüge¹—and probably there are others.² A short bibliography is attached to Mr. Norman Hodgson's report of a successful case, using small intestine, in the *British Medical Journal* of May 13th, 1933 (p. 822).

For the success of this case I am indebted to the skilful and patient co-operation of Miss Burton, the ward sister.

REFERENCES

- ¹ Liebl: *Zentralbl. f. Gynäk.*, 1919, xliii, 288.
- ² Wallace: *Buffalo Med. Journ.*, February, 1911 (unverified).

LEAKING CEREBRAL ANEURYSM

BY

G. D. KERSLEY, M.D., M.R.C.P.

MEDICAL REGISTRAR, ROYAL UNITED HOSPITAL, BATH

As cases of leaking cerebral aneurysm diagnosed *intra vitam* are unfortunately still rare, the case mentioned below seems worthy of record. It is of additional interest, moreover, in that the leak occurred during an eclamptic fit, which somewhat complicated the diagnosis.

CASE REPORT

A multipara, aged 38, was sent into hospital as a case of ante-partum haemorrhage. The history revealed nothing of note except that six months previously she had experienced a sudden intense occipital headache, which had gradually subsided. There was no history suggesting renal involvement. On admission the patient did not appear very ill, but there was marked albuminuria and a blood pressure of 215/140; there was no oedema. There were no labour pains, but the cervix was three fingers dilated; the placenta could not be

A = Ruptured aneurysm. B = Second aneurysm, unruptured.

felt. The following night the patient suddenly fainted, labour pains began, and there was a large discharge of blood from the uterus. The membranes were ruptured, and shortly afterwards a stillborn child was delivered. The mother was very collapsed, with a pulse rate of 130 and a blood pressure of 140. Two hours later she had a fit, lasting six minutes, with opisthotonos and clonic movements of the limbs. Under treatment for eclampsia with morphine and chloral she improved a little, but remained somewhat cyanosed. Till the fourth day there was little change, the patient remaining drowsy. The temperature rose to 101°, the breasts became very tender, and at the same time the patient became more lethargic. The blood pressure had now risen to 180.

On the sixth day after delivery the patient developed a right hemiplegia. The diagnosis at this stage appeared to lie between that of a leaking cerebral aneurysm or of an embolic cerebral abscess. A uraemic hemiplegia was also considered, as was a syphilitic thrombosis, but this was thought to be unlikely. After due consideration it was decided that the most probable diagnosis was leaking aneurysm, and this was confirmed by lumbar puncture, when the cerebro-spinal fluid was found to be intimately mixed with blood, with a coloured supernatant fluid after centrifugalization. The only other findings in the fluid of note were a urea content of 40 mg. per cent. and a negative Wassermann reaction. The patient gradually became more comatose, and died on the tenth day. On post-mortem examination a small ruptured