their bronchoscopic clinic. I have also to thank Dr. Gloyne and those working in the pathological department at Victoria Park Hospital for undertaking the bacteriological examinations.

REFERENCES

- ¹ Zinsser, Hans: *Textbook of Bacteriology*, 1927, Appleton and Co.
- ² Castellani, A.: *British Medical Journal*, 1910, ii, 868.
- ³ Ellman, Philip: *Proc. Roy. Soc. Med.*, 1931, xxiv, 528.
- ⁴ Stokes, Kiser, and Smith: *Journ. Amer. Med. Assoc.*, 1930, xcv.
- ⁵ Steinfield, E.: *Journ. Lab. and Clin. Med.*, 1923, viii, 744.
- ⁶ Oliver, H. Gordon: *London Hosp. Gaz.*, September, 1933.
- ⁷ Pijper, A.: *Med. Journ. South Africa*, 1917, xii, 129.
- ⁸ Idem: *Ibid.*, 1923, xix, 101.
- ⁹ Mendelson, R. W.: *Journ. Amer. Med. Assoc.*, 1921, lxxvii, 110.
- ¹⁰ Joeles, Th., and Simpson, R. H.: *Lancet*, 1923, ii, 108.
- ¹¹ Medlar, E. M.: *Amer. Journ. Path.*, 1927, iii, 305.
- ¹² Millar, W. S.: *Ibid.*, 1927, iii, 315.
- ¹³ Ferris, H. W.: *Amer. Rev. of Tuberculosis*, 1933, xxviii, 348.
- ¹⁴ Maret, P. Janvrin: *British Medical Journal*, 1933, i, 917; *Lancet*, 1932, i, 511.
- ¹⁵ Wade, H. W.: *Philippine Journ. Med. Sci.*, 1918, xiii, B, 271.
- ¹⁶ Stovall, W. D., and Greeley, H. P.: *Journ. Amer. Med. Assoc.*, 1928, xci, 1346.
- ¹⁷ Castellani, A.: *Journ. State Med.*, 1926, xxiv, 202.
- ¹⁸ Jackson, Chevalier: *Proc. Roy. Soc. Med.*, 1930, xxiv, No. 1, 1.
- ¹⁹ Craik, R.: *Med. Press and Circular*, April, 1928.

A NOTE ON THE STUDY OF PULMONARY TUBERCULOSIS IN INFANTS AND CHILDREN

BY

G. GREGORY KAYNE, M.D., M.R.C.P.

DOROTHY TEMPLE CROSS RESEARCH FELLOW IN TUBERCULOSIS

Considerable confusion still exists at the present time on the subject of pulmonary tuberculosis in children. The widely divergent views on frequency, type of disease, and prognosis, held by men of apparently equally high authority, may partly be accounted for by the following factors: (1) conclusions based on the observation of different types of material will vary with the nationality and status of the author, who may be a tuberculosis officer, medical superintendent of a sanatorium for children, physician to a general, children's, or chest hospital, morbid anatomist, or finally, a public health statistician; (2) diagnosis depends on the extent of investigations carried out, particularly as regards radiography, searching for tubercle bacilli when no sputum is available (the rule in children), and employing a tuberculin test; (3) the number of diagnoses confirmed or otherwise must depend on the opportunities for performing necropsies; and (4) the features of the "primary complex" or "primo-infection" are by no means universally appreciated.

I have had the opportunity of studying the methods employed by Armand-Delille and Lestocquoy on the vast material available in the children's department at the Hôpital Hérod in Paris. Their scheme seems logical and complete; moreover, it enables pulmonary tuberculosis in children to be studied in a manner impracticable otherwise. It consists of the following features.

1. Enquête Sociale

For every child admitted a complete inquiry is made by a specially trained health visitor into its past and family history, surroundings, and possibility of contact.

2. Clinical Examination

This implies: (a) a physical examination; (b) x-ray films of the chest, both in the antero-posterior and in the lateral or oblique position; (c) a tuberculin test. The von Pirquet test is used here, as it is almost exclusively elsewhere in Paris. While admitting the greater sensitivity of the intradermal test in dilutions of 1 in 1,000 or 1 in 100, the von Pirquet test is considered sufficient for all practical purposes; (d) the search for tubercle bacilli. These are

looked for as a routine in a gastric washout. A suitable-sized stomach tube is introduced in the morning before the first feed, and 80 to 100 c.cm. of warm water (with a trace of sodium bicarbonate) allowed to run in by holding the container about two feet above the child's head; the gastric contents are then siphoned out by lowering the vessel, and examined by a concentration method, and, if necessary, by inoculating a guinea-pig. Armand-Delille and Lestocquoy claim eminently satisfactory results from this method, and consider it no more inconvenient to the child than swabbing the back of the pharynx after a cough, or removing a small piece of mucus during a laryngeal examination.

3. Necropsy with Anatomical Control of X-Ray Photographs

The method employed here is of particular interest, and is the main object of this note. A necropsy is performed automatically on every fatal case unless the parents take the initiative of objecting immediately after death. The usual post-mortem technique employed elsewhere, which consists in detaching the lungs from the heart either in the thorax or after removal from the body and slicing them with a few vertical or radial cuts, may fail to discover small pulmonary lesions; moreover, the true relation of the lesions to the lung and the heart is

FIG. 1.

difficult to estimate. Preliminary hardening of the organs in formalin causes retraction and loss of normal outline, particularly as regards the costo-vertebral and costo-diaphragmatic portions. Accurate comparison of the lesions with the x-ray photographs is therefore impossible. To obviate these disadvantages Armand-Delille and Vibert devised the following technique.

Hollow moulds representing thoracic cages of various sizes are made by pouring plaster-of-Paris into the thoracic cavities after the contents have been removed through a small sternal opening in bodies with unopened abdomens; while the plaster sets pressure must be maintained on the abdomen to retain the convexity of the diaphragm; plaster bandages are applied over these solid blocks, as in the preparation of a plaster jacket, and the mould thus obtained is subsequently varnished with paraffin. A photograph of such a mould is reproduced in Fig. 1. At the necropsy the whole of the thoracic contents are removed *in toto*, cutting the oesophagus and trachea as high as possible in the neck and leaving the diaphragm *in situ*. A little formalin is injected into the trachea, and the whole mass placed in a mould of the appropriate size, where it is kept in position

by inserting on the inside of the mould a very thin layer of cotton-wool. The mould is then immersed in a 10 per cent. formalin solution for about fifteen days. The whole block, which keeps its form extremely well, is easily dislodged afterwards and cut into frontal slices 1/2 to 1 cm. thick with a brain knife. The slices are washed in running water for two days, then immersed in methy-

FIG. 2.—Sketch of x-ray photograph showing left perihilar shadow.

lated spirit for twenty-four to forty-eight hours, which causes the return of the normal post-mortem tissue colour. To obtain permanent specimens the sections are placed in large Petri dishes filled with the following solution: gelatin, grains 105; thymol, m5; glycerin, 31.75; distilled water, 31.5. Sagittal, oblique, or lateral sections may be obtained in a similar manner.

Examination of the sections enables their situation, form, and relation to other parts to be studied accurately, and permits easy comparison with x-ray photographs taken with the same orientation.¹ Two examples may be

FIG. 3.

FIG. 3.—Sketch of sagittal section of thorax at *a b* (in Fig. 2), showing area of infiltration in left lung (anteriorly) giving rise to the left perihilar shadow. *H* = heart; *V* = vertebra; *R.L.* = right lung; *L.L.* = left lung.

FIG. 4.

FIG. 4.—Similar sagittal section showing area of infiltration in posterior part of left lung, also giving rise to a left perihilar shadow in an antero-posterior x-ray.

given to illustrate the great importance of such a comparison. A good deal has been written about "perihilar tuberculosis." Figs. 2, 3, and 4 explain how an antero-

FIG. 5.

FIG. 5.—Sketch of x-ray photograph showing shadow of varying density in right lung.

FIG. 6.—Sketch of lateral section of right lung at axis *c d* (in Fig. 5), showing that varying density of shadow is due to superimposition of two areas of similar density.

FIG. 6.

posterior photograph may easily suggest a perihilar lesion if one only thinks in terms of two dimensions. An antero-posterior x-ray photograph will often show a shadow of varying density, suggesting different types of lesion. Comparison with the post-mortem sections may, however,

demonstrate that the difference in appearance is due to the superimposition of lesions situated at different depths in the antero-posterior diameter, and actually at the same stage of the pathological process (Figs. 5 and 6). It may therefore be pertinent to conclude by stressing the importance of lateral or oblique x-ray photographs in every case of pulmonary tuberculosis.

It is a pleasure to express my thanks to Dr. Armand-Delille and to Dr. Lestocquoy for their kindness in giving me the opportunity of becoming familiar with their methods, and for the loan of the photograph.

REFERENCE

- ¹ Armand-Delille, P. F., Lestocquoy, Ch., and Huguenin, R.: *La Tuberculose Pulmonaire et les Maladies de l'Appareil Respiratoire chez l'Enfant et l'Adolescent*, Legrand, Paris, 1933.

ON THE LACK OF EVIDENCE OF TRANSMISSION BY HUMAN BEINGS OF TUBERCULOSIS DUE TO THE BOVINE TYPE OF BACILLUS

BY

GILBERT WALKER, M.B., CH.B., M.R.C.P.

LATE ASSISTANT MEDICAL SUPERINTENDENT, GLENLOMOND SANATORIUM, KINROSS

The stimulus for research into the incidence of disease due to the bovine type of tubercle bacillus, and the virulence of that organism for the human subject, was supplied by Koch when, at the Washington Congress in 1908, he asserted that it was less virulent than the human type, and that it had never been demonstrated as a cause of chronic pulmonary tuberculosis in human beings. Investigation has gone on in several countries since that time, and has led to a fuller appreciation of the role of the bovine bacillus in the pathogenesis of human tuberculosis.

The results were summarized in a paper by Griffith and Munro,¹ which also gave an account of ten new Scottish cases of chronic pulmonary tuberculosis due to bovine organisms. Munro, in 1925, began to isolate and type the organisms from every case admitted to the sanatorium with tubercle bacilli in the sputum, and this work is still going on. For the past two years I have been privileged to assist in this, and am indebted to Dr. Munro for permission to use his figures and clinical notes. Up to September, 1933, 595 Glenlondmond strains have been isolated and typed, and of these twenty-three proved to be bovine, while in one case a mixture of bovine and human type bacilli was found, giving a percentage incidence of bovine type of 4. In addition two bovine strains were isolated from a total of forty specimens examined from Ayrshire. Isolation was effected chiefly by cultural methods, rarely by animal inoculation, and in no case did failure to isolate occur, so that no anomalous virus was missed.

Consideration of the group of cases of chronic pulmonary tuberculosis due to the bovine organism reveals several interesting points, and we are concerned here with only one of these. While they are clinically and radiologically indistinguishable from cases due to human bacilli, there is an apparent epidemiological difference at present unexplained—namely, the complete lack of proof of transmission from one human subject to another. How otherwise it is with tuberculosis due to human type bacilli is well known. It is recognized that the disease is produced as the result of contact, more or less prolonged and repeated, with some source of infection, usually an open case of pulmonary disease, and therefore becomes manifest as a "familial" or, more accurately, a "household" disease. So much is this the case that careful inquiry is always made into the family history with a view to