glycogen value for this liver was below what one might expect as normal, but normal control figures for the human do not seem to be readily available. All that can be said is that apparently this liver was poor in glycogen. But in the absence of further evidence, and remembering that the patient took or retained little food for twenty-four hours (apart from the intravenous glucose given about eleven hours before death), it would be rash on this score to attribute the chronic hypoglycaemia to a hepatic cause. To account for the chronic hypoglycaemia one is apparently driven, in the present state of knowledge of this complicated subject, for want of a better explanation, to consider absolute hyperinsulinism; but here again there is very little evidence to go upon. Finally, it is interesting to speculate as to whether the fulminant

course of the bronchopneumonia may have been determined to some degree by the chronic hypoglycaemia.

NOTE.—Since the above was written we have come across a case of apparently true hyperinsulinism, or what certain authors might call dysinsulinism. This was a man, aged 28 years, who complained of slight mental confusion, slight dimness of vision, and weakness about three hours after the heavier meals. The fasting blood sugar was only 80 mg. per 100 c.cm. After ingesting 50 grams of glucose the blood sugar in half an hour was 90, in one hour 135, in two hours 97, in three hours 64, in four hours 66, and in five hours 78 mg. per 100 c.cm. His physical examination was normal in all respects, and we could find no evidence to implicate either the ductless glands or the liver.

REFERENCE

¹ Moore, H., O'Farrell, W. R., Malley, L. K., and Moriarty, M. A.: *British Medical Journal*, 1931, ii, 837.

A CASE OF OVARIAN PREGNANCY*

BY

W. C. SPACKMAN, M.B., B.S.LOND., F.R.C.S.ED.
LIEUTENANT-COLONEL I.M.S.

PROFESSOR OF MIDWIFERY AND GYNAECOLOGY, GRANT MEDICAL
COLLEGE, BOMBAY

(With Special Plate)

The following case of ovarian pregnancy appears worthy of record in view, first, of the great rarity of the condition, and, secondly, of the circumstances leading up to it.

Mrs. C., aged 38, a European, formerly a maternity nurse, consulted me privately on April 26th, 1933, for sterility.

Although married for only twenty months, in view of her age and the fact that her menstruation was becoming scanty she was anxious to become pregnant as soon as possible.

Menstruation.—Every twenty-six days, duration five days; latterly duration three and two days. Last menstrual period April 20th to 22nd.

Physical Examination.—Perfectly healthy type. Pelvic organs normal.

Advised a Rubin inflation of the Fallopian tubes about ten days after first day of each period for three or four months, and to take agomensin for a week before each period.

Treatment.—A Rubin inflation of the Fallopian tubes was carried out on May 1st; pressure 120, both tubes patent. The patient had sexual connexion two or three times in the following few days as advised, but it was later ascertained that she had also had connexion three days before the test was performed.

The patient took agomensin (two tablets three times a day) from May 11th, and was seized with severe lower abdominal pain on the 14th. The pain was so severe that she was admitted to a nursing home. I was out of Bombay, and so am unable to particularize further about this attack. She was given hyoscine. The pain subsided, and she left the home in two days.

She herself attributes the pain to the agomensin. The period expected at the middle of May did not appear. She came to see me again on May 31st. She then had a slight uterine haemorrhage, like a mild menstruation, but no pain. She stated that, since the pain of May 14th, she could not lie comfortably on her left side.

She was not examined, in view of the probability of the missed period indicating pregnancy and the slight haemorrhage a threatened abortion, but she was ordered rest and sedatives, and told to return again in a few days and to keep in touch with me meanwhile.

* This case was reported at a meeting of the Bombay Branch, held in the Grant Medical College, Bombay.

She returned on June 14th and reported that she had "miscarried" on June 8th. From her description of what she passed, and her experience as a nurse, I at first considered that this was probable, but on examining her I found the uterus anteverted, normal in size, and a lump, situated behind and to the left of it, slightly tender, about the size of a plum, and firm in consistency. There was pulsation between this lump and the left side of the uterus. She had a slight blood-stained discharge, brownish red in colour, and slight pain in the lower abdomen. She was told that the condition indicated a possibility of extrauterine gestation, was cautioned to keep very quiet, to remain in communication with me, and to report any developments. In the event of any pain or feeling of illness she was instructed to go to the nursing home immediately and to ring up her doctor.

In the absence of such developments she returned on June 19th, when the following condition was found.

Slight uterine bleeding as before. Slight pain in the hypogastrium. Per vaginam, the lump was found to be nearly double the size noted on the 14th. She was ordered immediately into the nursing home as a case of extrauterine gestation, and I operated on the same day, assisted by Captain H. S. Waters, I.M.S.

On opening the abdomen, there was extremely little free blood in the peritoneum, but, on raising the sigmoid flexure of the colon, the left ovary was found enlarged to the size of a peach, deep red in colour, and with loose adhesions formed by blood clot to the lateral wall of Douglas's pouch, which was subdivided into two halves by oedematous appendices epiploicae. The right half of Douglas's pouch was quite normal. The left half, besides this enlarged ovary, contained a small amount of blood-stained serum. The ovary presented exactly the appearance of a twisted ovary, or ovarian haematoma, clearly limited to the ovary, except for a shell of blood clot on the outer side.

The left Fallopian tube was entirely normal throughout its length, its abdominal ostium being patent and entirely free from any swelling or blood. On freeing the ovary a small amount of this blood clot detached itself from the outer wall of Douglas's pouch, but the ovary remained adherent to the back of the left broad ligament by the ovarian ligament. In fact, the ovary maintained its normal relation to the unaltered mesosalpinx and corresponding tube.

The ovary was resected and the ovarian ligament tied off. The right tube and ovary were both normal. The patient had specially requested the operator to leave both tubes if possible (!), and, the operation being done under spinal anaesthesia, she kept inquiring as to the condition found; for these reasons the left tube, being entirely normal, was allowed to remain. The patient made a good recovery from the operation.

The specimen was immediately examined by the operators. On incising the ovary towards its outer side a gestation sac was found deeply embedded, and, on opening this, an ovum, 0.6 cm. in length, was demonstrated.

Pathological Report.—The specimen was sent for examination by Dr. P. V. Gharpure, professor of pathology, Grant Medical College, who reports as follows.

The specimen is approximately of the size and shape of a small hen's egg, or resembling a small truncated cone, elongated and tapering at one end, and broader at the other. Near the broader end on one of the faces is an oval rough area, 3 by 2 cm., indicating attachment of the ligament, by severing which the specimen has been removed at operation. The specimen measures 5.5 by 4.5 cm. On section it exhibits a non-homogeneous wall nearly uniform in thickness, with a number of blood sinuses and interstitial haemorrhages: the wall has an average thickness of 0.5 cm. It encloses a smooth, shining cavity, lined by a smooth membrane. Inside this cavity, towards the broader end, is attached a tiny white object by a slender pedicle. This is an embryo, 4 mm. at its broadest and 1 mm. at its longest part. It has lost its "C" shape, and has a broader middle and tapering posterior end. Stalks of appendages are recognizable. The embryo does not show any formation of a face. The age, by process of exclusion, is beyond completion of five weeks but before completion of the seventh week. Its internal anatomy has not been studied.

On microscopical examination the wall around the sac contains abundant vascular channels and well-formed chorionic villi, with proper cell content. Some sections show ovarian stroma in anatomical continuity with the chorionic tissue. No localization of a formed placenta can be made out. The entire wall surrounding the gestation sac shows uniform appearances.

COMMENT

This case of ovarian pregnancy satisfies Spiegelberg's¹ criteria, which are as follows:

1. The tube on the affected side must be intact.
 2. The foetal sac must occupy the anatomical position of the ovary.
 3. It must be connected with the uterus by the utero-ovarian ligament.
 4. Definite ovarian tissue must be found in the sac wall.
- This case also satisfies an additional criterion, put forward by Whitridge Williams² in the original edition of his *Obstetrics* (but not included in the latest edition), that ovarian tissue must be present in several portions of the sac wall at some distance from one another.

Although the tube was not removed, and therefore was not examined microscopically, it presented no evidence of abnormality to one who is constantly examining normal and abnormal tubes *in situ*. In this connexion it would appear theoretically necessary to remove and examine both tubes to avoid possible cases where a fertilized ovum has been extruded from one tube, and has found a nidus on the surface of the ovary of the opposite side, such cases being inadmissible as true ovarian pregnancy.

It seems likely that ovarian pregnancy does not occur in a strictly normal ovary. Alternatively, a pregnant ovary undergoes certain histological changes in its structure, with disappearance of immature Graafian follicles. From an examination of the sections of the present specimen, as well as from published illustrations and descriptions of others, it appears that in cases of ovarian pregnancy the ovary itself may be very deficient in Graafian follicles. It is to be noted in the present case that menstruation was becoming less profuse than normal.

The mechanism by which ovarian pregnancy is possible is uncertain. Probably a follicle ruptures with a pinhole

orifice, or the tension of the liquor folliculi is below normal, so that the egg-cell contained in it is not detached. A spermatozoon penetrates, and the fertilized ovum, by the erosive action of its trophoblast, burrows deeply into the ovary. In the present case, as shown in the photograph, and I think in other typical cases, the ovum is found deeply embedded in the ovary and surrounded entirely by ovarian tissue.

Dr. Gharpure, in a recent discussion, has put forward the very interesting and, I think, original view (to which he has kindly permitted me to allude), that all cases of ovarian, and many of tubal, pregnancy may be explained by minute inclusions in these organs of endometrial tissue similar to what occurs in endometriosis; such inclusions or rests would exert a chemotactic attraction for the fertilized ovum. It is possible that careful histological study may afford support for this view.

From the estimate of this ovum being of six weeks' development, it appears that pregnancy followed shortly after the Rubin test. The suggestion that a fertilized ovum, free in its passage along the tube towards the uterus, was driven back by the insufflation, and settled on the ovary must be faced, but it appears improbable (1) because one would expect to find such an ovum on the surface of the ovary rather than in its interior, and (2) because the ovum would be somewhat more advanced, just over seven weeks. Ovulation, moreover, is held to occur about the fifteenth day of the menstrual cycle—that is, about May 5th in the present case, four days after the Rubin test was performed.

Professor Gharpure and myself would be glad to lend the blocks of the ovary from which sections were made to anyone interested.

REFERENCES

- ¹ Spiegelberg: *Arch. f. Gynäk.*, 1878, xiii, 73.
- ² Williams, J. W.: *Obstetrics*, first edition, New York, 1903, p. 537.

The Codex Revision Committee of the Pharmaceutical Society of Great Britain has issued four subcommittee reports. That of the Pharmaceutical Chemistry Subcommittee gives a summary of the principal standards for chemical substances recommended by it, and accepted provisionally for inclusion in the *British Pharmaceutical Codex, 1934*. The Pharmacy Subcommittee has summarized the principal new or revised formulae recommended by this subcommittee for inclusion in the *Codex*. It advises also the restoration of a number of preparations from earlier pharmacopoeias which are not in the *British Pharmacopoeia, 1932*, but are still in more or less frequent demand. For many of the preparations tests have been worked out with a view to the constitution of a *Codex* standard, and alcohol limits for the concentrated infusions, spirits, and tinctures have been noted. The hope is expressed that it will be possible to obtain the permission of the Board of Customs and Excise to use methylated spirit in making preparations not taken internally, such as liniments, paints, and ointments. The report of the Action and Uses Subcommittee summarizes the descriptions and standards submitted for the inclusion of certain substances in the *Codex*, among which are antitoxins, toxins, and gland products. The Dressings Subcommittee proposes for inclusion in the *Codex* basic materials, such as jute, silk, and wool, descriptive monographs of their characters having been prepared. In the case of such dressings as phenol gauze and tow, no quantitative standard for the proportion of medicament present has been recommended. This subcommittee's report relates, therefore, to the more important dressings for which revised requirements have been prepared. It is hoped that this summary will prove particularly useful to manufacturers and others interested in the standardization of dressings. These four reports are published by the Pharmaceutical Press, 23, Bloomsbury Square, W.C.1, the first two costing 2s. 6d. each and the last two 1s. 6d.

W. C. SPACKMAN: OVARIAN PREGNANCY

FIG. 1.—Embryo in situ in centre of ovary.

FIG. 2.—Reverse view showing resection through hilum of ovary.

R. W. BROOKFIELD AND H. V. CORBETT:
DIABETES WITH DEGENERATION OF SUPRARENALS

Photomicrograph of suprarenal gland showing thickened capsule and marked cellular changes in cortex (see text).

H. MORTON ANDERSON: METHOD OF REDUCTION OF FRACTURES OF THE SURGICAL
NECK OF THE HUMERUS

FIG. 1.—Before reduction.

FIG. 3.—Position on fourteenth day.

FIG. 2.—Immediately after reduction, showing bone awl in position.