pneumonia at this stage. Also, a similar rise may be demonstrated in a normal person whose arterial CO_2 is not diminished.

Cyanosis

There is usually little or no change in the cyanosis. What there is cannot be due to increased arterial oxygenation (see table), but might be explained by an increase in the minute-volume of the heart or by a local vaso-motor change. Otherwise it is difficult by the bedside to point to any immediate change in a patient as a result of small quantities of CO_2 at all comparable to the effect of O_2 in lobar pneumonia.

CONCLUSION

The practical conclusion of this paper is that in the absence of respiratory failure, as evinced by shallow

breathing, the clinical benefits of CO_2 administration in air are not sufficiently demonstrated by these observations to warrant its routine use in lobar pneumonia.

REFERENCES

- ¹ Hanson, J. F., and Calhoun, A. W.: *Arch. Int. Med.*, 1932, 1, 269.
- ² Alison, J. F.: *South. Med. Journ.*, 1932, xxv, 386.
- ³ Henderson, Yandell: *Arch. Int. Med.*, 1930, xlv, 72.
- ⁴ Hilton, R.: *Journ. Physiol.*, 1933, lxxvii, Proceedings, February 18th.
- ⁵ Haldane, J. S., Meakins, J. C., and Priestley: *Ibid.*, 1919, lii, 420.
- ⁶ Meakins, J. C.: *Arch. Int. Med.*, 1920, xxv, 1.
- ⁷ Meakins, J. C., and Davies, H. W.: *Journ. Path. and Bact.*, 1920, xxiii, 435.
- ⁸ Uyeno: *Journ. Physiol.*, 1923, lvii, 203.
- ⁹ Binger, C. A. L., Brow, C. R., and Branch, A.: *Journ. Clin. Invest.*, 1924, i, 127.
- ¹⁰ Henderson, Yandell: *Journ. Amer. Med. Assoc.*, 1933, ci, 261.
- ¹¹ Dale, H. H., and Evans, C. L.: *Journ. Physiol.*, 1922, lvi, 125.

RENAL TUBERCULOSIS

WITH A SURVEY OF SEVENTY CASES

BY

ARTHUR JACOBS, F.R.F.P.S.

UROLOGIST, GLASGOW ROYAL INFIRMARY; UROLOGICAL SURGEON,
GLASGOW ROYAL CANCER HOSPITAL

Renal tuberculosis is an insidious disease, for the focus in the kidney is usually well established before its presence gives rise to symptoms. In the majority of cases the outstanding symptoms are referred to the bladder, and consist of increased frequency in urination, with pain before and after the act. The frequency may be variable, but tends to increase, and, as the disease progresses, the patient may require to urinate as often as every twenty minutes, night and day. The urine is acid or neutral, contains albumin and pus, and, frequently, blood. Pain in the renal area is often absent altogether, or is insignificant in degree. Usually a diagnosis of "cystitis" is at first made, but the symptoms fail to respond to the medical treatment commonly prescribed for this complaint. All such cases of persistent frequency, particularly when occurring in adults between 20 and 40, should be suspect.

It is the present consensus of opinion that a normal kidney cannot filter tubercle bacilli from the blood stream into the urine. If, therefore, tubercle bacilli are found in the urine, and a genital tuberculosis can be ruled out, a renal focus must be present, because there is no such thing as a primary tuberculosis of the bladder. Recognition of a genital tuberculosis in the male is usually easy, but it should not be forgotten that this condition is frequently accompanied by a coincident renal infection. In the female the diagnosis of a genital tuberculous lesion may not be so obvious. In the following case a tuberculous cystitis was found to be secondary to infection of a Fallopian tube.

A woman, aged 24, a school teacher, was referred for consultation because of a persistent frequency and dysuria of one year's duration; she required to urinate several times in an hour. She had had a period of sanatorium treatment in Switzerland four years previously for pulmonary tuberculosis, but it was considered that this lesion had healed. The urine contained a heavy deposit of pus. Examination of numerous smears failed to reveal the tubercle bacilli, but a guinea-pig inoculation was positive. Cystoscopy showed a generalized cystitis, and, in places, superficial ulceration. There was considerable adherent purulent lymph on the right side. The base of the bladder to the right of the left ureteric orifice was markedly elevated. The ureteral orifices looked normal, and both ureters were catheterized to the renal pelves. An abundant flow of clear urine came from either side, and both specimens were found to be free from pus, and sterile. Bilateral ascending pyelograms were normal. The x-ray films, however, showed a large, rounded, calcareous shadow in the left side of

the pelvis, above the level of the ischial spine. It was obvious that the tuberculous cystitis was not of renal origin. The diagnosis was settled by a vaginal examination, which revealed an extensive pelvic mass, due to a tuberculous pyosalpinx.

The discovery of the tubercle bacillus is not essential in order to establish a diagnosis. The presence of pus in the urine, but no organisms, is strongly suggestive of a tuberculous infection, and a guinea-pig inoculation should be made. I cannot, however, agree with Wade, who states that "if, on examining the cellular deposit, no organisms are found in it, it is safe to assume that the infection is tuberculous and that you have failed to detect the causal virus."¹ This infers that every case of aseptic pyuria is tuberculous; contrary to this, I have had under observation several such cases, in which an investigation of the complete urinary tract, including guinea-pig inoculation, failed to reveal evidence of a tuberculous lesion.

Since January, 1928, I have seen in hospital and private practice seventy cases of tuberculosis of the kidney. Forty of the patients were males and thirty females; forty-eight were between 20 and 40 years of age, ten were under 20 (the youngest being 7), nine were in the fifth decade, and three in the sixth. Renal pain was a symptom in twenty-two, and in four of these bladder symptoms were absent. The tubercle bacillus was found by examination of direct smears of the centrifugized urine in forty-five, and by guinea-pig inoculation in another ten. The only specimen submitted to the bacteriologist in many instances was the urine taken from the bladder at cystoscopy. The proportion of positives would undoubtedly have been higher if further specimens, taken from twenty-four-hour collections of urine, had been examined. Thirteen patients out of forty-six who were subjected to nephrectomy had the operation performed in the absence of positive urine findings. In two of these a pre-operative diagnosis of hydronephrosis had been made, and the tuberculous nature of the lesion recognized only at operation. In the other eleven a correct pre-operative diagnosis was made in the absence of the positive urine smears, and without waiting for confirmation by animal inoculation.

INVESTIGATION OF A CASE OF RENAL TUBERCULOSIS

(A) *Cystoscopy*.—The appearances of the bladder in cases of renal tuberculosis often make a diagnosis possible in the absence of the causal organism. The evidence thus obtained is frequently characteristic and of immediate diagnostic significance. The first point usually noticed is the reduced capacity of the bladder, which normally is eight to twelve ounces. In addition, the viscus is easily irritated, so that, if too quickly filled with the distending lotion, spasm is induced and completion of the cystoscopy may be prevented. Generally speaking, if the capacity is

less than two ounces cystoscopy is impossible. Where renal pain rather than urinary frequency is an outstanding symptom it is common to find little or no impairment in the bladder capacity. This may be due to the bladder having successfully resisted the infection, or to occlusion of the ureteric orifice on the diseased side. There may then be no appreciable bladder manifestations of the condition. The appearance of the ureteral orifice on the involved side is usually altered. At first it is swollen, reddened, and congested. Tubercles may infrequently be observed near or on the margin. As a result of ulceration the latter becomes irregular, and the meatus gapes; the meatus may, however, become almost closed by oedema of the surrounding mucosa. Finally, the orifice becomes dragged upwards as a result of contraction occurring in the affected ureter, and an appearance aptly described as "golf-hole ureter" obtains. The changes in the bladder itself are not usually of a specific nature. Thus it is only occasionally that tubercles are actually observed. Ulceration following rupture of the tubercles is more frequently seen either in the vicinity of the ureteric orifice or on the fundus. The margins of the ulcers are undermined, the edges irregular, and a slough may occupy the base. Bullous oedema is very common, and may cause the ureteric orifice to be entirely hidden from view. Later, if secondary infection occurs, exuberant polypoid excrescences may form, and may render the ureteric orifices unrecognizable. The whole of the vesical mucosa will then become involved in the inflammatory process, and will look reddened, swollen, and lustreless. This is the stage at which a large percentage of the cases are sent for investigation. The impaired capacity and irritability of such bladders present the cystoscopist with one of his most difficult tasks.

(B) *The Indigo-carmin Test*.—This is of little value in making a diagnosis. It may be useful in helping to locate the diseased side, when the data to be had from the appearance of the orifices are not sufficient, or when the inflammation is so generalized that both sides appear to be affected. I do not consider that a normal dye secretion proves that the corresponding kidney is free from infection. Conversely, a delayed and weakened secretion does not necessarily mean that the kidney is infected. The function of the non-tuberculous kidney may, for example, become impaired from a secondary nephritis, or from dilatation due to back pressure resulting from contracture of the inflamed bladder, and these conditions can be responsible for impairment in the elimination of the dye.

(C) *Catheterization of the Ureters*.—This is the most important step in the whole examination. It is done in order to obtain specimens of the secretion from the separate kidneys for bacteriological and cytological examination, as well as for the urea concentration test. The catheters should be passed up the ureters to the renal pelves, and the urines collected in the second hour after a urea meal (15 grams of urea and 100 c.cm. of water). If leucocytes and tubercle bacilli are absent from one side the disease may be regarded as unilateral. Suspicion must be entertained if leucocytes alone are present, and a guinea-pig inoculation should be carried out. When cystoscopy and intravenous pyelography (see below) reveal a definite involvement of one side, I consider that it is only necessary to catheterize and obtain specimens from the supposedly healthy kidney. If 2 per cent. or more of urea is discovered in the urine the function of the kidney may be regarded as satisfactory.

(D) *Pyelography*.—I do not carry out ascending pyelography as a routine in tuberculous cases, because of the supposed possibility of disseminating the causal virus from the diseased kidney into the circulatory system. Moreover, the examination in many instances is superfluous.

I have no hesitation in using it on suspected cases, however, when there is failure to demonstrate the tubercle bacillus, and I am unaware of any harmful sequelae resulting. The pathognomonic pictures usually obtained frequently allow of an immediate and definite diagnosis. Since the advent of intravenous pyelography I have employed this method, either as a first step in the investigation of cases known to be tuberculous or as an additional check-up after cystoscopy and ureteral catheterization. The radioscopic appearances obtained in this way vary greatly with the phase of the disease. Experience teaches one to recognize as characteristic certain features, such as thinning and lack of definition in the calyces, knobbing and woolliness in their terminal ends, irregular dilatation and fuzzy outline in the renal pelvis, and areas of dilatation and constriction in the ureter. Minor defects which would be observed with the ascending method may not, however, be apparent. The indefiniteness of the shadow frequently makes it impossible to form an opinion as to the nature of the lesion without the help of cystoscopic data. In certain unilateral cases the dilatation in the ureter and pelvis that can occur in the other kidney as a result of back pressure from a contracted bladder may, with an intravenous pyelogram, give an erroneous pictorial impression of bilateral disease. If there is very advanced renal destruction, then, of course, no picture at all is obtained by the intravenous method.

Diagnosis.—The object of the urological examination is to confirm the diagnosis of renal tuberculosis, and to ascertain whether the disease is unilateral or bilateral. It is desirable that this information be obtained with the minimum possible instrumentation. In cases in which the tubercle bacillus is known to be present in the urine it may, as I have indicated, be sufficient to make an intravenous pyelographic examination, followed by a cystoscopy and catheterization of the ureter from the apparently healthy kidney. If the secretion from that side be uninfected, a diagnosis of unilateral tuberculosis can be made. In the event of an intravenous pyelogram giving conclusive evidence of a bilateral involvement, instrumental examination may not be required at all.

Illustrative Cases

Case 1.—Female, married, aged 34, was referred from the out-patient department, where she had been sent because of frequency and dysuria of seven months' duration. The nocturnal frequency was five to six times, the diurnal hourly. A bacteriological examination of the urine had already been made and tubercle bacilli found. Intravenous pyelographic studies showed extensive dilatation in all the calyces of the left kidney; the right side looked normal. At cystoscopy, carried out in the second hour after a urea meal, the bladder capacity was found to be reduced to 50 c.cm. There was a generalized inflammation, but both orifices looked normal. A catheter was passed up to the right renal pelvis, and specimens of clear-looking urine collected. Subsequent examination of the specimens was negative for pus cells or for tubercle bacilli, and the urea concentration was 4.3 mg. per cent. The diagnosis was therefore a left renal tuberculosis. The absence of infection in the right kidney was further confirmed by a guinea-pig inoculation with some of the urine taken from that side, which proved negative for tubercle.

Case 2.—Male, labourer aged 34, complained of frequency of micturition and dysuria of twelve months' duration. The nocturnal frequency was two-hourly, the diurnal hourly. He had had excision of tuberculous cervical glands in childhood. The urine contained albumin, pus, and blood. The bladder capacity was 175 c.cm. There was a mild generalized cystitis. A deposit of muco-pus lay on the base behind the inter-ureteric bar. The left orifice looked dilated, the right normal. A mediumly concentrated secretion appeared at the right ureteric orifice five minutes after an intravenous indigo-carmin injection; no dye appeared from the left orifice after ten minutes' observation. The specimen

of urine collected at the examination was found to contain tubercle bacilli. At a subsequent cystoscopy a specimen was collected from the right kidney. This had a urea concentration of 2.4 mg. per cent., and contained pus cells and tubercle bacilli. A diagnosis of bilateral renal tuberculosis was therefore made.

Case 3.—Male, aged 7, referred from the sanatorium, where he had been under treatment for the previous eight months. He was complaining of generalized abdominal pain, and frequency of two years' duration. He was considerably emaciated. A guinea-pig inoculation had proved positive for tuberculosis. A series of intravenous pyelographic studies was made. Films were exposed after 5, 10, 20, 40, 60, 90, and 105 minute intervals. The left kidney pelvis and ureter were visible onwards from the first film. The pelvis and calyces were grossly dilated, and the ureter was about the calibre of the small intestine. The right shadow was not apparent until the forty-minute film was taken. The dilatation on that side proved to be even greater than on the left. This was deemed to be sufficient evidence of extensive bilateral involvement, and further investigation considered unnecessary.

Case 4.—Female, aged 28, complained of attacks of right renal pain of five weeks' duration. For the previous nine months she had been suffering from intermittent bouts of frequency, which would clear up spontaneously. The urine contained albumin and pus, but was sterile. The bladder capacity was 250 c.cm. The right ureteric orifice was dilated and its margin injected, otherwise the bladder appearances were normal. Catheters were passed up both ureters, and specimens collected from each kidney. The secretion from the right kidney was definitely cloudy, that from the left clear. A bilateral ascending pyelographic examination was made. On the right side the pelvis was slightly dilated, the calyces were all clubbed, and the upper group had a "moth-eaten" appearance. The appearances on the left side were normal. The urine from the right kidney was found to contain a considerable amount of pus, but no tubercle bacilli. Some diphtheroids grew on culture. The left kidney urine contained no cells or organisms, and no growth on culture. A diagnosis of right renal tuberculosis was made because of the pathognomonic appearances of the pyelogram.

TREATMENT

The important clinical fact in the treatment of renal tuberculosis is that, with the exception of cases of the miliary type, only one kidney is involved in an appreciable percentage of sufferers. In making this statement I am not unmindful of the evidence which has been brought forward in recent years to show that, pathologically speaking, the disease is frequently, if not always, bilateral, and that small lesions in the apparently normal kidney can often be found when the opportunity arises to examine it.² But a caseo-cavernous renal tuberculosis is usually unilateral in its early stages, and although it may be possible for a small incipient tuberculous lesion in a kidney to heal, it has never been shown that this can occur with a clinical demonstrable caseous focus. In the majority of cases, if the condition is not successfully combated, there is a progressive spread of the disease throughout the genito-urinary tract. When the bladder becomes seriously involved the patient may be reduced to a state of misery and discomfort seen in few other conditions. If, however, nephrectomy is carried out, a cure can be obtained, and the secondary bladder lesions, if not too far advanced, often heal spontaneously.

As it is considered that renal tuberculosis is always part of a general tuberculous infection,* and that the disease is at least latent in other organs, such as the lungs or mediastinal glands, every patient after operation should have a prolonged period of treatment by means of rest, open air, and nutritious diet. Confinement to bed for

from one to six months is advisable, the period varying with the response of a patient to the operation. A course of tuberculin treatment, given over a period of from eighteen months to two years, may have a beneficial effect on the bladder.

Advanced bilateral renal tuberculosis is not usually a surgical condition, though occasionally a nephrectomy may be justified when there is severe pain, haemorrhage, or obstruction on one side. Such a patient should have sanatorium treatment in the hope that, by building up the defence mechanism, the lesions may be controlled. If this is successful, bilateral disease may be compatible with life, and even with useful work, for a number of years.

Of the seventy cases that I have investigated forty-seven were found to have a unilateral infection and twenty-one were bilateral. The data in the other two cases were incomplete; as the patients refused the necessary examination. Thus the percentage of unilateral infections (69 per cent.) was considerably lower than the figure commonly quoted, which is 80 to 90 per cent.^{3, 4} This question of the relative frequency of bilateral involvement is of the utmost importance, and will be referred to again under prognosis.

RESULTS AND SEQUELAE OF NEPHRECTOMY

Of the forty-seven patients suffering from a unilateral infection, I operated on forty-five and removed the affected kidney. Surgical intervention was not advised in one case because of multiple calculi in the other kidney, and in the other it was refused by the patient. There was one death after operation, and this occurred on the fourth day from duodenal ileus (necropsy). Only one case of advanced bilateral disease was subjected to operation. This was in a woman of 41, who suffered from constant severe left renal pain and begged for relief. It was thought that the right kidney was capable of carrying on the total renal function, as it concentrated a medium-blue dye within ten minutes of an intravenous indigo-carmine injection. The patient, however, never secreted more than two ounces of urine in twenty-four hours after operation, and, in spite of a nephrostomy on the remaining kidney, died on the tenth day.

Healing of the Lumbar Wound

A frequent cause of disappointment after nephrectomy for renal tuberculosis is the breaking down of the wound and the development of slow-healing sinuses. Even when the wound heals per primam a breaking-down may occur several weeks or even months later. This is a universal experience, though scantily referred to by many authors. The complication has apparently little effect on the end-results, but is serious in that it may cause a very protracted incapacity in a patient who would otherwise be well. Crabtree and Cabot⁵ reported only 25 per cent. out of seventy cases as healing by first intention, and Scholfield⁶ reported 40 per cent. out of fifty-seven cases. Twenty-nine of my cases healed by first intention, but ten of these developed sinuses in periods varying from two weeks to two months after operation. Thus the percentage healing by first intention and remaining healed was 42. Post-operative drainage was carried out in ten cases, and six of these developed sinuses; thirty-five were closed without drainage, and out of that number sinus formation occurred in twenty. Neither the presence nor the absence of drainage would therefore appear to be more favourable to sound healing.

Scholfield considers that persistent sinuses are not due to the continuation of the infection from the ureter or from the perinephritic fat, but rather to the development of a new tuberculous process in the traumatized tissues

* Coexisting tuberculous lesions in other parts were noted as follows: pulmonary, seven cases; testicular, five cases; knee-joint, three cases; hip-joint, two cases; sacro-iliac, one case; cervical glands, one case; tibia, one case; foot, one case; cutaneous, one case.

of a patient with a lowered resistance. Removal of the complete ureter, as sometimes advocated, will certainly not ensure healing. One of my patients still has discharging sinuses four years after ureterectomy performed twelve months after removal of the kidney. In most cases, however, with persistent antiseptic treatment, the wound will heal within the year. This has occurred in all but three of my cases: one of these was the patient referred to above, whose wound has been discharging for four years; one other remained unhealed for two years; and another for twenty months.

Effect on the Bladder

Most patients suffering from renal tuberculosis seek relief because of bladder symptoms, and they naturally expect that, after operation, the frequency and dysuria from which they have suffered will be alleviated. After nephrectomy the secondary bladder lesions often heal spontaneously, and the symptoms for which they have been responsible progressively abate. Sometimes this occurs almost immediately, but usually it is a gradual process. It has been stated by Wildbolz⁷ that the bladder may take two to five years to heal. Cystoscopy may show areas of persistent localized ulceration to be responsible for the continuation of the bladder symptoms, and their healing may be brought about by light fulguration. When no such localized lesion can be found, instillations into the bladder of an antiseptic sedative, such as gomenol oil, may be helpful.

In some cases, because of the diffuse inflammatory involvement of the bladder walls, the viscus becomes so contracted that a normal frequency can never be obtained. The pain on urination usually disappears, however. This state of chronic contraction in the bladder renders the remaining kidney liable to a progressive back-pressure destruction. This sequela is well illustrated in the following case.

A girl of 20, who had had a left kidney removed for tuberculosis four years previously, was referred because of total urinary incontinence. This condition had gradually replaced the frequency which had been present prior to her operation. She never at any time passed a proper stream of urine from the bladder, but was constantly wet from dribbling. There was no pain. The urine contained pus and *B. coli*, but no tubercle bacilli, and a guinea-pig inoculation was negative. A cystoscopy was found to be impossible, as the bladder could not be made to retain a drop of fluid. An intravenous pyelographic examination showed an enormous distension of the right renal pelvis and of the complete ureter.

In one case of this series, where there was no appreciable amelioration in the bladder symptoms after the removal of a kidney which was the seat of an extensive caseous destruction, I carried out the operation of resecting the presacral nerve. The result has been most satisfactory.

The patient, a woman of 42, lived in the country and, after the nephrectomy (which was performed in January, 1932), was treated most assiduously by her doctor with tuberculin and instillations into the bladder. She continued to have an hourly frequency night and day, and suffered from a constant irritable feeling in the urethra and severe pain on urination. Tubercle bacilli were still being secreted in the urine, and there was a generalized hyperaemia of the bladder mucosa, which bled easily. There were two localized ulcerated areas present, and I treated these with diathermy without giving any relief. Two months later I carried out the presacral resection. The dysuria disappeared immediately and the frequency became a three-to-four-hourly one. This complete cessation of symptoms has continued since the operation.

PROGNOSIS

How many patients are permanently cured of their urinary tuberculosis by removal of the diseased kidney? In a paper by Wildbolz⁷ of Bern, in 1929, details were given

of 270 cases, all of which had been operated on at least ten years previously, and many dated back to over twenty years. It was claimed that 59 per cent. of these 270 patients were alive, and, with the exception of three, were free from any evidence of urinary tuberculosis. As against this the mortality of unoperated unilateral cases is, according to Person,⁸ 82.5 per cent. within five years.

None of the cases in my series goes back further than six years, but the follow-up which has been carried out has elicited some interesting data. Out of the forty-five unilateral cases that were nephrectomized four have died. Death occurred in two months, nine months, thirteen months, and two years respectively after operation. By a curious coincidence these four cases all belong to the 1931 group. A necropsy was performed on only one of these—namely, on the patient who succumbed two years after operation. He had continued to have bladder symptoms, and a positive tubercle bacilli urine. He refused to go to a sanatorium, and carried on with his work as a road labourer. Three weeks after last reporting he was admitted to Duke Street Hospital suffering from tuberculous meningitis. The necropsy confirmed this diagnosis. The remaining kidney was reported to be macroscopically studded with tubercles. The patient who died two months after operation had active pulmonary tuberculosis, and had been resident in a sanatorium until immediately before the operation. This was carried out under spinal anaesthesia, and, following a rapid recovery, he was transferred back to the sanatorium on the tenth post-operative day. Death was reported to be due to generalized tuberculosis. Information about the cause of death in the patient who died nine months after operation was not available. The patient who died in the thirteenth post-operative month was a boy of 12, and he also appears to have succumbed to a generalized tuberculous infection, though definite details were not obtained.

Taking into account the one post-operative death, there are left forty survivors out of forty-five cases. All have been traced except a girl of 18, operated on in 1929, and last seen in May, 1931, when she was in excellent health, had a four-hourly frequency, and a tubercle-free urine. One patient, operated on four years ago, has developed a faecal fistula in the lumbar wound, and is considered to be dying; in her case also the urine is tubercle-free. Of the remaining thirty-eight cases, twenty-one were operated on before the end of 1932, and twenty of these have had the urine examined within the past few months. In four the organism was found in examination of direct smears, and in five others by guinea-pig inoculation. Thus nine out of twenty cases which were considered to be unilateral are still secreting tubercle bacilli, although the offending organ has been removed. The general condition of eight of these patients is very good: they have, on an average, a two-hourly frequency, and no dysuria. This group includes the patient upon whom a resection of the presacral nerve was carried out. The ninth case has a frequency which varies intermittently from half- to two-hourly, and he also suffers from dysuria.

If the percentage of cases with post-operative positive urines is added to the percentage seen in the same period which were found to be suffering from a clinically bilateral infection, then the relative frequency of the latter would appear to be in the neighbourhood of 57 per cent. If my interpretation is correct, the conception held, until recently at any rate, by most urologists, that renal tuberculosis is, as a rule, a unilateral disease, is erroneous. This, however, does not, in my opinion, infer that a nephrectomy is indicated in only some 43 per cent. of cases, for I consider that a kidney which is the seat of a well-established destructive lesion should be removed,

even if the other side is secreting an occasional bacillus of tuberculosis. If such a kidney has a normal anatomical outline on pyelography, exhibits a good renal function, and is secreting little or no pus, the case can be regarded, clinically, as a unilateral one, and a nephrectomy should be carried out. The marked improvement in most of the nephrectomized patients, even though they still have the organism present in the urine, supports this contention. Further support is to be had, I think, from a follow-up of unoperated cases, which were considered, clinically, to be bilateral. An endeavour was made to trace fourteen such cases which had been seen in the years 1928 to 1931 inclusive. It is significant that it was possible to ascertain the fate only of eight, and of these five had died. Of the remaining three, one was in poor health and had a discharging lumbar sinus, following on the rupture of a perirenal abscess. The two others, although still in fairly good general condition, had a half-hourly frequency night and day.

CONCLUSIONS

1. Renal tuberculosis is clinically bilateral in at least 30 per cent. of cases.

2. A case should be regarded as unilateral, and be nephrectomized if the kidney secretion from one side does not contain pus or tubercle bacilli, and if the pyelographic appearances of that side are normal.

3. The presence of tubercle bacilli without pus in the urine of a kidney which has a normal pyelographic outline should not negative removal of the other kidney if it is the seat of an established destructive lesion. The kidney which exhibits occasional bacilli of tuberculosis in its secretion as the sole evidence of its involvement has probably only an initial tuberculous lesion, and it has been demonstrated that these frequently heal.

4. All patients after operation should undergo a sanatorium regime. This is particularly necessary if sinus formation occurs in the wound and if bladder symptoms persist.

My thanks are due to Dr. Robert Cruickshank, bacteriologist, Glasgow Royal Infirmary, under whose supervision all the bacteriological work has been done. I also wish to thank Mr. Mack, assistant in the urological department and McCunn Research Scholar (in bacteriology). He has, during the past year, undertaken a considerable share in the bacteriological investigations, assisted at the examinations and operations, and enthusiastically helped in the follow-up of cases.

REFERENCES

- ¹ Wade, Henry: *Edinburgh Med. Journ.*, 1933, xi, No. 3.
- ² Medlar, E. M.: "Renal Infection in Pulmonary Tuberculosis: Evidence of Healed Tuberculous Lesions," *Amer. Journ. Path.*, 1926, ii, 401.
- ³ Eisendrath and Rolnick: *Urology*, p. 704.
- ⁴ Thomson-Walker, Sir John: *British Medical Journal*, 1927, ii, 625.
- ⁵ Crabtree and Cabot: *Surg., Gynecol. and Obstet.*, 1915, xxi, 669.
- ⁶ Scholfield, B. C.: *Journ. Urol.*, 1928, xx, No. 3.
- ⁷ Wildbolz, Hans: *Ibid.*, 1929, xxi, No. 2.
- ⁸ Person, M.: *Ann. of Surg.*, 1925, lxxxii, No. 4.

A new edition of the well-known handbook of the American Medical Association of Vienna has been issued to cover the academic year 1934-5. It is based on the June, 1931, edition, revisions, corrections, and additions having been effected to bring it up to date. Written in English; it affords a detailed and comprehensive guide to the medical and cultural opportunities for training in that University, special attention being devoted to courses of instruction given in English. For German-speaking visitors details are given of the university courses which are open to them, as also of the many facilities for private tuition and study. Various items of general information reveal the magnitude of the help rendered by this society, membership of which carries various privileges as regards study in Vienna. Copies of the handbook may be obtained from the headquarters of the American Medical Association of Vienna, Alserstrasse 9, Vienna VIII.

THE EFFECTS OF SUBCUTANEOUS AND INTRAVENOUS INJECTION OF TOXINS COMBINED WITH FINE EMULSIONS OF OILS

BY

V. G. WALSH, M.B., B.Ch., B.Sc.

AND

A. C. FRAZER, M.B., B.S.

(From the Department of Physiology, St. Mary's Hospital Medical School)

Experimental work has been carried out by many investigators on the adsorption of toxic substances by colloidal suspensions, and finely divided charcoal and kaolin have been used with considerable success for the adsorption of toxins in the intestinal tract. Particles of kaolin carry a negative charge, and cholera toxin, acting like a positively charged colloid, is adsorbed by the kaolin, and is consequently not absorbed by the intestinal mucosa. Finely divided charcoal will adsorb both tetanus and diphtheria toxins *in vitro*.

AIM OF INVESTIGATION

In order to investigate the properties of finely divided fat particles with regard to the adsorption of toxins and at the same time to overcome the difficulties attendant on the use of solid particles for injection purposes, experiments were conducted with very finely divided emulsions of cod-liver oil, arachis oil, and olive oil. Toxins, such as diphtheria and tetanus toxins, adsorbed on to these fat particles were found to be non-toxic when injected subcutaneously in large doses. The emulsions used in the following experiments consisted in all cases of 3.5 per cent. oil in water, using sodium carbonate as the emulsifying agent, the finished emulsions being adjusted to pH 8 (using the indicator method). Emulsions were prepared with the apparatus previously described by the authors,¹ the oil being very finely dispersed, the size of the particles averaging $1/2$ to 1μ . These emulsions are extremely stable, and will withstand sterilization by boiling.

First Series

Eight rabbits were selected and divided into two groups, A and B. Group A consisted of two rabbits; Group B of six rabbits. Group A—The animals in this group were given a subcutaneous injection of 5 minimum lethal doses of diphtheria toxin mixed with 0.5 c.cm. normal saline. Group B—The animals in this group were injected subcutaneously with 5 minimum lethal doses of diphtheria toxin mixed with 0.5 c.cm. cod-liver oil emulsion. Results: Group A—Death of both animals within thirty-six hours; Group B—No toxic symptoms shown by any of the animals, and all survived.

Conclusion 1.—The toxin-emulsion mixture was non-toxic when injected subcutaneously.

The experiment was now repeated, the animals being divided into Groups A and B as before, but on this occasion the injections were given intravenously. Group A animals were given 5 minimum lethal doses diphtheria toxin with 0.5 c.cm. normal saline injected into ear vein. Group B animals were given 5 minimum lethal doses diphtheria toxin with 0.5 c.cm. cod-liver oil emulsion injected into ear vein. Results: Group A—Death of both animals within twenty-six hours; Group B—Death of all animals within about the same period.

Conclusion 2.—The addition of emulsion to the toxin afforded no protection when the mixture was given intravenously.