A CASE OF HAEMOCHROMATOSIS

BY

S. WATSON SMITH, M.D., M.R.C.P.Lond., F.R.C.P.Ed.

HONORARY PHYSICIAN

AND

HENRY BLAIR, M.B., B.S.Lond., M.R.C.S., L.R.C.P.

RESIDENT MEDICAL OFFICER

ROYAL VICTORIA AND WEST HANTS HOSPITAL, BOURNEMOUTH

Bronzed diabetes is so uncommon, and this case so unusual, that its publication seems to us to be worth while, particularly because of the absence of diabetes mellitus, the extreme obesity, the deep bronzing of the skin, the atrophoderma of the face, all in a girl of 19 years; and because of the post-mortem findings.

CASE RECORD

A girl, aged 19, single, was admitted to the Royal Victoria and West Hants Hospital on 24/10/31 and discharged relieved on 14/12/31; she was referred by Dr. C. G. H. Morse. The diagnosis made was of supranrenal insufficiency, dyspituitarism, and Addisonian anaemia. The condition on discharge was "improved."

History.—Headache two days previous to admission; went to bed in the afternoon; slept soundly, but could not be wakened next morning. Continued comatose for twenty-four hours, and was then admitted to hospital. There was no evidence whatever that she had taken drugs of any kind. Her past history showed that she had always been large and obese, but grew rather suddenly fatter at the age of about 15 years. She had never menstruated. The mentality before this attack was fairly good; she had won prizes for music when at school.

Examination.—Comatose, but resented interference. Was inclined to throw herself about rather violently if attempts were made to rouse her. Would not, or could not, swallow. There was a large erythematous patch over the right thigh and buttock, with a less distinct patch on the left thigh; the patient was very obese, especially about the hips and thighs. Complexion was dark, skin of face coarse. On admission, temperature 102.5° F., pulse 100, respirations 22. There was no paralysis found in any muscle group; pupils reacted to light; knee-jerks were not obtained; ankle-jerks present; plantar response not definite; abdominal reflexes not obtained; ophthalmoscopic examination not possible. Lumbar puncture, although tried on four separate occasions, failed owing to patient's enormous bulk. The blood pressure was 140/90; the heart was clinically not enlarged; there were no murmurs. Urine: S.G., 1022; neutral; contained some albumin, a few red cells, and a trace of sugar.

Progress.—On 29/10/31 the patient had become conscious and seemed to be mentally unbalanced; gave way to silly laughter at times; erythematous patch on right buttock worse, and inclined to break down; had a loose cough; rales were heard at both bases, especially at the right. The pupils reacted to light and accommodation; no paralyses; muscle power seemed to be rather less on the left than on the right side. There was general dulling of sensation, though not a complete loss. There was no pain to be felt on pinprick. Reflexes as before.

20/11/31.—Pigmentation of skin and hair since admission, similar to that of a mulatto, had now an added jaundice. By the appearance of the abdominal striae there was still a considerable recent laying on of fat subcutaneously. There was no slowing of the pulse, or pruritus, or any abdominal pain, or vomiting. The blood count was now beginning to approach that of an Addisonian anaemia, and confirmed the halometric reading of 4.3. A continuation of thyroid extract per os was ordered, and one daily dose of adrenaline mix with pituitary extract mix was given.

1/12/31.—Atrophoderma reticulata symmetrica faciei, brought about by the too rapid stretching of the skin by overgrowth of subcutaneous tissues and skeleton. Blood cultures and subcultures were negative. 4/12/31: Much better during the past few days; extraction of dental stumps caused considerable bleeding, stopped only with difficulty.

7/12/31: Temperature normal for nearly three weeks now; mental condition fairly good, though a little silly; could get about on her legs quite easily; fit for discharge. On 19/11/31 her blood examination had been as follows: red cells, 2,260,000; leucocytes, 9,700 per c.mm.; haemoglobin, 48 per cent.; colour index, 1.06. The stained blood showed large lymphocytes 10 per cent., small lymphocytes nil, polymorphs 85 per cent., transitional cells 5 per cent.; very slight anisocytosis; halometer reading 4.3. Until 26/11/31 the temperature had been remittent, the diurnal variation being between 98.4° and 103.2°, the pulse rate was 76–110, and the respirations 20–24. The bowels moved each day.

Readmission

The patient was readmitted on 26/9/33 and died on 7/10/33. The history was that she had developed oedema of the legs; that she had felt well until January, 1932, and had then had headaches at night. In February, 1932, she found that she was losing her hair. In April albumin was found in the urine. In August, 1932, her father died suddenly and unexpectedly: he was found to have advanced arteriosclerosis and an unusually small pituitary body and fossa. In December, 1932, she complained of pain in the left side, and in May, 1933, of headaches and giddiness.

Present illness.—For two weeks the patient has had swellings of the ankles and thighs, has had puffy eyes in the mornings, and been excessively tired; she has also been very thirsty and has polyuria, with a nocturnal frequency of 5–6. She has indigestion, with flatulence and heartburn. The bowels are constantly constipated. Sleeps badly and invariably has headaches. No past illnesses.

Examination.—Deeply pigmented generally; obese, especially about hips and thighs; no loss of hair; has a number of vertical lineae albicantes on each side of the lower abdomen, and many over each hip region and thigh. Bitemporal hemianopia present. Heart: loud first sound at apex and a systolic bruit there. Appearances suggest dyspituitarism. 29/10/33: Disks and whole of retinae are pale in colour. There is marked atrophy of retinal pigment, which in places is gathered into small masses or thin lines lying in front of the retinal vessels. This pigment is in the deeper layers of the retina or in the superficial layers of the choroid. The condition seems more of a pigmentary retinal and choroidal degeneration than an active inflammatory state. On 6/10/33 Mr. Percy Ross reports that the condition of the fundi is unchanged; that the left external field shows a concentric contraction rather more marked on the temporal side, and in the right eye a twenty degree contraction of both temporal and nasal fields.

6/10/33.—Haematemesis of four ounces. Patient states she had haematemesis when an in-patient previously; spleen felt two fingerbreadths below the left costal margin. Pain in left loin, which has become oedematous; perinephric abscess believed to be present there. Temperature now varies between 100.6° and 102.6° F.; pulse, 100–136; respirations, 22; urine shows albumin in small amount, but no sugar, a deposit of many pus cells, some epithelium, many phosphate crystals, no red cells, and no casts. Blood count much as before, but the leucocytes now number 13,000. The fasting blood sugar 0.143 per cent.; bitemporal hemianopia; slight jaundice. On 7/10/33 she had repeated haematemesis, and gradually sank and died.

PATHOLOGIST'S POST-MORTEM REPORT

Dr. C. G. H. Morse, honorary pathologist to the hospital, kindly made an exhaustive post-mortem examination, and reports as follows.

The body is that of a dark-skinned and fat girl, very large and tall for her age. Face, neck, and hands are darker than the rest of the body. Skin generally is deeply bronzed, with pigmentation no more marked in axillae or folds of the skin. There are no localized masses of fat. Facies slightly mongol in type. Neck short, and thick at the base. No thyroid enlargement. Breasts small, nipples indrawn, and areolae deeply pigmented.

Skull.—Hair soft and black. Skin soft, and cuts like that of a child. Nothing abnormal in size or shape of head; calvarium of normal thickness; dura mater normal. There

is slight general oedema of the brain with some excess of cerebro-spinal fluid; no haemorrhage or tumour found. Pituitary gland normal in size. The right frontal sinus contained a calculus about the size of a pea lying loose in otherwise healthy surroundings. No disease of sphenoid, ethmoid, or cavernous sinuses. Cerebral vessels normal. Nasopharynx normal.

Neck and Thorax.—Thyroid gland slightly small for size of patient. Thymus gland not found. There is passive congestion of both lung bases; findings otherwise normal. Some small amount of clear fluid is present in both pleural cavities. The heart is normal in size, the muscle soft and greasy; the endocardium and aortic lining are stained red, and there are two small patches of recent aortitis near the coronary openings. The valves are healthy.

Abdomen.—The abdominal walls are fat. There are many lineae about the lower abdomen and hips; slight amount of clear fluid in peritoneum; no evidence of fat necrosis. Stomach very distended; contains much gas, and a small quantity of grumous fluid and mucus; no source of haematemesis found; intestines distended but otherwise normal.

Liver.—Small, contracted, green in colour, and feels very hard—weight 2 lb. Gall-bladder distended and full of green bile; contains two pigment stones. On section, the whole liver consists of pale yellowish-brown nodular masses in a thick fibrous stroma. The prussian-blue test is very markedly positive.

Kidneys.—Large, soft, and congested. The left kidney has a large perinephric abscess behind it, with one or two ounces of thick creamy pus; no pus found in the kidney itself.

Pancreas.—Quite black in colour and foul-smelling. On section the tissue does not appear diseased apart from the black colour. Prussian-blue test very markedly plus—in fact, rather more so than in the case of the liver. Great distension of splenic vein is present.

Spleen.—Not enlarged, very soft, and rotten.

Suprarenals.—The right cannot be found. The left suprarenal is very friable, and in a mass of fat, but some of it can be removed in pieces. These appear darker in colour than normal. Uterus, tubes, and ovaries are small, and of infantile type.

Microscopical Findings

Liver.—In addition to data already reported, sections of the liver show an active perilobular fibrosis. The interlobular fibrous tissue is very much thickened, and vascularity is increased. Near the external capsule of the organ, which is very thick, many small areas of six or seven cubical cells, resembling liver cells, are to be seen in the fibrous tissue. The cell outlines are indistinct, but the nuclei are clear and granular. Similar groups of cells are also seen in the dense fibrous tissue at the angles of the lobules near the vessels. There are no more than seven cells seen in one group, and such groups have a thin capsule. The liver tissue itself is completely degenerated everywhere. The cells have hardly taken up the stain; no nuclei can be seen. There are masses of fat globules, especially noticeable near the periphery of the lobules. In most of the lobules the central veins are not seen, but one area shows a sclerosed mass in that site, with a few small lymphocytes round it.

Pancreas.—Sections show an advanced fibrosis. The gland cells are atrophied and partly necrotic. No cells can be recognized at all; iron reaction very marked.

Pituitary Body.—Sections show a healthy gland.

Suprarenal (Left).—Sections show normal gland.

Thyroid.—No marked abnormality, except a slight thickening of the interacinar tissue, with some infiltration by small lymphocytes.

Ovary (Right).—Normal ovarian tissue with active ovulation. (This does not conform with the statement that the patient had no menstruation or menstrual symptoms.)

Spleen.—No marked change; passive congestion only.

Kidney.—Advanced cloudy swelling in almost every part. Glomeruli shrunken and cells indistinct. A few tubules only show almost normal structure.

The condition of the pancreas and liver, which are extensively fibrosed and necrotic, show this case to be one of atrophic cirrhosis of the liver. The origin of the condition is, however, obscure. The pathological findings agree with the diagnosis of bronzed diabetes.

DISCUSSION

From all appearances this was a case of bronzed diabetes. In only one specimen of urine was there any sugar to be found, and that simply a trace. With the skin pigmentation of considerable depth and of a definite bronze hue, especially on exposed parts, and with two occasions on which the patient was jaundiced so that the bilious conjunctiva was clearly evident and the cirrhosis of the liver suspected, one could only suppose that the girl, even at her age of 20 years, had died of haemochromatosis. At no time was the liver palpable; it had become shrunken and small, so that one was uncertain about its size on clinical examination. During life, although a provisional diagnosis of haemochromatosis was made, it was not then possible to confirm it; this was done only after the post-mortem examination. The cause and the pathogenesis of the atrophoderma of the face, rare as this condition is, are capable of easy explanation. In the case of this girl the subcutaneous tissues and skeleton increased so rapidly in growth that the skin could not accommodate itself to the increased bulk, and became overstretched, as it did at the same time about hips and thighs. The lesions of the face occurred in the course of two weeks or so, and, pinkish in colour at first, faded after some weeks.

Clinical Memoranda

EFFECTS OF UTERINE CONDITIONS UPON THE BARIUM MEAL: FIBROIDS SIMULATING CARCINOMA COLI

(With Special Plate)

Abnormal conditions of the uterus may at times have curious effects upon the barium meal examination. The following case is submitted as of possible interest in this respect.

The patient, who was in her fifty-sixth year, was on a visit to this country, and sought advice concerning "weakness, anaemia, and constipation," for which, during the previous year, she had been under constant treatment with practically no improvement. From the extreme pallor the anaemia was obvious; but investigation proved it to be of a secondary character. She was still menstruating somewhat profusely and slightly irregularly, but she did not appear to regard these symptoms as of importance, because she "had been accustomed to them all her life." The history of a myomectomy thirteen years previously was then elicited, and one naturally suspected further fibroid growth.

Attempts to carry out an ordinary examination per rectum or per vaginam met with scant success, for the patient at once and involuntarily fell into a state of clonic spasm, doubtless consequent upon an atresia of the vagina, which all operative efforts had failed to relieve. Her marriage, indeed, had never been consummated. Repeated attempts to examine would no doubt have been successful, for the patient was a sensible woman, and anxious to attain some measure of health; but, out of interest in the influence of uterine conditions upon the barium meal, I gave her a small meal, and asked her to return in eight hours. I felt reasonably certain that I should then obtain a radiograph, which would clearly demonstrate the abdominal condition, and which would be of use to me in discussing with the worried relatives the drastic treatment which I had in mind. The result is shown in the illustration (see Special Plate), which presents apparently the characteristic appearances of carcinoma of the bowel, itself a not unlikely diagnosis in this case. Another x-ray photograph was taken twenty-four hours after the meal, and this presents the same appearances, except that the ascending colon is empty, and a few small masses of the meal, roughly a half to one inch in diameter, have found a way into the rectum.