is it more desirable to maintain the biological outlook. We must think in terms of decades and generations. Once the tubercle bacillus has obtained a foothold whether, as rarely, during intrauterine life or at some subsequent date from infancy onwards, the possibilities resulting from bacillary implantation are manifold, and should be kept in view by the thoughtful physician without fuss or exaggeration.

If the influence of the harboured bacillus be frequently beneficial, making for immunity against further attack from outside, it is to be borne in mind that the bacillus is highly resistant, and may remain dormant for many a day, encapsuled perhaps in some out-of-the-way corner, still capable of reactivity and reproduction, and, given favouring conditions, of spread in all sorts of directions. No need for me in a company like the present to dilate on those favouring conditions which occur in numerous ways and at numerous stages in conventional human life. These occurrences explain much that were otherwise hard to interpret in the life-history of the tuberculous individual

and in mortality tables. It is when you have watched in countless cases the "ups and downs" in the persistent struggle between the invading bacillus and the resisting forces of the host—strong or weak as these may be from time to time—that you begin to understand the mystery of the drama of tuberculous infection, action and reaction, and the interlocking of successive events.

These are a few of the reminiscences and musings which remain with me from fifty years' wandering up and down in the wonderful garden. These rambles have occupied a large part of my life and have brought both pleasure and solace. You have given me the privilege of recalling these happy times aloud. Your kindness in inviting me and in listening has enhanced the pleasure of recollection. I hope that I have not wearied you or unwittingly trodden on anyone's toes, or seemed to make light of the labours of others in the garden. Their friendliness and good will have made me happy, and my admiration of their successful efforts has been unbounded.

PARASITIC DISEASES COMMON TO MAN AND ANIMALS*

BY

THOMAS W. M. CAMERON, M.A., D.Sc., Ph.D.Ed.
RESEARCH PROFESSOR OF PARASITOLOGY, AND DIRECTOR, INSTITUTE
OF PARASITOLOGY, MCGILL UNIVERSITY, MONTREAL

Human beings have acquired their parasites from two main sources: from their own simian ancestors and from environmental circumstances, the most important of which is association with animals. It is not always possible to place any given parasite immediately in its own special category—inquiries into the distant origin of animals and plants must always contain an element of speculation—but most of them can be catalogued with some degree of accuracy. To the first category belong such parasites as malaria, filaria, and pinworms, while to a borderline group between it and the second category belong the hookworms and similar forms, which have very close relations in domesticated animals, but which, acquired in very early days, have now become exclusively human. It is interesting to observe that the human *Ascaris* and *Necator* are probably undergoing a reverse process of evolution, and are becoming parasites of pigs as well as of man.

The purpose of this paper is to discuss parasites which even now are being carried to man from animals, especially animals to be found in Canada. In parenthesis it may be mentioned that all such forms as occur in animals in Britain also occur in Canada. Like the white man in North America, they also were immigrants. However, this continent provided its own quota, and the forms discussed here include a few which are not found in the United Kingdom.

For purposes of convenience we may consider these parasites in the following groups: (I) those which live as adults both in man and in animals; (II) those which live as adults in man and as larvae in animals; and (III) parasites which live as larvae in man and as adults in animals.

GROUP I

The group of parasites living as adults in man and in animals can be subdivided into two, the first division including forms which are sufficiently frequent in human

beings to be considered as "normal," and the second including forms only rarely found in man, which are obviously "accidental" parasites.

"Normal" Parasites

The broad tapeworm (*Diphyllobothrium latum*) of man and carnivores has been known for a long time in Europe, where its modern endemic focus is in the Baltic area; it also occurs, however, in Switzerland and Ireland. It was introduced to the North American continent many years ago, and is now located mainly around the Great Lakes and other lakes from Ontario to Alberta. A number of cases of infection with this parasite have been recorded from New York, but the source appears to be pike from the Great Lakes. The incidence of the tapeworm in Canada is rapidly increasing, and the fact that it infects a variety of wild mammals, such as bear and mink, as well as domesticated ones makes control much more difficult.

The eggs of the tapeworm are passed in the faeces into water, where in due course they hatch. The larvae are swallowed by minute water fleas (*Diaptomus oregonensis*), in which they undergo a certain amount of development: this is completed when the first host is swallowed by a fish. Quite a number of fresh-water fish can serve in this way, but the most important is the pike (*Esox lucius*). The infective stage is reached in the fish, especially in summer, and particularly in medium-sized or smaller specimens. Infection results, of course, from eating raw or improperly cooked fish. This worm is generally associated with a form of anaemia, but the latter is certainly not present in all cases, although there is evidence that the worm does cause a small increase in size of the red cells. There is a distinct eosinophilia in the early stages of the infection; this reaches its maximum about the time of maturation of the worm, after which it rapidly diminishes almost to normal.

The presence of the parasite in Canadian fish is of some importance, and every effort should be made at control: (1) by keeping summer-caught pike off the market; (2) by cooking or freezing fish for consumption by both man and animals; (3) by discouraging the feeding of raw fish to dogs; and (4) by periodically dosing the latter for tapeworms. In the presence of wild reservoir hosts absolute control is difficult, but any steps to prevent the pollution of lake water with infected faeces should be of assistance in reducing the incidence.

The human whipworm, *Trichocephalus trichiuris*, which is quite cosmopolitan, but nowhere very common or serious, also occurs in pigs, while the dwarf tapeworm, *Hymenolepis nana*, which is fairly common in the Southern States, is probably identical with the form found in rats and mice in Canada. The dog flea occurs

* An address delivered before a joint meeting of the Central Canada Veterinary Association and the Ottawa Medico-Chirurgical Society, at Ottawa, April, 1934.

frequently on man, and the human flea on the dog, so that both of these parasites may also be included in this section. The human lice, on the other hand, are peculiar to man, and none of the animal forms can find more than a very temporary lodging on him.

Abnormal Parasites

A very large number of parasites of animals have at one time or another been recorded in man. Many of these are of purely academic interest, but a few occur with sufficient frequency to enable us to regard the animals as "reservoir hosts" for these parasites.

The number of species of *Paragonimus* is still far from being settled, although the consensus of opinion is that probably all are variants of a single form. However, it seems advisable to retain the name *P. kellicotti* for the species of lung-fluke which occurs in North America. Here the normal host is the mink, but it is also common in the musk-rat, and it has been recorded from pig, dog, cat, wild cat, goat, and man. Its distribution in Canada is still very incompletely known, but it seems to be fairly widespread, especially in the east.

The eggs, coughed up, are swallowed and passed to the exterior in the faeces. In about three weeks they hatch, and the larva gains admittance to the operculate snail *Pomatiopsis lapidaria*. This snail is amphibious, so that the eggs on damp ground, as well as in water, have an opportunity of continuing their life-cycle. From three to four thousand cercariae emerge from the snail in about three months after infection, but they develop further only if they are able to penetrate into the thin chitin between the segments of crayfish (*Cambarus spp.*). In the second intermediate host they proceed to the heart region, where they become encysted. (In the Asiatic species they are often found in the musculature.) These continue to develop after encystment, and several weeks may elapse before they become infective.

Infection is by eating the crayfish in an uncooked condition, when the young flukes penetrate the intestinal wall and migrate directly to the lungs. Needless to say, many worms never reach their final habitat, and they are occasionally found alive in various other parts of the body.

Human beings may act as accidental hosts for the liver-flukes of sheep, but this is very rare, and there is reason to believe that many of the recorded cases, which were diagnosed from the presence of eggs in the droppings, were not really infected. The eggs are merely contaminations of liver eaten as food, and are in course of leaving the body. The liver-flukes usually found in man and carnivorous animals belong mostly to the family Opisthorchidae, and while the only genus commonly seen in human beings is *Opisthorchis* (including *Clonorchis*) there is every reason to believe that, given the opportunity, any one of the members of this family could attack human beings. All appear to be fish-carried, and in certain parts of the Far East and in Northern Asia and Northern Europe, where conditions are quite analogous to Northern Canada, human infections are very common. The two genera so far found in North America are *Opisthorchis* and *Parametorchis*; there is increasing evidence that the latter is of considerable importance to dogs in the North. A heavy infection causes progressive weakness, emaciation, and anaemia, and its significance in an animal which provides the sole method of transportation for many months of the year can well be understood.

The members of this family, in addition to the usual snail intermediary which all flukes require, have a second intermediate host in fish. The cercariae which escape from the snail encyst on scales or in the flesh of various species of fish, and the definitive host is infected by eating this. The available evidence suggests that any fish-eating animal may be infected. Man owes his freedom, as a rule, to cooking, as heat quickly destroys the cysts. For this reason fluke is not common in human beings here, but it must be admitted that very little search has yet been made for it. Accordingly, the parasites must be considered as of potential human, as well as actual veterinary, importance.

Dipylidium caninum is an exceedingly prevalent parasite in dogs and cats, which also occurs in children; in fact,

it is probably the commonest human tapeworm seen in Great Britain. The intermediate stage is found in fleas and dog lice, and these are occasionally swallowed by children. It is not a very dangerous parasite, but its presence in children indicates that, hygienically, all is not as it should be. Children also become infected with several species of ascarid worms found in dogs and cats. Here, of course, infection is produced by swallowing the embryonated egg, and implies opportunities for faecal contamination of food or utensils.

The genus *Trichostrongylus* includes a number of very small parasites related to the hookworms living in the alimentary tract from the stomach to the caecum. They have a wide geographical range and live in a great variety of mammals. Their main hosts are domestic ruminants and, possibly, horses, but there are a number of records from man; probably these could be greatly increased if examinations were made more frequently. The life-cycle is somewhat similar to that of the hookworm, except that infection is mainly by ingestion. The eggs resemble those of the hookworms, but can be fairly easily distinguished. It is doubtful if these parasites do much harm to human beings.

Man is sometimes infected with mange mites from the lower animals (particularly *Sarcoptes*), chiefly from horses, cows, and dogs. All the species in this genus are similar morphologically, but strains appear to have evolved for each species of host, and so human infections are generally very amenable to treatment with sulphur dressings. The human strain is much more resistant to treatment, however.

GROUP II

Of the parasites living as adults in man and as larvae in animals *Taenia solium* and *Taenia saginata* are both exclusively human parasites in their adult state. The larva of the former, however, is found in pigs (*Cysticercus cellulosae*), whereas that of the latter occurs in cattle (*Cysticercus bovis*). Their distribution, therefore, corresponds with the distribution of these two animals, but their frequency depends on whether or not meat products are habitually eaten uncooked or only slightly cooked. *Taenia solium* is, accordingly, commoner in Germanic and similar communities, whereas *Taenia saginata* is more often seen than *Taenia solium* in Anglo-Saxon peoples. Both occur in Canada, although their distribution is somewhat erratic. Osler found *Taenia solium* in Montreal, and there are recent records of its local frequency in the Slav communities in the west. It is, of course, a rare parasite in Great Britain and elsewhere among people of British extraction—reports to the contrary notwithstanding—because of their habit of overcooking pork. *Taenia saginata* is found in Great Britain, and, although we have little information of its exact distribution in Canada, it also is most frequent in the west; however, it is quite generally distributed elsewhere, although never very common, as in Britain. *Taenia solium* is quite a dangerous human parasite, as its larval stage can develop in man as well as in pigs; it does not always choose the musculature, but is found subcutaneously or in the central nervous system. Quite a number of cases of lunacy are attributable to it. There is no such danger with *Taenia saginata*.

The Trichina Worm

The most important of the parasites which produce trichinosis in man is the trichina worm, or *Trichinella spiralis*. The parasite itself was only recognized in 1828, when Peacock observed the larvae in London, but it was thirty years later before Zenker was able to show its connexion with the serious disease with which we now associate it; at one stage it caused more public apprehension in Germany than did cholera. For some ten years American pork was excluded from Germany on this account, and from this the present very efficient American system of meat inspection directly resulted.

Canadian as well as British meat inspection ignores the possibility of trichina infection of pork, and no attempt is made to search for it. However, in Europe, elaborate measures are taken and large staffs and special apparatus are employed for this purpose. The importance of trichina

infection depends entirely on the habits of the people. Where raw, undercooked, or smoked pork is customarily eaten it is a parasite to be feared, and it is in nations with such habits that we encounter it most commonly. Where, however, as in Anglo-Saxon countries, it is customary to overcook all pork and pork products, the dangers are negligible; this precaution is probably more effective than the most perfect system of meat inspection, which, after all, may merely lead to a sense of false security. In the United States, where a fair proportion of pigs are infected, there have been about 2,000 human cases reported since 1842, although the actual numbers infected but not diagnosed must be very much higher. Most cases are traceable to the consumption of raw sausage or ham, usually prepared at home or in small establishments. Where pork is prepared on a large scale the dilution of the worms is generally such as to make infection negligible and too small in amount to cause disease. Practically all these American cases were confined to people who had still retained their Central or South European habits, while those of British or French origin were not affected. Quite a number of cases have had their origin in the eating of bear flesh, and we have found the trichina worm in a variety of wild carnivores. It is, of course, essentially a parasite of flesh-eating mammals, and in certain parts of the United States it is fairly common in dogs and cats.

Generally, the disease is chronic in the pig; in fact, there may be no symptoms at all and the flesh is unchanged macroscopically. It is unnecessary here to discuss the course of the disease in man, but it may be of interest to draw attention to the frequent diagnosis of typhoid fever or "ptomaine" poisoning in the acute cases, and to "rheumatism" in the subacute later stages of trichinosis. Like most parasitic infections, in the earlier stages there is observed considerable eosinophilia, which, in this case, may reach as high a figure as 40 per cent. Sir William Osler was probably the first to draw attention to the great diagnostic value of this fact.

The distribution of the trichina worm in Canada is unknown, as it is in Great Britain. Human cases are occasionally recorded, but it must be remembered that symptoms are directly proportional to the numbers of parasites ingested. Cases with slight infection probably show no symptoms at all, are not likely to come to the attention of the physician, and still less likely to be diagnosed as trichinosis.

GROUP III

All the parasites living as larvae in man and as adults in animals are obviously "accidental" forms, although some are of very great importance, such as hydatid cyst, fluke larvae causing cercarial dermatitis, and hookworm and other larvae causing "creeping disease."

Hydatid Infections

Echinococcus granulosus as an adult lives in carnivorous animals, the dog being considered the commonest host, but wild Canidae are often infected and play no small part in its spread; cats are only infrequently infected. The intermediate hosts are legion; practically any mammal may harbour the hydatid cyst. Sheep are most often infected, but horses, pigs, and wild ruminants, such as the moose, are also important hosts. The disease is therefore essentially one of pastoral countries. There sheep constitute a large reservoir from which dogs are continually infected, and the opportunities for human infection are thus enormously increased. Although in human pathology hydatid is generally studied in the adult, it is in youth or even in infancy that the disease is usually contracted. Accordingly, in most cases, the cyst of the adult is one already old—a fact which explains the great proportion of abnormalities found in human hydatids.

It has been considered that drinking-water was the commonest source of infection, but echinococcus eggs, like most other helminth eggs, do not float in water in the state in which they are usually passed in the faeces. The drinking of surface water should not, therefore, cause infection. Contagion occurs most frequently through contact with dogs, especially by acquiring dried eggs on the

hands from a dog's coat contaminated with dried faeces. Faecal contamination of foodstuffs, particularly vegetables, is also probably a common source of infection.

Cercarial Dermatitis

The infective stages of a number of trematodes gain admission to the host by actually penetrating the skin. The most important of these are the blood-flukes or bilharzia worms. If they penetrate the skin of the correct host they continue to develop to maturity; if, however, they happen to have entered the skin of an unsuitable host, they die. Unfortunately, whether they are in the correct host or not, they cause a pronounced reaction at the site of entry, and the name "cercarial dermatitis" has been given to the resulting lesion.

There are no human flukes in north temperate climates which have such a mode of entry, but quite a number of wild birds and mammals are so affected. Their larval stages are common in many lakes, both in Canada and in the United States, and closely related forms occur in many parts of the British Isles and elsewhere. These minute cercariae have consequently become a considerable source of annoyance to bathers, and in some localities are so bad that fully half the people entering the water become affected. The condition starts with a prickly sensation a few minutes after exposure to infection, and urticarial blotches quickly develop round the site of entry. The itching may be intense, and papules or even pustules appear. In as little as four or five days the sores disappear with simple treatment, but in some cases they persist from four to seven weeks. The lesions vary to some extent with numbers, dirtiness of the water, and scratching, as well as individual susceptibility, but the history generally makes the origin of the disease obvious.

Analogous to cercarial dermatitis is the condition called "creeping eruption," in which a variety of parasites of lower animals penetrate the skin of man, and, finding the environment unsuitable, move about in or under the skin. Chief among these are immature hookworms of dogs and cats. While a number of these are present in northern latitudes, we have few records of the condition from there. It has been shown that relatively high temperatures, sufficient to cause visible perspiration, are favourable for the disease, and this may account for the fact that it is most frequently found in the south.

CONCLUSION

Of the parasites commonly found in man in north temperate climates only pinworms (possibly ascarids) and lice are exclusively human. Practically all the others are associated in some way or another with lower animals. The broad tapeworm carried by fish is also found in dogs, cats, and wild carnivores; the whipworm occurs in pigs; the two taenias and the trichina worm are conveyed to man by pork or beef; and hydatid cyst is the larval stage of a dog tapeworm. All of these, as well as cercarial dermatitis, are of considerable public health interest. Of potential importance in Canada are the lung-flukes and liver-flukes of carnivores, *Dipylidium* (the common urban tapeworm of dogs and cats), and the mange mites. It must be remembered, however, that modern parasitology is still a very young subject, and as it develops many more forms will probably have to be added to this list. Most of these parasitic diseases are easily preventable; in fact, the evolution of modern civilization and the activities of the sanitarian, the veterinarian, and the cook have already done much to reduce their incidence. Many, however, are still with us, and it is important to know not only where they occur but how frequently they occur, in order that adequate public health measures may be taken. Bacterial diseases are coming more and more under control, and the education of the public has done much to bring this about. Similar steps with regard to the animal parasites should have similar results, and there are signs that such steps would be not inopportune. These are particularly necessary in communities where habitations are widely scattered and sanitary services, so effective in our towns, are still in a rudimentary state.