to point out that of the twenty-two infants who had been vaccinated with B.C.G. over six weeks before (sixteen over three months before)—all by mouth except two, four a second time since birth—only two were positive to a tuberculin test. Thus:

(a) Vaccinated subcutaneously nine weeks before test. Had been in contact with sputum-positive mother for a few days before admission. Negative Pirquet; positive to Mantoux (1 in 100).

(b) Vaccinated at birth by mouth seventeen months before test. Had been in contact for eleven months with a father who coughs, but whose examinations were stated to be negative as regards tuberculosis. Positive Pirquet and Mantoux (1 in 1,000).

Conclusions

In spite of the relatively small number of children here investigated we feel justified in drawing the following conclusions:

1. The Moro test is the least sensitive of the three tuberculin tests commonly employed. The Pirquet (pure tuberculin) and the Mantoux test (1 in 1,000) are about equally sensitive, the Mantoux (1 in 100) being definitely more sensitive.

2. For use in infants and young children the Moro test* has definite disadvantages as regards technique and interpretation of results. When employed, a control test with ether only should always be carried out.

3. The Pirquet test is, for infants and young children, the simplest and easiest test to perform, and should therefore be employed first. If found to be negative, an intradermal test with a dilution of 1 in 100 should then be carried out. In older children it is worth while to repeat the test, if still negative, with a dilution of 1 in 10.

4. The Mantoux test in a dilution of 1 in 10 should always be carried out with a control injection of the broth used for the preparation of tuberculin.

5. A Pirquet test should be regarded as positive only if there is at least 1 mm. of erythema on each side of the scarified area.

6. The minimum required for a positive Mantoux test with any dilution is an area of erythema 10 mm. in diameter, associated with some swelling to touch, or a well-defined erythema greater in area than this. Reactions showing an erythema 5 mm. in diameter or more should be regarded as doubtful, re-inspected on the fourth day, and, if still doubtful or negative, the test should be repeated with a stronger dilution. If a reaction obtained with a dilution of 1 in 10 is on the fourth day definitely more marked than the control test, but still under the limit of a positive reaction as indicated above, it is advisable to regard the test as doubtful, and repeat it either within ten days or in three months' time.¹

7. It is suggested that instead of maintaining the controversy as to the relative value of the Pirquet and Mantoux tests, the test indicated here (Pirquet followed by Mantoux 1 in 100), combining both, should be adopted as a routine under a different name—for example, "the combined tuberculin test," or "the cutaneous intradermal test."

We should like to express our thanks to Professor Ombrédanne for allowing us to test the patients under his care, to Dr. Guérin, of the Pasteur Institute, for supplying the broth used for the control test, and to the ward sisters for their help during the course of the investigation.

REFERENCE

¹ Hart, P. D'A.: Medical Research Council Special Report No. 164, 1932.

* The tuberculin ointment now prepared by Löwenstein and sold in small tin tubes as "dermotublin" contains only 40 per cent. tuberculin and is recommended to be used on the arm, back, chest, abdomen, or front of leg, after preliminary cleansing with benzene, ether, alcohol, or soap and water.

CHLOROFORM-HYOSCINE AMNESIA IN LABOUR

BY

JAMES CLARK, M.B., CH.B.ED.

In the *British Medical Journal* of May 26th, 1934, there is an admirable article on hyoscine amnesia in labour by Mr. Trevor Barnett. It stresses the value of hyoscine alone in producing an amnesia which is preferable to hyoscine-morphine narcosis. Mr. Barnett has verified that the use of hyoscine without the initial injection of morphine eliminates the oligopnoea of the newborn infant. In his conclusions he stresses three conditions which make it impossible for the general practitioner to employ the method he advocates. These are:

1. The need for constant supervision during labour.
2. Occasional necessity for two nurses to control the patient.
3. That the method cannot be employed unless one is assured of the presence of the possibly necessary skilled attendants. (These, perhaps, meaning a qualified medical assistant and two nurses.)

This valuable method, then, can only be utilized in a nursing home where such skilled staff is provided.

It is my intention to try to show that a modification of the hyoscine method has proved to be of great value as a domiciliary measure. During the last ten years in general practice I have conducted eighty-four confinements on the lines about to be described. The usual antenatal examination and care are observed, and the cases selected are those which one, as a general practitioner, would select for home treatment.

Details of Procedure

Treatment is commenced when the patient complains of pains which are regular and of moderate strength. The strength and duration of pains are best gauged by abdominal palpation. I have not found that any arbitrary division between first and second stages of labour assists one's judgement. The nervous constitution of the patient and the consequent degree of fear and sensitiveness to pain are more important. As the practitioner's efforts are directed towards saving the patient from any avoidable suffering, it is better to institute the treatment too early rather than let the pains assume too severe a character. Erring on the early side does not seem to protract the labour. After all, one is dealing with a psychosomatic process and not with a stereotyped mechanism which must conform to an average standard set by textbooks.

The patient is put to bed. One nurse is in attendance. An initial light anaesthesia is induced by chloroform until the patient is quiet and sleeping between pains. (It is the experience of most general practitioners that women in labour take chloroform very well.) A hypodermic injection of hyoscine (1/100 grain) is given immediately. The patient is lightly under. The hyoscine is repeated in doses of half this amount at intervals varying from half an hour to an hour. The ideal to be aimed at is that the patient should be definitely drowsy and quiet between pains. During pains discomfort is evidenced merely by subdued groans and slight restlessness. Any indication of returning awareness can be controlled by immediate inhalations of chloroform and a subsequent injection of hyoscine. This use of chloroform as an intermediary often makes all the difference between a perfect amnesia and one in which a severe pain is recalled with extraordinary intensity.

One also finds that chloroform controls any restlessness which hyoscine may produce. This, however, is rarely

necessary, as the initial administration of chloroform seems to have a subduing effect throughout. The quiet nature of a confinement conducted in this manner is such that the assistance of only one nurse is ample. After experience of the method, one learns to judge for how long during labour the patient may be left in sole charge of the nurse. She receives instructions within the discretion of the medical attendant as to the times of the next one or two injections of hyoscine. As labour progresses the dose of hyoscine is diminished to 1/300 or 1/400 grain, and the intervals of injection lengthened or shortened according to the judgement of the practitioner. I always avoid giving any hyoscine during the hour immediately preceding delivery, and administer light chloroform from the time that the anus assumes the D shape until the head is born. Should labour be delayed, as it may be in any confinement, pituitrin may be used as in ordinary cases, with certain efficiency. Ordinary everyday stimuli should not affect the patient. By these are meant daylight or artificial light, sounds of outdoor traffic, sounds of quiet voices, and of walking in the room. There is no need, therefore, to darken the room, stuff the patient's ears with cotton-wool, or preserve in the house the silence appropriate to a death-chamber.

In no case does the amnesia seem to affect the condition of the infant at birth. The delivery of the placenta is often delayed, but not unduly so; in this one respect more patience is required than in the ordinary confinement.

Maternal loss and after-recovery remain unaffected. The mother sleeps or remains drowsy for some time. Anyway, the amnesia lasts beyond delivery for a period which is directly proportionate to the extent of the amnesia induced during labour. The mother usually emerges from her amnesia with a strongly expressed disbelief that her baby has arrived.

Advantages of Method

This method is particularly valuable in general practice for the following reasons:

1. Primiparae prefer to be confined at home; the first baby is an event, and the medical attendant is expected to get the patient out of her pain as soon as possible.
2. Multiparae who have previously had protracted or exhausting labours do reasonably demand that they should not "have to go through the same again."
3. It reduces the number of instrumental labours. The practitioner is relieved of the exhortations of the patient and her relatives to "get her out of her suffering soon."

It was in such a case that the writer developed the chloroform-hyoscine technique. A young primipara was sufficiently advanced in labour as to justify vaginal examination. This she subconsciously resisted. All that could be ascertained was that the head *might* be on the perineum. The patient and her relatives demanded early relief. Chloroform was administered with a view to applying Milne-Murray forceps. Under light anaesthesia it was found that though the os was paper thin, the head was not entirely on the perineum. Hyoscine (1/200 grain) was given immediately and two injections each of 1/400 grain at half-hourly intervals. (Morphine was avoided on account of the advanced stage of labour.) A further administration of chloroform was given on the first appearance of the head, which was born under control two hours after the first light chloroform was given. Three hours after delivery the patient was asleep, the amnesia having continued until she was in a natural sleep of fatigue.

4. It enables the general practitioner to enhance his reputation or to establish it. Here is something that clinics do not do, something that general nursing-homes might, but as a rule do not, countenance; yet something that still maintains the status of a medical man as *the family doctor*.

Clinical Memoranda

CARCINOMA VENTRICULI IN A WOMAN OF 90

In view of the extreme age of the patient and the scarcity of recorded cases of stomach cancers in old people, I think the following case worthy of record.

Mrs. A., aged 90, was admitted to the City Hospital, Plymouth, on May 30th, 1934, complaining that for the previous ten days she had had constant retching and vomiting, and had been "unable to keep anything down." The bowels had not been open for the last three days, but there was no history of previous ill-health. On admission the patient was obviously practically moribund and very wasted and cachectic, and there was also cyanosis of the lips and slight oedema of the ankles.

Abdominal examination revealed a hard, freely movable mass, with well-defined margins, situated in the mid-line just below the umbilicus. The mass was about 2 inches by 3/4 inch, and was dull to percussion. Examination of the heart and pulse showed that well-marked auricular fibrillation was present. A probable diagnosis of carcinoma of the transverse colon was made, but the patient was too ill for operation to be contemplated, and she died on June 1st.

At the necropsy it was found that the mass was due to a hard, sclerosing growth of an annular type involving the pyloric end of the stomach; the edges of the growth were extremely ill defined: there was no visible ulceration of the gastric mucosa, and a few hard glands were seen lying in the lesser omentum close to the growth. No visceral deposits were present apart from the primary growth. Microscopical section of the mass demonstrated the growth to be a typical columnar-celled carcinoma. There was extreme gastric dilatation above the growth, proving that almost complete obstruction had occurred. The heart showed extreme fatty degeneration.

COMMENTARY

The literature concerning cancer in the aged is very scanty, and no case of cancer of the stomach in a patient of this age has, I believe, been reported in recent years. According to Osler the ages at which cancer of the stomach occurred in 150 cases at the Johns Hopkins Hospital were:

Age	Cases	Age	Cases
20-30	6	50-60	49
30-40	17	60-70	36
40-50	38	70-80	4

In the *American Journal of Cancer* (February, 1934, xx, 477) Shore reports that in Finsterer's series of resections of the stomach for carcinoma, one-third of the patients were between the ages of 60 and 83. Finsterer found age *per se* no contraindication to operation; in fact, while the immediate mortality in the group (60-83) was 9.5 per cent., compared with 6 per cent. in the younger patients, the percentage of five-year cures was 40 in the aged, compared with 22 in the under 60 class, showing that carcinoma of the stomach in the aged is a relatively benign condition.

It is well known that cancer of the stomach can advance to the inoperable stage without giving much evidence of its presence, and in this case, though a growth was found post mortem which must, by its size, have been present for several months, no symptoms were complained of until a few days before the patient's death from pyloric obstruction.

It is, of course, possible that carcinoma of the internal organs in the aged occurs more often than is supposed, and that the diagnosis is seldom made owing to the rarity with which post-mortems are performed in such patients, and the difficulty in getting histories from persons of advanced years.