ON THE CONTROL OF OBESITY*

BY

A. H. DOUTHWAITE, M.D., F.R.C.P.

ASSISTANT PHYSICIAN, GUY'S HOSPITAL

Sixty million pounds are spent annually in the beauty parlours of London. A considerable proportion of this sum is devoted to preventing the development of obesity or to combating an unsightly rotundity which has crept insidiously upon its victim. Yet this aim can be achieved with certainty without expense—in fact, with the exercise of economy. Two widespread fallacies in treatment should be at once exposed. The first is the wholesale exhibition of thyroid extract even though the gland be functioning perfectly. The other is the prescription of strenuous exercise to individuals already exhausted by the task of carrying about several stone of surplus fat.

DANGERS OF OBESITY

The obvious danger to a fat-laden body of microbic infection, or major operation, is too well known to need emphasis. On the other hand, the association of rotundity with many minor disabilities is less widely recognized. For instance, an extraordinary degree of impairment of muscular efficiency and signs indicative of cardiac embarrassment can be found in many people whose increase of weight at first glance appears to be trivial. All who have conducted post-mortem examinations will testify to the deplorable condition of the heart in the fat person. Not only are the usual pericardial deposits increased a hundredfold, but we find fat penetrating and destroying the myocardium relentlessly. The resulting impairment of the heart's action is almost always accompanied by deficient diaphragmatic movement, which is itself induced by the masses of omental fat which bulge the abdominal wall. No doubt it is a combination of these factors which is responsible for the high incidence of bronchitis and the invariable basal catarrh in the fat middle-aged individual.

The effect of fatness on mental processes is an interesting one. The old dictum of "Laugh and grow fat" should not be taken too seriously. In fact, it is questionable whether the cart has not been put before the horse. The fat person, as a rule, avoids effort, mental and physical, his critical faculties are dulled, and a tendency to laugh rather than to grumble shows his bias towards taking the easier path. This is why women urge their husbands to overeat. They exhibit the same attitude with regard to smoking. The affectionate term "fatty" conjures up a picture of the beaming subject of practical jokes. Yet, from time to time the habitual unruffled serenity of the obese is violently disturbed by an outburst of rage not infrequently occasioned by some quite trivial slight. Few persons realize what they are losing in respect of alertness by being fat until they take measures to restore their normal girth. It is during recovery that these patients exclaim with delight at the return of energy, interest in life, and the sensation of youth which accompanies it.

Loss of sexual attraction is an important consideration in keeping the population from obesity, but it is a feature which has not always been operative. At certain times in the world's history and in certain countries, extreme obesity in women has been regarded as attractive. Conversely, until recent years we have seen the younger females abolishing so far as possible every typical feminine curve. The explanation of this remarkable state of affairs is to be found in the increase of frankly or potentially

homosexual males; who arose from a childhood spent during the war, when they were subjected purely to feminine influence. We see now how cause and effect are on the wane.

The effect of obesity on fertility is more worthy of study. I have several patients whom I have assisted to normal outline whose temporary delight was marred by unexpected pregnancy. These women, having had one or two children, had become very fat and had gradually discarded contraceptive practice on the assumption that they could no longer bear children. The reduction of weight by dieting *alone* has restored fertility. In the male, impotence is more prone to arise.

Infiltration of the abdominal and perineal muscles with fat is responsible for the constipation and, in women, moderate incontinence of urine. The latter, in its turn, aggravates the existing intertrigo inherent in the flexures of every fat person. The bad stance induced by a pendulous abdomen commonly results in osteo-arthritis of the spine. The mere effect of excess weight and interference with the vascular supply of the joints by deposits of fat will induce osteo-arthritis of the knees in many cases. The alterations of the metabolism in the obese are still but imperfectly understood. It is certainly a fact that in the majority the basal metabolic rate is found to be normal, or even raised, provided the calculations take into account the increased body surface arising from the dead weight of fat in which metabolic processes are, to all intents and purposes, absent. A subnormal temperature is very common, and is quite compatible with a raised basal metabolic rate. From time to time outbursts of fever relieve the monotony as an expression of a spasmodic attempt on the part of the organism to destroy the material which is threatening its existence. Dyspepsia of a peculiar kind is to be observed in most of the gross obesities. It is characterized by a spurious appetite, by which I mean an inordinate desire to eat large quantities of food in an attempt to alleviate a sinking feeling in the epigastrium. As the meal progresses this sensation is overcome by drowsiness, and the victim is usually to be found snoring through a very liberal "forty winks" in the afternoon. If, added to all this, we recall the unpleasant body odour of the fat person, we need no more to convince us that obesity, if avoidable, is disgusting.

CAUSES OF OBESITY

It may be accepted as an axiom that fatness is due to excessive eating or to endocrine deficiency. Gluttony may, of course, be relative—that is to say, food intake has to be considered in relation to energy output. This is too obvious to need further stress. The excuses for over-indulgence are many and ingenious. The commonest is a denial that it exists. Nearly all fat people say they are small eaters, and often they are right. What they fail to recognize is that bulk and calorie value do not run parallel. It is worth while taking down from the patient's description a note of a day's food and analysing its energy equivalent. To find 500 calories in excess of requirements is quite common.

The necessity of keeping up strength, the sinking feeling without a meal, sleeplessness, or the remarks of neighbours that the patient is looking so ill—all or any of these may be used as an excuse for overfeeding. The most difficult of these arguments to overcome is that of sleeplessness.

* Address to the East Sussex Medico-Chirurgical Society, February 20th, 1934.

Prompted by posters of alluring beauties asleep with the aid of a hot drink of considerable food value, our patient, already disturbed by the indigestion of obesity, flies to the remedy and finds that it works—for a time. The sleep is induced, no doubt, by the diversion of blood to the splanchnic area, but in three hours the victim is wide awake again. What does she do? Reaches out for her thermos flask of hot milk, and so adds more fat to her greasy deposits.

Malt liquors are well known for their fattening properties, which are only slightly related to the trivial alcohol content. On the other hand, the avoidance of alcohol and smoking often drives the excessively righteous to an orgy of feeding which is more revolting than any vice he may have abandoned.

to function through glandular insufficiency. As an example, the photographs of a case of Frölich's syndrome are instructive.

This boy is aged 14, measures 5 ft. 10 in., and weighed $13\frac{1}{2}$ stone. The feminine distribution of fat is well seen on the buttocks and flanks. The distribution of pubic hair and tendency to knock-knee is likewise characteristic. The remarkable height of the boy is due to the disproportionate development of the lower limbs—namely, $38\frac{1}{2}$ in. as compared with $31\frac{1}{2}$ in. from pubes to the crown of the head. This feature is the result of testicular insufficiency secondary to pituitary failure. Note the typical venous stasis in the lower limbs. To give such a boy thyroid would be futile. The absence of thyroid involvement is always clearly shown by the normal shape of hands, fingers, and toes, the healthy condition of the hair, and the facies. Actually this purely glandular

FIG. 1.

FIG. 2.

January 26th, 1934

FIG. 3.

FIG. 4.

March 2nd, 1934. Note change in facial expression.

As to endocrine complicity in the development of obesity, my experience has been that this is present in some 20 per cent. of cases in clearly recognizable form. In a series of ninety-four cases seen privately, endocrine deficiency was present in eighteen, and of these there were ten examples of hypopituitarism, five of myxoedema, and three of gonadal insufficiency (two post-operative). Of twenty-eight hospital cases there were two examples of myxoedema, four of hypopituitarism (Frölich's syndrome), and the remainder were alimentary in origin.

It is particularly instructive to note that thyroid deficiency was unquestionably responsible for only seven out of 122 cases of obesity in view of the widespread and mistaken habit of prescribing thyroid extract to all fat patients. I quite realize that many of the other cases which I have classed as "alimentary obesity" had also some evidence of hypothyroidism, particularly those women whose pre-menopausal obesity increased rapidly with cessation of the periods. To these the administration of thyroid is justifiable, but it should be appreciated that by suitable dieting, if the case has not gone too far, all the symptoms can be abolished, and a moderately low basal metabolic rate may right itself in the process. This can never be said of true myxoedema.

It is unnecessary for me to describe in detail the characteristics of the various types of endocrine deficiency. In any case they are so often combined that it is impossible to be certain of the extent to which one of them contributes. Any hereditary factor which is present appears

obesity, with its associated drowsiness and lethargy, has been overcome in a few weeks solely as the result of dieting.

In hypopituitarism of later life the fat is deposited in masses on the body from the waist downwards, and often on the arms. Characteristically, the hands and feet are quite normal, as also are the forearms, ankles, and wrists. The face is not affected proportionately. The great increase in weight which affects many women after they have given birth to one child, and attacks a minority of patients after an acute illness or even operation, is presumably to be attributed to loss of activity of the anterior pituitary. The distribution of fat certainly favours this.

TREATMENT OF OBESITY

Whether endocrine or alimentary, the treatment of obesity is almost entirely a matter of dieting, with the one exception of the fatness of myxoedema. Preparations of pituitary, given by mouth and by injection for weeks on end, have proved valueless. Dr. P. Bishop, experimenting on rabbits with a well-known preparation of whole pituitary, has shown that it is in no way protected from digestion by the use of a keratin capsule, although the same material given intravenously displayed well-marked activity.

DIETETIC TREATMENT

What, then, is the principle of dietetic treatment? It is to give a diet of calorie value below that required for the optimum (not actual) weight of the patient. The

deficiency of calories is made up by the subject's using his own fat. It follows that all forms of obesity can be corrected by dieting. A gentle, but very efficacious diet, for which no calculation is essential, is as follows:*

On Waking.—A glass of water.

Breakfast.—Tea or coffee, sweetened with saccharine; one slice of dry toast, no butter; 3 oz. of cold tongue, boiled so'e, haddock, or whiting; fresh fruit.

1 p.m.—A tumbler of water.

Lunch.—4 oz. of chicken or meat (except pork), but no gravy; vegetables as given below, cooked without fat; salad without oil; one square of vita-weat; fresh fruit.

5 p.m.—Tea without milk or sugar.

Dinner.—Bouillon; fish; 2 to 3 oz. of game or meat (except pork), no bread-sauce or bread crumbs; vegetables as mentioned below; salads; one slice of toast; dry wine; fresh fruit; coffee with saccharine.

Condiments Allowed.—Worcester and anchovy sauces, ketchup, pepper, mustard, vinegar, walnut pickle, horse radish, salt sparingly.

Vegetables Allowed.—Green vegetables (except peas), celery, seakale, asparagus, and salsify.

Anything not mentioned above may be assumed to be forbidden. This is safe even for elderly subjects with cardiac weakness, although the heart should always be carefully watched during reduction of weight.

A more accurate and much more rapid treatment has been devised by Evans and Strang.¹ Their diet is built up around 1 gram of protein per kilo body weight. This is necessary to maintain nitrogen equilibrium, provided that some carbohydrate is given with it. Without carbohydrate some of the protein would be diverted for its antiketogenic properties. Thus, not less than 0.6 gram is given for each gram of protein. One of these authors' diets which has been quoted recently² allows about 8 calories per kilo instead of the normal 25 to 30 calories. It is very efficacious, though drastic.

Evans and Strang Diet (about 650 calories)

Breakfast.—One egg and 1 oz. of bread.

Lunch.—One egg and 4 oz. of vegetables as below.

Dinner.—One cup of bouillon and 3 oz. of lean meat; 4 oz. of vegetables as below.

Vegetables Allowed.—Lettuce, cucumber, spinach, asparagus, endive, celery, mushrooms, tomatoes, sprouts, watercress, cauliflower, radish, cabbage.

Water as much as needed; no fried foods; a teaspoonful of bicarbonate of soda daily.

It is essential to eat all the amount mentioned in this diet.

The tendency to acidosis is relieved by the use of bicarbonate. Many patients on this diet lose 1/2 to 1 lb.

* Modified diet from *Obesity*, Leonard Williams.

a day, but may feel listless and suffer constipation. This is the diet used for the patient illustrated above.

Kenyon's modification³ is a useful alternative, though capable of less accurate control. Its food value is approximately 1,000 calories. It allows of high protein, vitamin, and mineral salt content, and leaves the patient to consume his own fat.

Breakfast.—One portion of fruit; one egg and white of one egg; coffee, tea, bread substitute (for example, Heudebert's breadsticks).

Luncheon.—3 oz. lean meat, fish, or fowl; 1/2 pint 5 per cent. vegetables; one portion fruit.

Tea.—Tea.

Dinner.—3 oz. of meat, fish, or fowl and vegetables as at lunch.

9.30 p.m.—Half-cup orange juice; 1 oz. bemax.

In addition, 1½ glasses skimmed milk daily (for calcium content); one capsule haliverol t.i.d. (for vitamin content).

A fruit portion = one orange; half a grape-fruit; one medium apple; two medium peaches; one small pear; one cup strawberries; one cup blackberries.

When the desired weight is attained the diet should be slowly increased with vegetables, and later butter and bread. With the exercise of a little care, weight may be maintained at the ideal figure.

DRUG TREATMENT

Cutting, Mehrtens, and Tainter⁴ investigated the pharmacological action of a yellow dye, alpha-dinitrophenol, which had given rise to fatal poisoning in the munition factories of France during the war. They found that the metabolic rate of animals could be raised 50 per cent. by a dose of 10 mg. per kilo, and over fourfold by larger doses, which would produce death from hyperpyrexia. In human beings the drug in doses of 3 to 5 mg. per kilo daily promptly raised the basal metabolic rate by 40 per cent., and this was accompanied by an average loss of weight of 2 lb. a week. Dodds and Pope,⁵ working with dinitro-*o*-cresol (dekrysil), found that it would produce the same effect as dinitrophenol, but in appreciably smaller doses—namely, 1 mg. per kilo daily.

Following this publication I tried the drug* in various doses on ten patients. With 1 mg. per kilo body weight given thrice daily, the temperature rose to 101° in two days, and this was accompanied by tremor, tachycardia, and profound malaise. On the third or fourth day the patients (three in all) were yellow in colour, just as if they were jaundiced. This is presumably due to the drug, which is a yellow dye; there was no true jaundice. The pigmentation passed off completely in five days to three weeks; the symptoms subsided in two days after cessation of administration. The remaining seven patients were treated with 1 mg. doses per kilo body weight once a day. No general constitutional effect was noticeable, but weight was lost, if at all, very slowly—for example, 1/2 to 1 lb. weekly.

Judging by Cutting's experiments with the more toxic dinitrophenol given for three months in full doses to animals, it is probable that the regular administration of dinitro-*o*-cresol is a safe procedure, if given in small doses. Subsequent to my somewhat alarming experiments, an illuminating article by Dodds and Robertson⁶ appeared showing how impossible it was to judge the effect of the substance on the metabolism by temperature or pulse readings: only estimation of the basal metabolic rate can be of help. The metabolic rate can be raised 50 per cent. or more, and yet be unaccompanied by any appreciable alteration in the heart rate or body temperature.

It view of the fact that it is possible to control obesity even if of frankly endocrine origin by means of dieting

* Supplied by the courtesy of British Colloids Ltd.

alone, it is clearly undesirable to use a drug which is poisonous in doses only a little above those which are effective. In obstinate cases its use may be countenanced, but only when strictly controlled by the doctor, who must be guided by knowledge of the basal metabolic rate. If these capsules be given to patients in bulk it will not be long before one of them doubles or trebles the dose in the same light-hearted way with which thyroid is treated, and then we shall have further scares from the coroner's courts.

REFERENCES

- ¹ Evans, F. A., and Strang, J. M.: *Amer. Journ. Med. Sci.*, 1929, clxxvii, 339.
- ² Dodds, E. C.: *Practitioner*, 1934, cxxxii, 61.
- ³ Kenyon, J. H.: *Journ. Amer. Med. Assoc.*, 1933, ci, 97.
- ⁴ Cutting, W. C., Mehrtens, H. G., and Tainter, M. L.: *Ibid.*, 1933, ci, 193.
- ⁵ Dodds, E. C., and Pope, W. J.: *Lancet*, 1933, ii, 352.
- ⁶ Dodds, E. C., and Robertson, J. D.: *Ibid.*, 1933, ii, 1137.

CUTANEOUS SENSITIVITY TO ACID-FAST BACILLI IN SUSPENSION

BY

PROFESSOR S. L. CUMMINS, LL.D., M.D.

AND

ENID M. WILLIAMS, M.D.

TUBERCULOSIS DEPARTMENT, WELSH NATIONAL SCHOOL OF MEDICINE,
CARDIFF

Muir in India and Mitsuda and Hyashi in Japan have shown that, while advanced cases of nodular leprosy are negative to intradermal injections of "leprolin," a suspension of leprosy nodule in a state of fine division, the mode of preparation of which has been described in their articles on the subject, the more resistant types of anaesthetic lepers react positively to these injections. They report also that, while infants and young children are negative, healthy non-leprosy adults are usually positive to such injections, just as are the more favourable anaesthetic cases. Muir, being anxious to ascertain for certain whether healthy adults in countries where leprosy is no longer endemic reacted as do non-leprosy adults in India, suggested to one of us that he would be prepared to supply "leprolin" if these tests could be performed. We received from Calcutta a supply of Muir's "leprolin" early in October. This preparation, examined microscopically, was found to contain very large numbers of acid-fast bacilli irregularly distributed as single rods, small clumps, and large bundles. Tested for sterility, it gave no growth either aerobically or anaerobically, and a guinea-pig, into which 1 c.cm. was inoculated subcutaneously, has remained free from signs of infection to date.

RESULTS OF TESTS

Through the kindness of Dr. P. K. McCowan and his staff it was possible to carry out these tests on a group of twenty-five non-tuberculous male adults at the Cardiff City Mental Hospital, where successive series of tuberculin tests, designed to throw light on the degree of sensitivity of the population of a mental hospital, have been in progress for some years. As the tests proceeded it became clear that we ought to discover, as far as possible, how far the positive reactions noted were comparable, not only with tuberculin sensitivity, but with sensitivity to a suspension of killed tubercle bacilli, and our tests were accordingly supplemented to allow of this comparison.

On October 30th, 1933, each of the twenty-five men received in the skin of the right upper arm an intradermal injection of 0.1 c.cm. of "leprolin," diluted to 1 in 5 in sterile carbol

saline, and, at the same time, 0.1 c.cm. of a 1 in 2,000 dilution of old tuberculin in the skin of the left arm. The results were observed, and, where positive, measured in extent of raised palpable reaction area, on the third day and afterwards at weekly intervals up to the end of the sixth week.

All those tested gave the usual positive response to tuberculin, which reached its maximum in most cases on the third day, was still well marked, though fading, after one week, and, except in five cases, ceased to be measurable by the end of the second week.

With the "leprolin," thirteen cases were recorded as negative on the third day, and twelve gave more or less marked positive reactions, but the majority had become positive by the end of the first week, five only being recorded as negative or doubtful on the eighth day. All, with the exception of one patient, a Dane, who attained to a "doubtful" only on the eighth day, and was negative at all other dates, were "positive" sooner or later during the time of observation. Their reactions will be considered in greater detail later. The rapid disappearance of the well-marked reactions to old tuberculin, a filtrate containing the soluble elements of acid-fast organisms, as compared with the persistent though less extensive reactions following the inoculation of a suspension of whole leprosy bacilli, led us naturally to attempt a comparison between these latter and a suspension of unbroken tubercle bacilli, killed by heating, as had been the leprosy bacilli. To this end an egg culture of "human" tubercle bacilli, eleven days old, was suspended evenly in distilled water, counted, killed by heating, and diluted with carbol saline so as to contain 100 million bacilli in 1 c.cm. The sterilization was effected at a temperature of between 90° and 95° C. for one hour.

On October 20th, the remains of the tuberculin tests having by now practically disappeared, each of the twenty-five already tested were given an intracutaneous inoculation of 0.1 c.cm. of tubercle bacillary suspension, equivalent to 10 million dead bacilli, into the skin of the left upper arm, thus permitting a comparison with the leprolin reactions, many of which still persisted. As before, the results were read on the third day and at weekly intervals up to the end of the third week.

In examining cutaneous sensitivity to suspensions of acid-fast bacilli, the reactions, both to leprolin and to tubercle suspensions, were found to fall into three types, the distribution of these types, however, being very different in the tested group. These types are as follows:

- Type A.—"First high." Maximum reaction on third day.
Type B.—"First low." Maximum reaction on eighth day.
Type C.—"First negative." Negative on third day developing later.

The distribution of these types to leprolin and to tubercle bacilli respectively was as follows:

Reaction Type	Leprolin		Tubercle Bacilli	
	No.	%	No.	%
Type A	6	24	21	84
Type B	6	24	2	8
Type C	13	52	2	8

It will be noted that the great majority gave their maximum reaction to tubercle bacilli on the third day, just as they had done to old tuberculin; whereas the majority of the group were negative to leprolin on the third day.

The reaction size in millimetres—maximum transverse measurement—is given, as the average for those included in each type, on each day of observation up to the end of the third week, by which date all the reaction areas were rapidly diminishing in intensity.

Average Reaction Size in Millimetres

Reaction Type	Antigen	3rd Day	8th Day	15th Day	22nd Day	No.
Type A	Leprolin	9.7	7.5	4.3	4.3	6
	T. B.	17.6	9.6	7.7	4.2	21
Type B	Leprolin	6.8	9.5	6.4	6.0	6
	T. B.	9.0	13.0	12.0	8.5	2
Type C	Leprolin	0	3.2	3.5	3.8	13
	T. B.	0	5.2	6.0	5.2	2