and the scars remained free from relapses. In the rest of the cases great improvement was obtained. Only a few scattered nodules reappeared in a scar otherwise free from lupus. Probably a repetition of the light treatment, combined perhaps with electrocoagulation, will succeed in removing in these cases, too, the last remnant of the disease, as was in fact found with a number of other inveterate cases which had proved refractory to previous treatment with the original Finsen apparatus.

The two outstanding advantages of the Finsen treatment are its efficacy, and even more the ideal cosmetic results: soft, smooth, often hardly visible scars, which have preserved the full vitality and resistance of the natural skin without any disposition to atrophy, ulceration, or malignant degeneration.

THE EYES AS A CAUSE OF HEADACHE*

BY

ARTHUR GRIFFITH, M.B., F.R.C.S.

SENIOR SURGEON, ROYAL EYE HOSPITAL; SENIOR OPHTHALMIC
SURGEON, WESTMINSTER HOSPITAL

It is surprising to an ophthalmologist that physicians attach relatively so little importance to the eyes as a cause of headache. I have examined my notes of 1,000 private cases, and I find that 24 per cent. of the patients came to me complaining of headache. Not every headache, however, is due to use of the eyes, and the oculist, faced with a patient complaining of headache, must determine, if he can, whether the headache is an ocular one or not. In this he may be greatly helped by the history which he obtains from the patient.

First, as to the *situation* of the pain. Usually an ocular headache is frontal or orbital ("pain behind the eyes"); more rarely is it occipital or temporal, and almost never is it vertical. In the series of cases which I analysed, of 181 cases where the type of headache was noted 60 per cent. were frontal or orbital. Next the *time of onset* of the pain should be ascertained. Commonly with ocular headache the pain comes on in the late afternoon or early evening—that is, at the end of a day's work. Sometimes a headache is present on waking in the morning, and it may be difficult to persuade the patient that this morning headache has been earned by misuse of the eyes on the preceding day. Of course, other headaches may be earned overnight, such as that which comes out of a whisky bottle, but the patient always recognizes this type, and will say that if he ever gets a headache he knows to what to attribute it.

The delayed headache is probably due to the power of the healthy human being to ignore slight pain. It is only when he is tired, or when his control has been weakened by sleep, that he becomes conscious of pain. Sometimes patients will say that if, for any reason, they wake in the night, they find they have a headache. Another instance of delayed pain is the week-end headache; the patient may be free from pain during the week, but on Sunday, although using his eyes less, he gets headache. One patient said that he had not time for a headache during the week. Association of the pain with use of the eyes may be so clear as to make it reasonably certain that we are dealing with ocular headache.

Ciliary Muscle Contraction as Cause of Headache

Although the pain of eye headache may radiate through any of the branches of the trigeminal nerve, the initial

cause is painful contraction of muscle. The muscles concerned may be the ciliary muscles, the extrinsic eye muscles, or the occipito-frontalis. If one doubts that contraction of the ciliary muscle can be painful one should put strong eserine drops into an eye; the resulting pain may be severe enough to cause nausea.

The simplest case of pain from ciliary muscle contraction is that of a hypermetrope with equal errors in the two eyes. Such a patient can only see clearly by contracting the ciliary muscles, and throughout his waking day they are in a state of tonic contraction, the contraction being greater when close work is done. After prolonged close work these muscles tire, and the patient finds his sight varying, the type appearing alternately clear and confused. He may seek relief from this by stopping his work and gazing into the distance for some minutes, when he finds that he can continue to focus his work for a time. More frequently he discovers that by getting closer to his work, and so obtaining larger retinal images, he can appreciate it, in spite of confusion of vision, with less mental effort, but this diminution of mental effort is gained at the expense of accommodation and convergence, and a vicious circle is set up, which ends in headache or in breakdown of accommodation. Most children are guilty of the mental laziness which makes them read at too short a distance and so gain larger retinal images. The habit is accentuated by refraction errors and by imperfect lighting, and is one which should be combated constantly by teachers and parents.

When the hypermetropia is of different degrees in the two eyes the ciliary muscles are presented with a more difficult task. Not only must the patient accommodate in order to see clearly, but he must accommodate to a different extent in the two eyes. Patients will often tolerate a considerable under-correction of their hypermetropia or presbyopia better than a small imbalance of accommodative effort in the two eyes. In correcting refraction errors it is of the greatest importance to get the two eyes balancing in this respect.

In astigmatism there is introduced another abnormality of ciliary muscle contraction. If there is much astigmatism the ciliary muscle may make little or no effort to correct it, and there is no "eye-strain" and no headache. But if the degree of astigmatism is low or moderate the ciliary muscle contracts irregularly, more in some fibres than in those at right angles to them, and produces a lental astigmatism of opposite sense to the corneal astigmatism. Often enough the effort is overdone, and we find that a patient who, for instance, has a low degree of astigmatism "with the rule," chooses a cylinder "against the rule." Low degrees of astigmatism often cause more severe headache than do high degrees, although the importance of correcting very small refraction errors has been exaggerated.

More debatable is the question of "eye-strain" and headache being caused by contraction of the iris muscles. It is undoubted that the early cinema theatres were more productive of "cinema headache" than are the modern ones. Much of the improvement may be attributed to abolition of flicker and to reduction in the size of the screen, but another factor is that, whereas in modern cinema theatres a certain amount of general illumination is kept, in the old theatres the house was quite dark. Alternate gazing at the brilliantly lit screen and into surrounding blackness caused alternate contraction and dilatation of the pupils, and this may well have contributed to "cinema headache." In some of the newest cinemas the audience must traverse long corridors before reaching the theatre. One may be inclined to attribute this to love of display on the part of the management, or to ineptitude on the part of the architect, unless one observes that the illumination is gradually reduced as

* Read in opening a discussion in the Section of Ophthalmology at the Annual Meeting of the British Medical Association, Bournemouth, 1934.

one passes from the brilliantly lit foyer to the dimly lit theatre.

Before leaving the subject of ciliary muscle headaches I would refer to the fairly large group of composite headaches in which the ocular part is only a fraction of the whole cause.

As an instance, I would quote the case of a woman who consulted me for very severe and constant headache, which completely incapacitated her. She was taking enormous quantities of barbitone drugs with little relief. She had given up her work, and friends had taken her away on their yacht to try to improve her health, but she spent the whole time in bed, and returned as bad as ever. I corrected her refraction errors and ordered glasses. On wearing them she was free from headache for the unprecedented period of three weeks. Then began a clear association of the headache with her menstrual periods; it returned a day or two before the onset of the period, passed off, and returned for three or four days after the period. I sent the patient to a physician, who treated her with appropriate endocrine preparations, with the result that she has hardly had a headache in over five years.

Headaches Due to Extrinsic Muscle Abnormalities

Here in Bournemouth, where Maddox lived and worked for over thirty years, it is appropriate to discuss the use of prisms. Headache caused by imperfect muscle balance can often be relieved by the discriminating use of prisms, but few things require greater judgement than their prescription. Even more than in correcting refraction errors must one avoid the error of over-correction. It is better not to order prisms than to order too strong ones. Not always can the same prisms be worn for distance and for near work. An esotrope with muscle imbalance sufficient to cause confusion and even diplopia in the distance may find his latent convergence rather a help in reading. Such a person, however, is liable to fall into the habit—a habit which is responsible for much ocular headache—of reading at too short a distance, and so overtaxing his accommodation.

The movements of elevation and depression of the eyes are performed by a mechanism more complex than that used for the lateral movements, and it is perhaps for this reason that vertical heterophorias, although less common than lateral imbalance, are more productive of discomfort and headache; it is especially these vertical errors which call for relief by prisms. As a rule, it is the less good eye which tends to deviate upwards, probably because the elevating muscles are stronger than the depressors. The factor which decides which eye shall tend to deviate is usually the refraction, the eye with the more complicated refraction error tending to rise. When there is little difference in refraction the hyperphoric eye is usually that which is not the "master eye": one might call it the "servant eye." We are therefore correct in speaking, as we usually do, of a right or left hyperphoria, and not of hypophoria.

As rough rules for the prescription of prisms for heterophoria Maddox suggested as maximal corrections two-thirds of the error in esophoria, one-half in exophoria, and two-thirds in hyperphoria. There can, however, be no fixed rules, and each case must be considered separately. A plan which I have found of the greatest value is to make the patient gaze at a distant spot of light with coloured glasses before his eyes, red glass before the right eye and green before the left. If he says, whatever the finding with the Maddox rod may have been, that the red and green lights are mixed together, his heterophoria does not need correction by prisms. If, on the other hand, he sees the red and the green lights separate, one must find the weakest prism which will enable him to mix them. This prism is to be regarded as a maximum, which is not to be exceeded in the prescription. It may even be found that some still weaker prism will give the patient comfort when reading the test types with

both eyes, making the letters appear steady instead of dancing and confused. In prescribing prisms, as in correcting refraction errors, over-correction must be avoided.

It is said that the prescription of prisms for heterophoria is purely palliative and not curative, but this is not always true. It is true that patients who have been relieved by wearing prisms may, when they return a year or so later, be found to have more heterophoria than before, but one finds at least as frequently that the tired muscles have recovered some of their tone, and that we can reduce or abandon the prisms in the glasses.

Occipital Headache

The last group of eye headaches to which I will refer includes those which can often be relieved by the volition of the patient. Many patients suffering from eye-strain screw their lids together. This may be done, as in cases of myopia, to cut off diffusion circles by narrowing the palpebral fissures, but in other cases it would seem that the excessive innervation of the eye muscles overflows into the orbicularis muscles. The frontales muscles are inserted into the upper lids, and they participate in the contraction, and as the frontales, on contracting, pull on the epicranial fascia the occipital muscles are pulled upon, and these, too, contract and cause occipital pain. This is not the only, nor perhaps the commonest, form of occipital headache. We shall hear from the neurologists of occipital pain arising in the meningeal branches of the trigeminal nerve and associated with changes in the meningeal vessels or with alterations in intracranial pressure. In a recent case of Lindau's disease, in which I had been observing the angiomatosis of the retina, the first and only indication that the patient had developed a haemangioma of the cerebellum was the sudden appearance of severe occipital neuralgia.

The occipital headache of eye-strain, however, seems clearly to be localized to the occipital muscles, and to be relieved by pressing on them. It is a form of headache which is easily relieved by wearing suitable glasses. Much may be done for the patient by explaining to him that he is himself producing headache by screwing his lids and frowning, and that he must make himself conscious of the occasions on which he does these things and decide not to do them.

I have intentionally omitted reference to the headache of gross eye disease, such as iritis and glaucoma, of nasal sinusitis, and also of nephritis, cerebral tumour, and hyperpiesis, diseases in which ophthalmologists are interested by reason of the fundus changes which occur in them. I have confined myself to consideration of the headaches caused by misuse of the eyes, and of the ways in which we can relieve them.

A memorial fund to the late Madame Marie Curie has been opened by the hospital in Hampstead which bears her name and operates as a centre for the radium treatment of cancer in women. The appeal is being simultaneously launched in Paris and Warsaw. It has been thought fitting that the money thus received should be most appropriately expended on extension of this hospital in which she was so deeply interested. A department for deep x-ray therapy has recently been added to supplement radium treatment and for certain cases which respond better to x rays alone. Already it is overtaxed, and many would-be patients have to be turned away owing to the lack of facilities. The growing number of applicants, and the fact that for most of these radiotherapy offers the only prospect of relief and cure, make it urgent that the contemplated extension should be undertaken as soon as possible, especially since the establishment of this institution was one of the tributes this great investigator appreciated most highly during her life. Contributions may be forwarded to the treasurer, Marie Curie Hospital, 2, Fitzjohn's Avenue, N.W.3.