On October 19th the temperature was settling, the faucial oedema had gone (petechiae showed on the mucosa), and there remained a diffuse, hard swelling under the upper third of the right sternomastoid muscle. On October 20th the spleen was felt. On October 21st the temperature rose to 101°, the glands on the left side of the neck became more swollen and tender, and a general macular erythema appeared. On the 23rd earache started, the temperature rose to 103°, and the membrana tympani was found to be thickened, red, and pulsating. Next day the temperature was still high; paracentesis of the membrane was performed, with liberation of pus and relief of pain. This was followed by a gradual settlement of the temperature, and convalescence proceeded uneventfully, but for a persistently high pulse rate and occasional rises of temperature to 99–100°.

Case 3 (aged 13)

On October 8th this patient went to bed with a feverish cold. On October 11th his temperature rose to 102° and the right submaxillary gland was found to be swollen and tender. On October 13th he vomited and complained of dysphagia, and the glands in the posterior triangles of the neck were enlarged and tender. On October 16th the fauces were swollen, but not septic, the glands large and tender on both sides of the neck. Two days later he complained of headache and pain low down in the throat. On October 20th

CHART III: CASE 3.

he was more comfortable, but the glands were still tender and the temperature irregular. On October 23rd there was pain in the upper part of the chest and a sense of constriction, which was thought to be due to enlargement of the glands in the mediastinum; this subsided in forty-eight hours. The temperature settled, but for minor irregularities, and no further symptoms of note developed, apart from anaemia. A differential white blood count on October 30th gave the following: polymorphs, 48 per cent.; small mononuclears, 47 per cent.; large mononuclears, 5 per cent.

A fourth boy, who ran a very mild course for a fortnight, relapsed on the fourteenth day, and had a much more severe bout, with high fever, dysphagia, and glandular swelling lasting a further ten days.

Treatment.—This was necessarily empirical. Phenazone and quinine were tried without any obvious effect. Antiphlogistine externally and throat sprays of hydrarg. perchlor. 1 in 1,000 and of adrenaline 1 in 1,000 seemed to give relief.

SUMMARY

Certain of the cases described showed a very acute onset, a marked degree of periadenitis, and oedema of the palate and fauces, leading in one instance to laryngeal stridor. Unusual complications arose in the shape of an acute retropharyngeal abscess, and a cervical abscess (in a case not described in detail).

I am deeply grateful to Dr. F. J. Poynton for his advice on the outbreak in general, and to Dr. G. F. Still for his help and interest in one particular case.

REFERENCE

- ¹ *Proc. Roy. Soc. Med.*, 1931–2, 155; *New England Journ. Med.*, 1931, ccv, 1238; *Edin. Med. Journ.*, 1932, xxxix.

ON THE DIAGNOSTIC SIGNIFICANCE OF A POSITIVE SPUTUM REPORT

(DIRECT EXAMINATION FOR TUBERCLE BACILLI)

BY

G. GREGORY KAYNE, M.D., M.R.C.P.

DOROTHY TEMPLE CROSS RESEARCH FELLOW IN TUBERCULOSIS

A great deal of attention has lately been focused on the neglect of the early diagnosis of pulmonary tuberculosis. It may not, therefore, be untimely to stress a certain aspect of the *wrong* diagnosis of pulmonary tuberculosis, which has not received the notice it deserves in view of the important social and economic consequences to the individual concerned.

The student is rightly taught that one or even several negative sputum examinations do not justify the rejection of a diagnosis of pulmonary tuberculosis; but neither he, the general practitioner, nor the tuberculosis officer has his attention sufficiently well drawn, either in textbooks or post-graduate lectures, to the fact that a single positive sputum report by no means always establishes the diagnosis. As clinical assistant to the tuberculosis dispensary at the Hospital for Consumption, Brompton, I saw several patients sent by their medical attendant with the diagnosis of pulmonary tuberculosis based on a single report of "T.B. found in the sputum," in whom subsequent physical examination, x-ray of the chest, repeated sputum examinations, and prolonged observation (including a stay in hospital) showed no evidence of active disease. The reports of the sputum examinations emanated from a pathological laboratory of high repute, and the patients had been promptly notified by their private practitioners.

It is unnecessary here to emphasize the significance of such a label to the patient. But the position of the tuberculosis officer also is not an enviable one. He must continue to keep the patient under observation for several months at least. If at the end of this period he is still convinced that the patient has no active pulmonary tuberculosis, he can hardly remove him from the dispensary register as "cured"! Two alternatives lie before him.

(a) He can himself denotify the patient as "notified in error," or as "diagnosis not confirmed." Such action requires some courage, for occasionally a patient will develop pulmonary tuberculosis shortly afterwards, purely as a coincidence, and the tuberculosis officer will undoubtedly be blamed as having mishandled the case.

Thus Rist¹ found that of 192 patients, representing a consecutive series of hospital nurses (see below), referred to him at the Dispensaire Léon Bourgeois of the Laennec Hospital as diagnosed or suspected cases of pulmonary tuberculosis, and discharged by him as non-tuberculous, after sufficiently detailed investigation to ensure that no error had occurred, two developed active pulmonary tuberculosis within six months and one year respectively.

(b) The tuberculosis officer can communicate with the patient's private practitioner, giving his findings and suggesting that the latter denotify the patient. This procedure will, however, prove awkward for the doctor, who will probably be accused by the patient of bungling, and will certainly be blamed for the great inconvenience caused by the notification.

Rist discussed this subject at some length in a recent post-graduate lecture in Paris. Since 1923 he has been chief medical referee for tuberculosis to all the nursing staff of the hospitals of the Assistance Publique. While the latter correspond to the municipal hospitals now under the L.C.C. they actually include nearly all the hospital beds available in Paris, as voluntary hospitals consist of only a few religious institutions. All staff diagnosed as, or suspected of, pulmonary

tuberculosis are referred to him at the Laennec Hospital by the general medical referees. In 1925 he published¹ the results of the first year's work of this department. Of ninety nurses referred to him as cases of pulmonary tuberculosis with a report of tubercle bacilli in the sputum the diagnosis was not confirmed in twelve (13.4 per cent.).

SOURCES OF ERROR

The reasons for the findings quoted in the preceding paragraphs may be grouped as follows:

1. *Errors of Technique*.—These have been fully dealt with by Wilson² in a recent report from the aspect of research work. In carrying out routine sputum examinations the following causes of error must be considered.

(a) Mistakes in labelling of specimen jars, slides, and reports.

(b) Using slides which have been previously employed for making smears containing bacilli.

(c) Drying smears with blotting paper previously used.

(d) Employing tap-water for diluting stains or washing the slides. (Tap-water frequently contains non-pathogenic acid-fast bacilli.^{3 7})

(e) Using contaminated cedar-wood oil and bottles of stains.

(f) Occasionally mistaking artefacts due to scratches on slides, or crystals of stain, for bacilli.

The possibility of some of these errors should particularly be kept in mind when only an occasional bacillus appears to be present.

2. *Dishonesty*.—The sale of bacilli-laden sputum unfortunately proved not un lucrative during the war, and one can postulate circumstances even in civil life when such means might be adopted. In the Assistance Publique of Paris a nurse automatically receives leave on full pay, for at least two years, immediately the diagnosis of pulmonary tuberculosis is made, and temptation may therefore sometimes prove too great. Further, the increasing number of pathological "services," some with inadequately trained staff, may be made use of by charlatans requiring some justification for the prolonged and costly treatment advised to the patient.

3. *Tuberculous Staff*.—A laboratory attendant suffering from open pulmonary tuberculosis may contaminate material in a laboratory for some time before his condition is detected.

4. *Other Acid-fast Organisms*.—Finally, attention must be drawn to recent papers which have described the occurrence in man of other acid-fast organisms. Branch⁴ reviews the question and describes the occurrence of non-mammalian tubercle bacilli (some of them belonging to the avirulent avian group) in the skin, pleural fluid, and blood cultures. Cummins and Williams⁵ describe an acid-fast bacillus ("M" bacillus) found in the sputum of a female patient suffering from severe pulmonary disease and achalasia of the cardia; they were unable to decide whether it was an adventitious saprophyte or the causative organism of the disease. Baldwin⁶ later reported a similar case—that of a young female patient suffering from a phthisis-like illness and achalasia of the cardia, in whom a bacillus was isolated, resembling the strain of "M" bacillus: his patient, like that of Cummins and Williams, was completely restored to health. Of special interest is the description by Neisser⁷ of acid-fast bacilli found on practically all metal musical instruments, but less frequently on wooden ones.

CONCLUSIONS

The above considerations prompt the following practical conclusions:

1. A diagnosis of pulmonary tuberculosis based on a single positive direct sputum examination without any other evidence is not justified.

2. In any case, whenever possible, more than one sputum examination should be carried out.

3. In view of Cummins and Williams's and Baldwin's cases it would appear that "even when the clinical picture is convincing, the finding of acid-fast bacilli in the sputum is not necessarily proof-positive of tuberculosis. The importance of cultural verification in all doubtful cases cannot be too much stressed" (Cummins and Williams).

4. When there is any question of compensation, pension, etc., unless the clinical picture is quite clear, not only should more than one sputum examination be carried out, but the patient should be made to expectorate in the presence of the physician or pathologist, and, if necessary, after isolation in a hospital ward under observation.

5. It would be well if the practice were universally adopted of indicating in every sputum report an approximate estimate of the number of tubercle bacilli present—for example, twelve per field, one per four fields, etc.

REFERENCES

- ¹ Rist, E.: *Rev. de Phthisiol.*, 1925, vi, 80.
- ² Wilson, G. S.: Medical Research Council Special Report No. 182, p. 97.
- ³ Schwabacher, H.: *Ibid.*, pp. 104 and 124.
- ⁴ Branch, A.: *Tubercle*, 1933, xiv, 337.
- ⁵ Cummins, S. L., and Williams, E. M.: *Ibid.*, 1933, xv, 49.
- ⁶ Baldwin, E. R.: *Ibid.*, 1934, xv, 236.
- ⁷ Neisser, M.: *Journ. State Med.*, 1933, xli, 63.

Clinical Memoranda

SEVERE HAEMORRHAGE WITH SPONTANEOUS EVACUATION OF PERITONSILLAR ABSCESS: LIGATION OF COMMON CAROTID— RECOVERY

I have read Dr. P. Shackleton's two cases of pharyngeal haemorrhage, published in the *Journal* of December 23rd, 1933, with much interest. Severe haemorrhage from a tonsil abscess is, as Mr. E. D. D. Davis remarks,¹ rare, and I think the following case should be added to those already on record.

CASE REPORT

The patient, a sturdy, full-blooded brewery labourer, now aged 46, gave a history of having been troubled by tonsillitis all his life. On April 3rd, 1929, he retired to bed with "influenza" and a sore throat. The sore throat settled on the right side, and four days later an "abscess on the right tonsil" burst, giving him considerable relief. At 1 p.m. on the following day he felt a "hot, tingling sensation" in the back of his throat, and suddenly found his mouth full of blood, which, he said, he had to spit out quickly, otherwise he would have choked. This bleeding stopped in a quarter of an hour, but recurred alarmingly a few hours later, when Dr. A. T. McKay had him admitted to Mercer's Hospital under the care of Sir John Lumsden and myself.

On admission the haemorrhage ceased for a day, but the next morning the patient suddenly spat up a kidney-trayful of blood. Examination of the throat at this stage revealed a resolving quinsy, and I tied off what I thought to be a bleeding point low down on the posterior pillar. This had no effect, however, and haemorrhage continued at intervals throughout the following day, in spite of all our efforts to control it with tonsil clamps, haemostatic injections, and other agents. On the evening of the third day after admission the patient had become very bloodless indeed, and as his condition was obviously serious it was decided to tie the right common carotid artery. This was accordingly done under very light ether anaesthesia. Just before the anaesthetic was started the patient vomited up two pints of fresh clotted blood. This very nearly proved the last straw, for although blood transfusion was performed as soon as possible after operation, and again after an interval of two days, the patient's condition continued to be very critical indeed for several days. He told me afterwards that during this period if either arm fell out sideways from the bed he had not the strength to move it back again. He noticed this particularly