DIVERTICULITIS: A CLINICAL REVIEW

BY

HAROLD C. EDWARDS, M.S., F.R.C.S.

HUNTERIAN PROFESSOR AND JACKSONIAN ESSAYIST, ROYAL COLLEGE OF SURGEONS OF ENGLAND;
SURGEON TO KING'S COLLEGE HOSPITAL, LONDON

The circumstances of the anatomy and position of colonic diverticula predispose them to secondary pathological processes. They are bottle-necked, thin-walled sacs opening from a muscular tube, which contains faecal material rich in micro-organisms. The initiation of secondary changes depends entirely upon the retention of faecal matter within the diverticula, and it is clear that the more solid the stage which the faecal material has reached, the greater tendency there will be to retention. Hence the sigmoid colon is easily the commonest site for diverticulitis, not only because it is the commonest situation for the formation of diverticula, but because its contents are solid.

The presence of a faecal concretion may lead to a variety of secondary inflammatory changes, which are set out from the clinical standpoint in the accompanying scheme.

These changes vary from acute infection of the diverticulum, with perforation into the general peritoneal cavity, to a chronic reactionary fibrosis of the wall of the colon, extending in some cases to beyond the peritoneal covering. To such a condition the term "perisigmoiditis" is applied. Such a mass of inflammatory tissue may ultimately produce stenosis of the bowel simulating carcinoma, or cause adhesions between the colon and neighbouring organs.

Should acute diverticulitis supervene in a diverticulum walled off by fibrous tissue immediate perforation into the peritoneal cavity is improbable, and a localized abscess may form in the mass of fibrous tissue surrounding the colon. An abscess of large size may eventually rupture into a neighbouring hollow organ, such as the bladder, or into the peritoneum. Not infrequently an abscess of this kind will track along the side of the bowel to come to the surface under the parietes in the left iliac fossa.

The formation of intracolic and pericolic fibrous tissue may be due, in part, to the mechanical irritation of the faecal mass, but the main factor is probably infection from the lumen. This is due to the filtering of toxins or actual organisms through the damaged mucosa, which is stretched over the retained mass.

The Clinical Aspects

One hundred and forty-six cases, from various sources, of diverticula of the colon were investigated from the clinical standpoint. Of these, sixteen were cases of fistula into the bladder, the notes of which were loaned to me, and these will be excluded from the series for purposes of statistics. Of the remaining 130 cases, complications, due to the diverticula, occurred in nineteen, or 14.6 per cent. of the total. The complications were as follows:

	No. of Cases
Obstruction, large bowel	6
" " small bowel, from adhesions ...	1
Acute perforation	4
Abscess	3
Abscess with secondary perforation into the peritoneum	2
Fistula into the bladder	2
Fistula into the bladder and abscess in the pelvis	1
Total	19

This leaves 111 uncomplicated cases. In thirty-three of these other conditions were present, which were responsible for the symptoms either wholly or in part; thus seventy-eight cases of uncomplicated diverticulitis remain.

Uncomplicated Diverticulitis

Under this heading is included chronic diverticulitis with and without acute or subacute exacerbations, and this description is based on the seventy-eight cases in which neither complications nor associated abdominal disease were present. Though all these patients had diverticula of the colon, there was some difficulty in estimating the number who were suffering from diverticulitis as opposed to diverticulosis. Knowing that diverticula are present, by what criteria must one judge the presence of inflammation in them? In the more severe cases—those who are subject to acute or subacute exacerbations, or who have a palpable tender swelling—no doubt will exist, but in the greater number the symptoms are vague. While it is true that inflammation of mild degree may give rise to pain in the left iliac fossa, one must not lose sight of the fact that pain may arise from that disordered action of the bowel of which diverticula are the outcome.

The physical signs of thickening of the colon felt per abdomen must be accepted as evidence in favour of inflammation, but many of the patients are so obese that any such thickening cannot be felt. Mere tenderness over the colon is not an absolute sign, as it is frequently present in spastic constipation when x-ray examination

has proved the absence of diverticula. Involuntary spasm of the abdominal muscles is an absolute sign of subjacent inflammation; but if the colon is tender to palpation voluntary rigidity is also present, and may in some cases be very difficult to distinguish from reflex rigidity. X-ray examination may demonstrate spasm of the colon, but this may not be the result of inflammation in the diverticula; it may be the irregular action which is the prelude to their formation.

It will be conceded, therefore, that on the history, the physical examination, and the x-ray findings, it may be impossible to distinguish diverticulitis from diverticulosis, except in the severer cases, and I have failed in my attempt to estimate the percentage of cases of diverticulosis in which inflammation occurs. Rankin and Brown, in 1,300 cases in which diverticula were found in the colon, state that 17 per cent. were suffering from diverticulitis. If pain is regarded as a criterion of inflammation, then the following analysis of the symptoms will show that in the present series the percentage incidence is infinitely greater than this.

Analysis of Symptoms

In sixty-nine of the seventy-eight cases there were symptoms referable to the large intestine. The most constant of these was pain, which was present in fifty-four. This varied considerably in character; in some it was a dull ache, and in others it came in colicky attacks. Sometimes it was brought on by attempts at defaecation, but in others it was unaffected by this act. In forty-one of the fifty-four cases it was situated in the left iliac fossa. Other distributions were as follows: right iliac fossa, five cases; lower abdomen, four; umbilicus, one; and epigastrium, five, in two of which there was also pain in the left iliac fossa, and in the other three it was related to food. The site of the pain did not always correspond with the segment of the colon most affected by diverticulosis, as indicated by x-ray examination. In fifteen cases pain in the back was also complained of, but in the majority of these it may have been the result simply of constipation. There was no actual pain in fifteen cases, but the patients complained of disturbances of a more vague character, chiefly on the left side of the abdomen. These disturbances were variously described as rumbling noises, windy spasms, flatulence, soreness, and fullness of the stomach, and were felt more during defaecation, or when the patient was more than ordinarily constipated.

Constipation.—The bowel history was known in seventy-one of the seventy-eight cases. Only nine patients claimed to be normal in their bowel action. Forty-eight were constipated, and in thirty-three of these the constipation was of long standing and was severe. In thirteen it was of recent origin, and in two it was stated to be slight. There was a definite history of irregularity in six. The bowels were loose in three, and in five there was constipation alternating with diarrhoea; in three of these the bowel had previously been constipated. The latter condition must be regarded as significant of obstruction, but this is not always present, for the attacks of diarrhoea sometimes coincide with, and are due to, recurrent exacerbations of inflammation.

Blood and Mucus.—These were present together in three cases, and in three there was blood alone. Of these, one had severe cardiac disease and another had piles of moderate severity, but in the remainder no other cause for bleeding could be found. In a further series of twelve cases the test for occult blood was negative. Bleeding should not, therefore, be regarded as a characteristic of the condition. In the rare cases in which it is present it is probably due to associated colitis. Mucus was passed with stools in nine cases; this is a common accompaniment of severe constipation, however, and is not of special significance.

Bladder Symptoms.—Symptoms arising from the bladder were present in eight cases. Increased frequency and pain on micturition either together or alone were complained of on seven occasions, and in one case there was incontinence. It is difficult to say how far these urinary symptoms were directly due to the diverticula. In one case the patient

complained of frequency, pain, and a little difficulty. The urine was foul. The prostate was not enlarged, and the symptoms suggested that the formation of a fistula from the colon into the bladder was imminent. In two other cases the urinary symptoms were increased during attacks of abdominal pain.

Physical Signs.—Of sixty-four patients, thirty, or less than half, were obese. Of the remainder, twenty-five were regarded as of medium build, and nine as thin. In forty-four of the seventy-eight cases abdominal tenderness was present, in most instances in the left iliac fossa. This varied considerably in degree, from pain on deep palpation to severe pain on light palpation, with spasm of the overlying abdominal muscles. In twenty-four cases the sigmoid colon could be felt, and in five of these it appeared to be thickened. In eleven cases a definite lump was present, but in not all of these was the lump due to inflammatory thickening. In one patient, for example, a very large mass could be felt under the liver on the right side. It was diagnosed as a growth at first, but it disappeared with enemata, and subsequent x-ray examination showed diverticula in the distal part of the transverse and the sigmoid colon.

Clinical Diagnosis

Though the above analysis suggests that the clinical findings are for the most part vague, the diagnosis of diverticulitis is by no means a difficult one to make on clinical examination. The symptoms, taken in sequence as has been done above, do not furnish the true clinical picture, which is a characteristic one. Study of the case records shows the justification for this statement. A patient over the age of 45 complains of mild attacks of pain situated chiefly in the left iliac fossa. During the past few years he has noticed an increasing tendency to constipation. He feels a fullness on the left side, and puts his hand over the left iliac fossa and complains of tenderness there. The similarity to early malignant disease is patent, and one is not justified in accepting the clinical diagnosis of diverticulitis without excluding the possibility of growth—which not infrequently coexists—by x-ray examination.

Diverticula can occasionally be seen through the sigmoidoscope. In eleven examinations they were seen on two occasions. Sigmoidoscopy cannot be relied upon to any great extent for the diagnosis, however, which is far more readily made, and with less discomfort to the patient, by radiography.

Treatment of Uncomplicated Cases

The main treatment of uncomplicated cases of chronic diverticulitis is to avoid complications. Colonic lavage should be insisted upon, and every hospital which deals with this type of case should be equipped with facilities for this treatment in the out-patient department. The object is to wash away any retained products in the colon, and not to disinfect it, and it therefore matters little what lotion is used. Ordinary tapwater is probably as useful as any antiseptic solution, though one would not dispute the value of the natural waters at Bath and other spas.

The wash-out should be at body temperature, and the two details which *must* be insisted upon are: (1) not more than two pints should be used; and (2) the head of water should not exceed eighteen inches. This errs, perhaps, on the conservative side, but no one will deny the power for harm that lies in over-distension of the colon, especially the colon which has departed from the normal.

The diet should be mainly vegetarian; it need not be strictly so, but excess of cellulose should be avoided. The latter is an important item. The colon is not fitted to be a receptacle of the quantity of cellulose recommended by faddist societies, and to load the inflamed colon with undigested cellulose is clearly not to be desired.

Operative Treatment

The role of the surgeon in the treatment of diverticulitis is largely confined to the treatment of its complications. The nature of the disease and its irresistible tendency to adhesions make cases where local excision is possible very few and far between. In this series a radical operation was performed in five cases: there were three resections of gut, and in two the Mikulicz operation was performed. In only one case—a comparatively mild one—was the procedure limited to one operation. Though this patient remains in reasonable health to-day, six years after the operation, I regard the operation, in the light of additional experience, as unjustifiable, and a case of this low degree of diverticulitis I should now treat by medical means. The other four cases were progressing dangerously towards one of the complications, and it is only in such cases—one might almost say at this stage of the disease—that operation is justifiable. The difficulties that may be encountered, which make the decision to operate a weighty one, are well illustrated by one of my cases. The patient is now alive and well, but he underwent no fewer than five operations. The condition encountered at laparotomy did not justify an attempt to excise the mass, and hence the peritoneum on the left of the intestine was incised and the affected bowel mobilized and brought on to the surface of the abdomen.

Though the technique is difficult, when circumstances make it possible—that is, when the condition is localized—there is no doubt that the radical procedure in these borderline cases is the best line of treatment that can be offered. The alternative is an attempt to rest the diseased portion of the bowel by simple colostomy, with a promise to the patient that this will be closed at a later date. Such a method, however, not infrequently falls short of expectation.

Temporary colostomy—merely creating a fistula between the bowel and the abdomen—is doomed to failure. Such an operation only partially drains the bowel and does not allow of that degree of rest which is a *sine qua non* of recovery of the inflamed portion. The same applies with greater force to caecostomy, which is to be condemned except as a possible prelude to the radical operation. Apart from the distress it causes the patient, caecostomy gives only very imperfect drainage to the colon, and has a very limited effect upon the inflamed distal portion. This is clearly shown in one patient who suffered an acute recurrent attack of diverticulitis fourteen days after caecostomy had been performed as a preliminary to the radical operation.

It is essential, therefore, for the colostomy to be of permanent type if the patient is to derive any benefit. The question will arise, How long should the colostomy be allowed to remain? The question is unanswerable. There is no guarantee that symptoms will not recur after the colostomy has been closed by operation, even though all symptoms have subsided. A minimum period of at least twelve months must be insisted upon. The closure of a permanent colostomy is not unattended by risk to the patient, even if one of the ingenious extraperitoneal methods are used. And for such an operation, successfully achieved, to be followed at a later date by a recurrence of symptoms is a grave disappointment to both patient and surgeon. If such a recurrence is severe, and the danger of complication appears imminent, the only operation possible is a second colostomy, for the previous operation would in all probability make the task of excision a difficult one.

The following guiding lines for treatment are therefore submitted: (1) Uncomplicated mild diverticulitis—medical treatment. (2) Severe recurrent diverticulitis, in which complications appear to be imminent—laparotomy with a

view to determining the possibilities of excision, and if this be thought possible, a preliminary transverse colostomy or caecostomy. Operation should be postponed to an interval between the attacks. If the condition met with appears unsuited to radical treatment, a permanent type colostomy, made *as near as possible* to the inflamed area, and left open for a minimum period of twelve months.

Diverticulitis with Obstruction

Six cases of chronic fibrous obstruction of the large bowel due to diverticulitis appear in my series. The complication is therefore not uncommon. The similarity, both in history and in physical signs, and even on naked-eye examination, between this condition and carcinoma of the bowel is well and widely recognized. In one case the nature of the condition was only revealed by microscopical examination after the mass had been resected at operation, which was successfully performed in one stage. In a second case the patient had been admitted a year previously and treated by colon wash-outs. She was well on discharge, but neglected to continue with the treatment, and was readmitted with acute obstruction. A Mikulicz many-stage operation was satisfactorily performed. The sigmoid colon had become obstructed in a third case by a band passing from an inflamed diverticulum to the left Fallopian tube, while in a fourth a caecostomy had been performed, but, two years later, x-ray examination demonstrated that the condition of the colon had not improved sufficiently to warrant closure. In two further cases the patients, who were both over 70, refused operation.

The difficulty in distinguishing, both clinically and at operation, between obstruction due to diverticulitis and that due to growth makes it imperative to remove the mass, either by resection or by the Mikulicz method. Radiography cannot always be relied on to differentiate between them, although the method of dual exposure, devised by my colleague Dr. Graham Hodgson, may be of help in many cases. The presence of diverticula elsewhere in the colon is no criterion whatsoever that the mass is inflammatory, as diverticula and growth frequently coexist. (Six cases out of 130.)

Obstruction of the small intestine as a result of adhesions to a portion of the sigmoid affected by the diverticulitis is perhaps the most serious of the complications. It was accompanied in one case by chronic obstruction of the colon itself. The left iliac fossa was filled with a mass of fibrous tissue, to which a coil of the ileum had become firmly adherent. The ileum above the obstruction was dilated and its wall thickened. The surgical treatment here is a problem of considerable difficulty. It is manifestly useless to attempt to separate the adherent ileum from the colon, for recurrence is certain, and the obstruction must be short-circuited either by ileo-ileostomy or ileo-transverse-colostomy. In addition the obstruction to the colon must be treated. In the case in the present series the ileal obstruction was a chronic one situated about three feet from the ileocaecal valve. An ileo-ileostomy was made around the obstruction, and a transverse colostomy performed. The patient died three days later from ileus.

Acute Perforative Diverticulitis

This complication is, fortunately, rarer than might be expected. Diverticulitis is usually an insidious disease, and the diverticula are walled off by the formation of fibrous tissue. It occurred in four cases in the present series.

The history and pathological process are similar to that of perforation of the appendix into the general peritoneal

cavity, except, of course, that the early pain and the physical signs are on the left side of the body. There is usually a history of vague pain preceding the acute onset, but on occasion this is absent, the acute condition appearing with dramatic suddenness. Perforation may follow some strain, or occur during a prolonged attempt to defaecate. An important sign not obtained in perforation of the appendix is an early distension of the abdomen, with a tympanitic note on percussion due to the presence of gas.

It goes without saying that immediate operation should be performed in these cases. The conduct of the operation, once the abdomen is opened, will depend entirely upon whether the perforated diverticulum can be found or not. In one case this was possible. The diverticulum, which was at the upper end of the rectum, was excised, the opening in the bowel closed by a purse-string suture, and a temporary caecostomy used to relieve the distension of the large bowel. A drainage tube was placed down to the site of the opening into the bowel.

If the site of perforation cannot be determined, or the opening be buried in a mass of fibrous tissue so that it cannot be closed by any plastic operation upon the bowel, the great omentum is brought down and fixed over the area with one or two stitches, and a colostomy performed as close to the area of diverticulitis as possible, and the abdomen drained.

Though the prognosis of the former class of case is relatively good, that of the latter is poor. In the present series the one patient in the latter class personally operated upon succumbed.

Abscess with Diverticulitis

The formation of a localized abscess is a not uncommon sequel to diverticulitis. In the majority of cases its formation is accompanied by an attack of acute diverticulitis, with a high temperature, and, not infrequently, diarrhoea. The acute inflammatory condition may subside suddenly when the abscess ruptures into a neighbouring organ, such as the small intestine, rectum, vagina, or bladder. In three cases the abscess ruptured into the peritoneal cavity, and terminated fatally.

The appearance of a collection of pus under the abdominal wall may be accompanied by only the mildest symptoms, but a history of an acute attack of abdominal pain preceding the formation of the swelling is usual. The insidious manner in which the swelling appears may mislead one as to its true nature. In one such case a woman patient had a fluctuating swelling in the left iliac fossa, with no inflammatory signs, and only a small evening rise in temperature. The condition was regarded as tuberculous until investigation of the contents revealed its true nature.

From examination of one specimen, in which there is a small abscess cavity in the centre of the pericolic mass of fibrous tissue, it seems probable that a small abscess around a diverticulum buried in fibrous tissue may be absorbed, without having given any indication of its presence. Indeed, it is probable that such abscesses are frequently formed during an attack of acute diverticulitis.

If it is clear that an abscess associated with diverticulitis is present, it should be opened. Whether or not at the same time colostomy should be performed depends very much on the condition found. Generally speaking, a conservative policy, confined to dealing with the needs of the moment, is wisest. The abscess should be drained, and the inflammatory condition allowed to subside, before deciding upon the line of treatment to be adopted for the underlying diverticulitis. In one case so treated a small faecal fistula developed through the operation incision in the left iliac fossa. This persisted for four years, during

which time the patient was free from symptoms. At the end of this period, however, the fistula suddenly closed, and another peridiverticular abscess developed, which was opened, and a colostomy performed. The latter is still present, and the patient remains well.

Colo-vesical Fistula

Though rarely of itself a cause of death, colo-vesical fistula is one of the most dreaded complications that may follow upon inflammation of diverticula of the colon. Of nineteen cases of colo-vesical fistula due to diverticulitis sixteen were in men and only three in women. The oldest was 69 and the youngest 44; the average age at onset was 54. The characteristic symptom of the condition is the passage of air and faecal material per urethram. In all cases there was a history, varying from thirteen years to one month, of acute pain in the abdomen. The average duration, in fifteen cases, of abdominal symptoms before the onset of the passage of air and faeces was three years and nine months. In most of the cases the onset of urinary symptoms corresponded with an acute attack of abdominal pain accompanied by fever. On occasion urinary symptoms might come on at the height of an attack, and the latter suddenly abate. Thus in one case pus suddenly appeared in the urine during an attack of acute pain, which had lasted for a few days, and on the appearance of the pus the pain rapidly subsided. This was unquestionably due to the rupture of a perisigmoid abscess into the bladder. It may take some little time before a fistulous track from the colon into the bladder is established in consequence of this rupture, so that the passage of air and faeces per urethram may not follow the first appearance of pyuria for some days. In one case the interval between the onset of pyuria and the appearance of faeces in the urine was ten days; the latter followed a dose of castor oil taken to disperse the lump in the left iliac fossa. In another case the interval was two years, and in a third frequency of micturition heralded the onset of pneumaturia.

The quantity of both air and faeces passed once the fistula is established varies considerably. In some cases it appears only at intervals, and there is considerable improvement in between. In two cases faeces were passed, but the patient was never conscious of the passage of air. One of these was a man and one a woman. In a third case, in which this symptom was absent, a colostomy had been performed *before* the fistula developed.

The bladder nearly always becomes inflamed, especially in the early stages, and frequency of micturition both by day and by night is common. One of the most notable features, however, is the apparent immunity to infection by the faeces which the bladder ultimately develops. In these cases, except during the periods when faecal material is passed in considerable quantity, the urine is clear, and the patient relatively free from bladder symptoms. Cystoscopy reveals a healthy bladder mucosa.

Ascending infection from the bladder to the kidneys is of relatively rare occurrence, owing, presumably, to the acquired immunity of the bladder mucous membrane, and this customary freedom from attacks of pyelitis must be borne in mind when deciding upon the treatment to be carried out in each case. In this series only one patient suffered symptoms of pyelitis, and this was limited to one attack. Haematuria occurred in three cases; in one it preceded by a month the passage of air through the urethra. In one or two cases the patient has been troubled at intervals by the passage of large quantities of mucus per urethram. A phosphate stone

subsequently developed in the bladder in two of the cases, in one of which a colostomy had been performed previously to relieve the condition. The onset of symptoms due to the stone was six months after this operation.

The diagnosis of colo-vesical fistula is readily made from the clinical history. X-ray examination is of little assistance, except to exclude the possibility of a malignant fistula. The barium does not readily enter the fistulous track, and it is a matter for comment that in radiographs of some of the cases the bowel does not appear to be grossly affected by diverticulosis. The clinical diagnosis is confirmed by cystoscopy, which will reveal the opening into the bladder. In the majority of cases this is situated near the apex on the postero-superior wall to the left of the midline, though in one case the opening was to the right of the midline. In some cases the opening may be masked because of the oedema of the wall consequent upon infection, and owing to the ease with which faeces pass through the fistula it becomes difficult to obtain a clear medium in the bladder through which an adequate inspection can be made. In such cases a preliminary course of bladder lavage must be instituted, and an attempt must be made to keep the bowels constipated for a few days prior to the cystoscopic examination.

Treatment of Colo-vesical Fistula

The ease of diagnosis is but a poor compensation for the difficult problem which confronts one as to the wisest treatment to adopt for colo-vesical fistula. The histories of the nineteen cases were as follows: colostomy, three cases; radical operation, one; radical operation and colostomy, three; no operative treatment, twelve; colostomy advised but refused, five; and spontaneous healing, one.

Colostomy.—The results obtained in three patients indicate that the case for colostomy is not a strong one. In one patient a fistula formed a year after colostomy had been done for diverticulitis. One should not lose sight of the fact that a colostomy is not the most desirable thing for the patient to live with, and the patient should not be made to pay the price of a permanent colostomy without any guarantee of reward in the alleviation of the bladder symptoms. In five cases in which colostomy was suggested the patients, probably wisely, turned it down, preferring to put up with the inconvenience of a fistula into the bladder than of one on to the abdominal wall. It must be remembered also that the bladder frequently becomes tolerant to the infection, and attacks of pyelitis are rare. The diverticulitis abscess has ruptured, and is draining into the bladder, so that which colostomy aims to do by operation has been achieved naturally. Moreover, there seems to be just as much liability to the development of a secondary abscess with the bowel draining on to the abdominal wall as with the bowel draining into the bladder. In one case urine actually leaked through the lower colostomy wound when the patient passed water. On this evidence colostomy is not a sound line of treatment, and certainly should not be undertaken without the patient being fully alive (a) to the discomfort of a colostomy, and (b) to the absence of any guarantee of cure of the bladder symptoms.

Radical Operation.—In the one case in which the radical operation was performed in one stage the fistula recurred after four months. The one-stage operation is unsuited to the condition, and I do not now give it any place in the treatment; in my view it should be abandoned.

Radical Operation with Colostomy.—Of the three cases in which radical operation was combined with colostomy one patient has been lost sight of. In the successful case

a first attempt at closure failed. The essential difference between this and the second and successful attempt was that in the latter the suture lines in the bladder and colon were separated by interposing the omentum. Though all three patients survived their operations, reference to the literature shows that the operation carries a relatively high mortality. It is difficult of accomplishment, and the subjects are nearly all poor operative risks. Suitable cases in which operation is justifiable are therefore few and far between, and if doubt exists discretion will be the better part. So far the case for operation is not a strong one. Colostomy alone is disappointing, and the operation for closure is only justifiable in a small proportion of cases.

What is the result of non-operative treatment? One patient in this series apparently underwent a spontaneous cure. Ten of the remainder continued in fair health. Not one of them suffered the stormy incidents that figured in some of the operated cases. In nearly all of them the attacks of abdominal pain were less frequent and less severe after the fistula had formed than before. The colon had acquired a natural safety-valve. This, in conjunction with the high degree of bladder tolerance and freedom from pyelitis, already commented upon, allowed the patients to lead a comparatively happy existence. At least five of them preferred the fistula to a colostomy.

A great deal can be done to alleviate the bladder condition by frequent bladder wash-outs, either in a course or at regular weekly intervals. The technique is simple, and the patients have little difficulty in acquiring it themselves. Dilute antiseptics, such as 1 in 20,000 oxycyanide of mercury, should be employed.

Conclusions as to Treatment.—Owing to the extreme difficulty of operation, the lack of guarantee of success, the high mortality, and the poor operative risk, an attempt to close the fistula at operation should only be undertaken when the patient is comparatively young and the fistula of recent occurrence. The operation should be preceded by a preliminary colostomy.

I wish to thank the Council of the Royal College of Surgeons for permission to use material from the Jacksonian Prize Essay, 1932, in preparation of this review, and my medical and surgical colleagues at King's College Hospital for permission to utilize their cases. I also wish to thank Sir John Thomson-Walker for his very kind loan of the clinical notes of some of his cases of fistula into the bladder.

BIBLIOGRAPHY

- Edwards, H. C.: *Lancet*, 1934, i, 221.
 Fifield, L. R.: *Ibid.*, 1927, i, 277.
 McGrath, B. F.: *Surg., Gynecol. and Obstet.*, 1912, xv, 429.
 Mailer, R.: *Glasgow Med. Journ.*, 1928, cx, 193.
 Mayo, W. J.: *Ann. of Surg.*, 1930, xcii, 739.
 Rankin, F. W., and Brown, P. W.: *Surg., Gynecol. and Obstet.*, 1930, l, 836.

The fourth conference of the International Association of Preventive Paediatrics (medical section of the Save the Children International Union) will be held this year at Lyons on September 27th and 28th. The subjects to be discussed and the names of the rapporteurs are: (1) "The Prophylaxis of Malaria in Children," by Professors Cacace (Naples) and Gillot (Algiers), with whom will be associated Dr. Larrouy (a British rapporteur will be named later); (2) "The Prophylaxis of Rickets and Convulsions," by Professors Adam (Dantzig) and Monrad (Copenhagen). Those who desire to be present at the conference, as well as to take part in the discussions following on the reports, are requested to communicate with the secretary of the I.A.P.P., 15, Rue Lévrier, Geneva, Switzerland.