A NOTE ON THE TREATMENT OF ACUTE INFECTIVE ARTHRITIS OF THE KNEE-JOINT

BY

T. B. MOUAT, M.D., M.Ch.Ed., F.R.C.S.
HONORARY SURGEON, THE ROYAL INFIRMARY, SHEFFIELD

When a case of acute infective arthritis of the knee-joint is progressing unfavourably, in spite of repeated aspirations of the joint fluid and immobilization of the limb with effective extension, arthrotomy is usually performed. Arthrotomy on the ordinary lines—by antero-lateral and postero-lateral incisions—frequently fails to arrest the spread of infection, and it then becomes necessary to sacrifice the limb to save the patient's life.

During the performance of amputation through the thigh for this condition many surgeons, and particularly

FIG. 1.

those who had experience of surgery during the war, must have been impressed by the frequency with which an abscess is found to have spread up the thigh, from the subcrureus pouch to the cellular plane in front of the femur, deep to the quadriceps extensor muscles. The resemblance to the spread up the forearm in pyogenic infections of the common flexor sheath is striking. Kanavel described this, and the method of treatment he introduced, by prompt and early drainage of the infected sheath and of the cellular plane deep to the flexor muscles by the lateral incisions he described, which effectively drain the cellular space lying in front of the pronator quadratus and interosseous membrane, has completely altered the prognosis of that serious lesion.

In acute infective arthritis of the knee-joint the deep extension of infection to the subcrureus pouch and thence to the cellular plane in front of the femur is comparable with the line of spread of infection in the forearm, and early treatment on similar lines is advocated: by early drainage of the subcrureus pouch through lateral incisions, the outer of which is continued freely upwards through the vastus externus down to the bone, to drain the cellular space above the subcrureus pouch in front of the lower third of the femur.

The opportunity of making a complete dissection of the limb occurred some three years ago, at a necropsy in which death was due to septicaemia following infection of the knee-joint through a small wound from the point of a pick. The involvement of the joint was not originally suspected, and the acute infective arthritis which super-

vened had not been arrested by the incisions which were made a week later through the capsule of the joint. At the necropsy little pus was found in the joint itself, but a huge abscess, which contained over a pint of stinking pus, was found to extend upwards from the subcrureus pouch, in front of the lower half of the femur, deep to the extensor muscles. I was impressed by the resemblance to the spread of pyogenic infections of the flexor sheath up the forearm, and resolved to try the effect of treatment on similar lines at the first opportunity, which curiously enough arose the same night, in the following case:

A. B., male, collier aged 59, admitted to the Royal Infirmary, December 25th, 1931. He had been struck on the front of the right knee by a sharp corner of a steel plate, ten days earlier. The small lacerated wound on the outer side and above the patella became infected, and three days before admission he developed an acute infection of the knee-joint with rigors, and became seriously ill. On admission, pus spurted from the sinus on movement of the limb and on pressure over the swollen joint. A moderate amount of pus

FIG. 2.

was evacuated when the joint capsule was opened by lateral incisions. The outer incision was prolonged up the thigh through the vastus externus down to the femur. Some ounces of pus escaped from the subcrureus pouch, and a large abscess cavity, which extended up the lower third of the thigh, in front of the femur, was opened by the free upward extension of the incision. The infected subcrureus pouch, and the large abscess which ran from it up the thigh, were drained with strips of dental rubber. The infection gradually subsided, and the patient left the Infirmary at the end of three months with a stiff knee-joint but a useful limb, and resumed light work three months later.

Since then similar treatment has been employed, with success, in the following cases:

C. D., female aged 3½, admitted to the Sheffield Royal Infirmary on July 25th, 1932, with a history of three days' illness. The right knee-joint and lower thigh were swollen, with some redness of the skin, fixation of the knee-joint, and great tenderness over the lower end of the femur. Temperature 103.5°; pulse 135. Radiograms showed no evidence of any focus of infection in the bone. Pus was found in the joint on aspiration (*Streptococcus haemolyticus*). The capsule of the joint was opened by lateral incisions below the patella, and drains inserted down to the capsule, by the resident surgical officer. The temperature and swelling persisted, and on August 11th the subcrureus pouch and an extension abscess up the thigh, in front of the femur, were freely opened by lateral incisions, and drained with strips of dental rubber. The temperature thereafter remained down and the general and local condition slowly improved. The ultimate result was excellent, and the photographs of the case show that the range of movement is practically normal (Figs. 1 and 2).

E. F., male aged 31, on December 5th, 1932, ran a narrow chisel into the left knee-joint just above the patella. He thought it was merely a flesh wound, and applied pink lint and a bandage. The knee became very painful and swollen during the night, and on the following day the local condition was worse and he had several rigors. On admission to the Royal Infirmary on December 7th, the joint was distended with the skin inflamed around the small punctured wound, with streaks of lymphangitis up the thigh. Pus was aspirated from the joint (*Streptococcus haemolyticus*), and ether, introduced through the needle, bubbled out through the punctured wound, which had penetrated the subcrureus pouch. The subcrureus pouch, which was distended with pus, was emptied and drained through lateral incisions, the outer of which was carried upwards down to the femur. The drains were removed on the third day, and the discharge from the incisions and the joint reaction gradually subsided. Immobilization with weight extension was continued for three weeks, and he left hospital at the end of six weeks. He attended the massage department for four months, and when discharged had regained 45 degrees of flexion movement at the joint.

G. H., female aged 55, admitted to the Royal Infirmary, March 6th, 1933, diabetic, developed acute infective arthritis of the left knee-joint following extensive cellulitis of the leg. Haemolytic streptococci and *B. coli* were grown on culture of

the pus aspirated from the joint. The joint infection slowly subsided after transverse drainage of the subcrureus pouch and of the cellular space under the quadriceps through upward extension of the lateral incision. The limb was slung with weight extension, but the ultimate result was complete ankylosis of the joint with a fairly useful limb.

In the treatment of punctured wounds about the knee, when there is doubt as to whether or not the joint cavity has been penetrated, 2 or 3 c.cm. of ether should immediately be injected into the joint. Where the wound has involved the joint, the ether blows out from the external orifice, and immediate excision of the whole wound tract is indicated. The reaction set up by the ether in the synovial membrane appears to stimulate its resistance to infection.

Immobilization with suspension of the limb and weight extension and, if need be, repeated aspirations of the joint exudate are carried out. Where, however, acute infective arthritis of the joint develops or is already established when the patient comes under observation, the further steps described in the above article, on the lines of the Kanavel treatment of infections spreading up the forearm from pyogenic infection of the common flexor sheath, are recommended.

RENAL GLYCOSURIA MISTAKEN FOR DIABETES MELLITUS

BY

R. D. LAWRENCE, M.D., F.R.C.P.

AND

R. A. McCANCE, M.D., Ph.D.

THE DIABETIC CLINIC, KING'S COLLEGE HOSPITAL

It is over thirty years since Klemperer first drew attention to a harmless form of glycosuria, now generally known as renal glycosuria, but our clinical experience shows that it is still a frequent cause of diagnostic error. In this condition glucose appears in the urine at normal blood sugar concentrations because the renal threshold is low (under 0.17 per cent.), and not because hyperglycaemia exceeds a normal threshold, as in diabetes. The latter is particularly liable to be confused with cases of renal glycosuria in which the leak point is very low (under 0.100 per cent.) and the sugar excretion consequently heavy. Other reasons, too, contribute to mistakes in diagnosis, and unpleasant impositions of unnecessary treatment, as the following cases show.

CASE I

Mr. B., aged 47, underwent a complete examination before going abroad on an aeroplane expedition, for which perfect fitness was essential. Sugar was found in the urine, and a blood sugar curve was carried out, with the following result:

Time	Blood Sugar	Urine Sugar
	Per cent.	Per cent.
Fasting: 50 grams glucose	0.087	nil
½ hour after	0.063	0.4
1½ hours after	0.077	nil
2½ hours after	0.071	0.5

The subject was so anxious to go abroad, and the response to glucose was so unusual, that the man was suspected of having taken a dose of insulin before the test was made. When we repeated the curve (see table below) its general features were confirmed, showing a combination of an extremely high carbohydrate tolerance with a very low renal threshold.

Time	Blood Sugar	Urine	
		Sugar	Ketones
	Per cent.	Per cent.	
Fasting: 50 grams glucose	0.095	0.2	0
25 minutes after	0.125	0.8	0
40 minutes after	0.063	1.1	0
1½ hours after	0.111	0.4	0
2 hours after	0.108	0.5	0

CASE II

Mr. O., aged 30. Sugar was found in the urine during a routine examination when the patient had pityriasis rosea. He was at once put on a strict diet of 9 Lines—that was, 45 grams carbohydrate, 67 grams protein, and 135 grams fat—and, since the urine still contained sugar after six weeks, a blood sugar curve was performed, with the following results:

Time	Blood Sugar	Urine Sugar
	Per cent.	
Fasting: 50 grams glucose	0.116	
32 minutes after	0.176	+
64 minutes after	0.188	+
120 minutes after	0.148	

No record of acetone in the urine was made, but the curve was held to be definitely diabetic, as hyperglycaemia persisted after two hours, and diet was again prescribed. When first seen by one of us, after four months of rigid diet, the blood sugar was normal at 0.099 per cent. two hours after breakfast, and slight glycosuria was present, which suggested a low threshold. Full diet was then prescribed, and one month later the following absolutely normal, though irregular, curve was obtained.

Time	Blood Sugar	Urine	
		Sugar	Acetone (Rothera)
	Per cent.	Per cent.	
Fasting: 50 grams glucose	0.116	slight trace	very slight trace
½ hour after	0.087	trace	slight trace
1 hour after	0.137	1.5	0
1½ hours after	0.077	slight trace	0