BRITISH MEDICAL JOURNAL

LONDON: SATURDAY, DECEMBER 15th, 1934

PAIN AND THE MECHANISM OF ITS PRODUCTION*

BY

DAVID WATERSTON, M.A., M.D., F.R.C.S.ED.

HONORARY CONSULTANT, JAMES MACKENZIE INSTITUTE FOR CLINICAL RESEARCH; BUTE PROFESSOR OF ANATOMY,
ST. ANDREWS UNIVERSITY

You might anticipate that my task as an anatomist in opening this discussion would be to describe the structures concerned in pain production, the distribution and the connexions of the nerve apparatus, and to present to you a more or less finished picture of the anatomy of the appropriate parts of the nervous system, leaving you to discuss the bearing of what I might say on the clinical manifestations with which you are all so familiar. That method of approach to the subject would be a stereotyped one, but in the present instance it is inapplicable. Let us try another method, first seeing where we stand and what exactly we want to know about pain; then we can approach the problems from the experimental side, discuss the results of these experiments, and see where they lead and what the next step is to be.

Pain is the most important of complaints, and the commonest and most clamant symptom with which the physician and surgeon have to deal. It drives the patient to his doctor, whose most urgent task is often its relief, while the search for its cause provides some of the most difficult problems he has to face. On the clinical side volumes have been written about it; I need only mention the classical work of John Hilton, and the monumental works of Behan and Pottinger. The current literature dealing with it is enormous; there is hardly a single number of the recent issues of the *Journal of the American Medical Association* in which there is not a reference to, or abstract of a paper dealing with, some aspect of the subject or recording observations upon it. Interest in pain is not confined to the medical world; philosophers and ecclesiastics have pondered over it, and the problem of pain has raised issues of deep significance in philosophy and theology.

Much of the discussion about pain has dealt with its "protective" aspect, but I do not think that any good purpose is served by philosophizing about this aspect of the question. Pain in some instances serves a useful purpose by warning against agencies to be avoided, or by deterring the sufferer from attempting physical exertion which would be harmful to the condition underlying his pain; but this principle is not a universal one. Pain is not always "protective," nor does it invariably arise when harmful agencies are at work, and pathological changes are going on. A burnt child may dread the fire, but the sufferer from renal colic is warned by his pain neither what to avoid nor what to do to ease it; and pathological changes may take place in the body, such

as those which convert a pliable artery into a rigid pipe-stem, without a hint of pain being given.

The definition of pain is also an unprofitable subject of discussion. Like other forms of sensation it can be defined only in terms of itself. Visual phenomena can be described only in terms of sight, and are unappreciable by one who has never seen. We have all experienced pain and know what it is like; we can say of it only that it is painful and immediately recognizable whenever it occurs. It is different from a mere unpleasantness of sensation, which is a "tone" that may accompany other sensations, just as may its opposite, pleasantness or agreeableness.

Our first task must be to discover the mechanism of its production if we are to solve any of the later questions which arise. We cannot, for example, understand the action of anaesthetics or of any means employed to relieve pain unless we know it. Pain is so varied in its forms, and arises apparently from so many different causes, that at first sight it might appear almost hopeless to discover the underlying mechanism and the processes that are at work, and still more difficult and improbable that any common cause operating in all cases will be discovered. The search will be a prolonged and difficult one.

Mechanism of Pain Production

The study of the mechanism of pain production is a problem in which anatomy and physiology can take a part, leaving clinicians to determine the diagnostic value and the prognostic significance of its occurrence. At this age of physiological research, when so many secrets of the action of the nervous system have been discovered, it may seem strange that we still have to discover the mechanism of pain production, but this is indeed the case. We do not know what are its nerves, or, indeed, whether it has its own nerves or not, nor whether the "centre" is in the thalamus, if there is a "centre"; nor do we precisely know the pathways for its impulses within the central nervous system. The reason for this ignorance is doubtless due largely to the impossibility of carrying out such physiological experiments as can be employed for other forms of nerve activity. The existence of pain can be recognized only in a sentient animal, and man himself must undergo experiment.

Another reason why pain is not better understood is found in the failure to recognize in it a definite form of sensation, with its own nerve apparatus. In many people's minds there is present the idea that pain is peculiar and unlike other sensations: that it is merely a manifestation of some sort of increased or disordered activity of any sensory nerves, and that, for example, it may arise from "over-stimulation" of these nerves. There is also the idea that this form of sensation can arise

*The opening paper of the discussion on "Pain," held in the Section of Neurology, Psychological Medicine, and Mental Diseases at the Annual Meeting of the British Medical Association, Bournemouth, 1934. [By inadvertence a contribution to this discussion appeared in the *British Medical Journal* of November 17th before the opening paper had been published.]

in consciousness independently of the afferent impulses which are its ordinary cause; that there is what is termed, as I understand it, "psychogenic pain."

All recent work on sensory physiology is opposed to the former of these ideas. It has been shown that variations in the stimulus applied to afferent nerves alter the number of impulses which are transmitted by them, and increase or diminish the intensity of the sensation, but there is no alteration in its quality. Division of the optic nerve may excite some visual sensation, but it does not cause pain, and the cutaneous nerves of the epidermis, the nerves, as I believe, of tactile sense, can also be divided without the production of pain. With these examples before us it is difficult to imagine that pain is ever due to impulses along nerves of other forms of sensation.

Pain is often, but not invariably, accompanied by other symptoms such as pallor, fainting, and sweating, and these are sometimes ascribed to the pain. They are not, however, invariable concomitants of pain, and hence we must conclude that they are produced in a different way, by other afferent impulses than those which cause pain, travelling by a different path and producing their effects often by exciting the sympathetic system. These concomitants fall into Mackenzie's group of "associated phenomena."

Different Pain Response from Different Tissues

The mechanism of pain production may be studied in two successive steps: the first is to determine the tissues and organs that can give rise to pain, and the second, the effective stimulus in each of them for its production.

It is a well-recognized fact that different parts of the body are entirely different in their pain responses. I remember well a case of enormous prolapsus of the rectum which came under my care during the war, and the difference in the sensory responses which could be elicited from its surface. The prick of a pin near the margin of the anal orifice was painful, but whenever one passed beyond a definite line and stuck the pin into the mucous membrane higher up no pain whatsoever was elicited. This afforded a simple but dramatic illustration of "visceral insensibility," or, in other words, that stimuli which are effective or adequate for pain in some tissues, which we may call the "somatic," produce no sensation from the "viscera." We know that the alimentary canal is so insensitive to such stimuli that the intestine may be cut or pierced or even burned without any pain resulting.

We have not as yet equally precise and well-established information about other viscera, but there is evidence to show that the lungs, the heart, and the brain itself show the same sort of "insensitivity," in that pain is not produced from them by mechanical stimulation such as that described, though it may originate from other causes. This is a field of study in which clinical observation on the sensitivity of exposed organs will in time provide the necessary data. The problem of visceral pain presents special features, and my own view is that its mechanism will not be understood until we know more about "somatic" pain—namely, the mechanism of its production from the tissues and organs of the body wall and limbs.

Experimental Mechanical Stimulation of Different Tissues

The experiments which I have carried out in work on the first of these problems have been published, and I need only summarize the results. They were devised to test whether pain could be produced by direct mechanical stimulation of different tissues. The stimulus used was a sharp needle, the point of which was passed through

the skin and onwards into different tissues, muscle, fibrous tissue, periosteum, and also into the walls of veins and arteries. These experiments showed that among "somatic" tissues there are great differences in the responses elicited by stimulation in this way.

Epidermis has its own nerves, but pain does not arise from them, even when they are divided. The dermis gives an immediate and acute response, as everyone knows, to a wide variety of mechanical stimuli. The subcutaneous tissue is less responsive. For the greater part it does not give a pain response to mechanical stimulation. We all know that the introduction of a hypodermic needle is painless once the skin has been penetrated. The subsequent injection may, however, cause acute pain, and evidently in subcutaneous tissue there is a pain mechanism. Fibrous tissues are responsive to direct mechanical stimulation. Acute pain is caused by the passage of a needle through the fibrous aponeurosis of muscles.

The walls of veins are almost entirely insensitive or give but slight response. This is my experience, but some who have observed the results in a similar experiment have found that considerable pain was produced. One observer told me that in himself he found pain produced almost as severe as that which I have described from arteries. Arteries give a very acute response: the pain is immediate, severe, and sickening, and tends to radiate along the course of the vessel. In my own observation, the wall of my brachial artery was penetrated, and I have received confirmation of my observations from those who, in their clinical work, have had occasion to open an artery.

Ordinary striated tissue gave practically no pain response to mechanical stimulation; the sensation elicited was dull, felt like pressure, and was unpleasant, but not as a rule definitely painful. Periosteum may give a pain response, but often does not, and bone is definitely insensitive to this kind of mechanical stimulation. To strip periosteum from bone, however, is definitely painful.

The value of these observations lies in the demonstration they afford that the somatic tissues do not all give a pain response to mechanical stimulation, while we know also that severe pain can be produced even from those insensitive to mechanical stimulation, by the operation of other causes. The case of muscle is specially instructive. Severe pain can be produced from muscles which contract if they have not a proper blood supply, and Sir T. Lewis has provided the important discovery that the pain is due to the production locally of a chemical stimulus, which he terms the "p substance." There are reasons which lead one to suspect that a chemical agent may be the stimulus for pain in certain other tissues and perhaps in all of them. Pain, for example, is seldom momentary; it persists for a shorter or longer time after the exciting stimulus has ceased to act, and this can be due only to the continued operation of a local cause.

Pain and Hyperalgesia

The condition of hyperalgesia throws some light upon the mechanism of pain production. Hyperalgesia shows itself by the occurrence of a pain response to a mild form of "stimulation," such as usually, and in the normal condition, fails to cause any pain. Illustrations of this are numerous, and among those with which we are familiar are the tender scalp which is present in neuralgia, and the subcutaneous and the muscular hyperalgesia found in visceral abdominal disease and in angina pectoris, when pressure on a fold of skin or on muscle produces an acute pain.

The same sort of hyperalgesia may be produced by local causes. If, for example, the skin be reddened by the application of a mustard leaf, the affected area is hyper-

aesthetic to warmth, or hyperalgesic, for exposure of the surface to even a mild degree of warmth causes severe pain. Lewis has recently shown that if the skin be injured in a variety of ways a local condition which he terms "susceptibility" follows, in which stimuli usually painless cause pain from the affected area. He has ascribed this to the local production of a chemical substance, which raises the excitability of the terminals of the pain nerves. These facts throw a very important light upon the mechanism of the production of pain, showing as they do again that a chemical factor may be the agent.

"Referred" or "Heterotopic" Pain

While pain is often localized to an area at or near the site of its cause, there is also the large group of instances in which pain is, as it is termed, "referred," as in many forms of visceral disease. Examples of this are innumerable, but one may mention particularly the pain of angina referred to the arm, shoulder pain in liver and diaphragmatic disease, and the headaches which arise from defective ocular accommodation. In this group we should, I suppose, include the labour pains of childbirth, which are referred to regions remote from the uterus. The term "referred pain" for these forms of pain is not to my mind a good one, and its use has led to misconceptions. All pain is "referred" by the sufferer to some place; the point of distinction between the two kinds of pain is that in some forms the pain is referred or localized by the sufferer to the place where the disordered process is at work, while in the others it is referred to some distant region. The distinction between the two types is of importance, and for the former I would suggest that the word "homotopic" be used, while for the latter in contrast we might apply the term "heterotopic." An advantage found in these terms is that they provide a contrast, whereas to "referred" pain there is no suitable contrasting term.

The means of the production of "heterotopic" pain is yet uncertain. Mackenzie's provisional theory of its production, his "viscero-sensory reflex," postulated that an "irritable focus" was produced in the spinal cord by afferent impulses from the viscera, and that this focus was in some way responsible for the transformation of the ordinary impulses of touch and pressure in cerebro-spinal nerves of that segment into impulses which gave rise to pain. The facts at least are clear—namely, that the hollow viscera do not give rise to a sensation of pain from ordinary stimulation, while afferent impulses from them induced by a special local condition—spasm of unstriated muscle—do cause severe pain, localized to some portion of the distribution of the cerebro-spinal nerves.

There are many difficulties in accepting Mackenzie's theory in its details, and another theory based upon what we have come to learn about pain production may now be put forward. Areas of heterotopic visceral pain and tenderness are very puzzling; there are present in them severe pain and acute tenderness, but there is usually no other feature to distinguish them from their surroundings. The colour is the same, there is no added warmth, there is no pilomotor excitement, and the activity of the sweat glands is unaltered. Some observers have claimed that they have been able to detect slight alterations and especially a difference in the local response to the application of warmth, and some oedema may be present, but the information is as yet rather vague, and additional observations are necessary to establish it.

In my view the close similarity between an area of pain and tenderness due to local causes and one of visceral pain and tenderness render it extremely probable that the same underlying condition is present in both of them. I have found that an area of fibrous and muscular pain

and tenderness due to local causes may be abolished by the local injection of a small amount of normal saline solution. The mode of its action is uncertain; presumably it helps to get rid of the "p substance" which is present. It is known that areas of heterotopic visceral pain and tenderness can be relieved by the local injection of anaesthetics, and, further, Dr. Forman has found the striking fact that normal saline solution injected into these painful and tender areas can act in the same way, and relieves them.

This procedure can act in no other way than by modifying the local condition, and the observations demonstrate that in "heterotopic areas" the same underlying condition is present—namely, that there has been a local production of a "pain-producing" substance. So far we know of the production of a "p substance" from muscle by local causes. On the evidence provided it seems probable that this substance may be produced in other ways, as by a visceral reflex, and I think that we shall find that in the areas of "heterotopic" visceral pain and tenderness there is formed the same pain-producing substance as that which is formed from local conditions, gives rise to pain, and, in a lesser degree of concentration, causes hyperalgesia.

Summary

Pain, as I understand it, is a definite form of sensation, not to be confused with unpleasantness or with discomfort or mental conflict, served by its own nerves, which are part of the cerebro-spinal nervous system. The various tissues and organs show a wide range of difference in their capability of exciting a pain response; some do so in response to mechanical stimulation, others respond only to a special chemical stimulus.

There is evidence to show that in areas of pain and tenderness from local causes there has been formed locally a "pain-producing" substance, which acts upon the terminals of the pain nerves.

This substance is liberated where there has been skin injury and in muscles under certain conditions. Probably it is liberated in fibrous tissues also.

We know little as yet of the way it is produced nor whether it is liberated from other than local causes. The close resemblance of areas of "homotopic" pain and tenderness to those of "heterotopic" pain and hyperalgesia, and the relief to both conditions afforded by local injection of normal saline solution, point to its being occasionally liberated locally from reflex sympathetic impulses from viscera, and this would afford a satisfactory explanation of Mackenzie's "viscero-sensory reflex."

Citing the various ill effects of a sedentary life, G. Laroche, de Chaisemartin, and A. Grigaut (*Bull. de l'Acad. de Méd.*, October 16th, 1934, p. 291) present a study of the changes in the chemical composition of the blood caused by exercise in twenty-one cases. Tests made at the beginning and end of a six-weeks' course of regulated exercises showed a return to normal of the blood cholesterol, urea, and uric acid; the blood content in these substances is evidence respectively of fat, nitrogenous, and purine and protein metabolism. An increase of weight, due possibly to an increased density of the tissues produced by the exercise, and a stabilization of the blood pressure, due to activation of the general circulation, were also noted. As no dietary or medicinal treatment was given, these effects can only be ascribed to the exercise, which is taken to have a regulating influence on metabolism. It is emphasized that the exercises were regulated, and the authors suggest that sports undertaken without due preparation and medical control may have harmful effects.