Wolff-Parkinson-White–type ventricular preexcitation mimicking left ventricular hypertrophy and an inferoposterior myocardial infarct

D. Luke Glancy, MD

Figure. Electrocardiogram obtained during hospitalization for trauma. See text for explication.

The electrocardiogram (*Figure*) meets at least two commonly used criteria for left ventricular enlargement: the R wave in V_5 or $V_6 > 2.6$ mV, i.e., >26 mm with the usual standardization of 1.0 mV = 10 mm (here RV_5 is 35 mm); and $SV_1 + RV_5$ or $RV_6 > 3.5$ mV, i.e., >35 mm (here $SV_1 + RV_5 = 57$ mm). Broad and deep Q waves in leads II, III, and aVF with broad and tall R waves in leads V_1 and V_2 are consistent with an inferoposterior myocardial infarct of indeterminate age. The patient's age of 19 years, however, extends the limits of normal for $SV_1 + RV_5$ voltage to 60 mm (1) and makes myocardial infarction highly unlikely. Even more importantly, the electrocardiogram is typical of Wolff-Parkinson-White–type ventricular preexcitation, with a short P-R interval of 0.11

seconds, a long QRS duration of 0.12 seconds, and a delta wave visible in every lead except aVR and V_6 . Delta waves are responsible for the Q waves in leads II, III, and aVF and the R waves in leads V_1 and V_2 .

Wolff-Parkinson-White–type ventricular preexcitation is a notorious mimicker and can imitate right or left ventricular enlargement, right or left bundle branch block, and an infarct

From the Sections of Cardiology, Departments of Medicine, Louisiana State University Health Sciences Center and University Hospital, New Orleans, Louisiana.

Corresponding author: D. Luke Glancy, MD, 7300 Lakeshore Drive, #30, New Orleans, Louisiana 70124 (e-mail: dglanc@lsuhsc.edu).

of various segments of the ventricular myocardium, in each instance dependent on the location of the accessory pathway (2). This patient probably has a posteroseptal accessory pathway (3, 4). Although repolarization is nearly normal in this patient, in many of those with Wolff-Parkinson-White-type preexcitation the QRS abnormalities are accompanied by repolarization changes that may be mistaken for ischemia. In addition, false-positive exercise tests have been reported (2). Finally, when the atrioventricular reciprocating tachycardia that often occurs in these patients is antidromic, i.e., antegrade over the accessory pathway and retrograde via the atrioventricular conduction system, the arrhythmia is often misdiagnosed as ventricular tachycardia.

The electrocardiographic abnormality in this patient was an incidental finding. He was admitted to the hospital because of being struck by an automobile, which resulted in a comminuted

fracture of his right femur, a laceration of his right superficial femoral artery, and extensive soft tissue damage. He underwent multiple operations without cardiac difficulty, and at no time was a history of cardiac disease obtained.

1. O'Keefe JH Jr, Hammill SC, Freed MS, Pozwizd SM. *The Complete Guide to ECGs. A Comprehensive Study Guide to Improve ECG Interpretation Skills*, 2nd ed. Royal Oak, MI: Physicians' Press, 2002:495.
2. DePadua F, Pereirinha A, Lopes MG. Conduction defects. In Macfarlane PW, Veitch Lawrie TD, eds. *Comprehensive Electrocardiology. Theory and Practice in Health and Disease*. New York: Pergamon Press, 1989:459–509.
3. Josephson ME. *Clinical Cardiac Electrophysiology: Techniques and Interpretations*, 3rd ed. Philadelphia: Lippincott Williams & Wilkins, 2002:355–356.
4. Takenaka S, Yeh SJ, Wen MS, Yeh KH, Wang CC, Lin FC, Wu D. Characteristics and radiofrequency ablation in posteroseptal and left free-wall subepicardial accessory pathways. *J Electrocardiol* 2005;38(1):69–76.