the discharge was sudden, rapid, effectual, and uninterrupted till the whole was expectorated. The sides of the cavity would thus be brought more readily into apposition, and its obliteration effected. The steady, rapid progress towards cure of this case, after the discharge of the matter, independently of the physical indications, shows, I think, how little true solidification of lung there had been.

The absence of the signs of pneumothorax (except in the case of Connor, where the indications of a circumscribed pneumothorax were well marked) is a point of great interest in all the cases detailed. This circumstance may perhaps excite doubts in the minds of some, whether there were really any communication with the bronchi. But I know not in what other way we are to explain the sudden removal of the pleuritic effusion, and the circumstances attending that removal. In the first case detailed, there cannot be the least doubt that a very large accumulation of fluid obtained sudden exit by the bronchi, so as for a time to place the life of the patient in extreme jeopardy. In the last case, the sudden discharge of a small quantity of fetid, gangrenous matter, succeeded by more copious purulent expectoration, was immediately followed by marked changes in the physical signs, which indicated the rapid disappearance of an accumulation of fluid in the pleura. Whether, in this and the other cases, the absence of all signs of the presence of air in the pleural cavity, is to be ascribed to the existence of extensive adhesions, or to any peculiarities of the communication with the bronchi, I am unable to say.

Such are the more important features of the thoracic inflammation, to which I venture to call the attention of the profession; and they are sufficient, I think, to show that it differs, in many most material respects, from the ordinary form of either pneumonia or pleuritis. Upon what does this difference depend? Not, I think, merely on peculiarities of individual constitutions. I have seen the disease in various constitutions, and in persons moving in very different social spheres. It has no necessary connexion with any form of continued fever. Is it connected with erysipelas? I am disposed to think that it is: not that it is generally associated with the external manifestation of erysipelas. The pneumonia that is associated with fever, erysipelas, and cachectic states of system generally, it is well known, is usually of the asthenic type; but I am disposed to regard the pleuro-pneumonia now under discussion as itself a form of erysipelatous disease, and for these, among other reasons: 1st. In its pathological features, it closely resembles erysipelatous inflammation; 2nd. It has prevailed contemporaneously with external erysipelas; 3rd. An erysipelatous form of cynanche has been very common, and, in many cases, has been an attendant on the thoracic inflammation; sore throat having been frequently complained of in the beginning, and, as this has subsided, bronchial symptoms have occurred, to which the pneumonia and pleuritis have succeeded. I feel, however, bound to admit that it differs in some material respects from the ordinary forms under which erysipelatous inflammation of internal organs has hitherto been known to us.

In respect of the treatment of this disease, it is manifest that it must differ materially from that of the ordinary forms either of pneumonia or pleuritis. General bleeding is rarely if ever admissible: I have never employed it. Local depletion, to a limited extent, especially in the early stages, is sometimes decidedly useful, especially from the relief it affords to the pain and the congested state of the parts implicated. Counter-irritation may be more early employed than is desirable in the more sthenic forms. Blisters, indeed, I have found of the most marked benefit, and have been in the habit of repeating them frequently. Calomel and antimony, if used at all, must be very cautiously and sparingly employed. Dover's powder, in doses of five grains, with a grain of calomel, three or four times a day, has appeared to me very useful. But I have not given the calomel with the view of affecting the system, so as to induce any of the symptoms of mercurialisation. Ammonia has proved the best expectorant; and this I have given usually with the acetate of ammonia, sometimes with the

camphor mixture, and at other times with serpentary or senega. In almost all cases, I have allowed the patients beef-tea, and very frequently found a certain portion of wine to be necessary. But, under any plan of treatment, the affection is attended by great danger, and, in its worst forms, very fatal. So rapid has been the progress in many cases, so great the prostration, even with the most cautious treatment, that I have, in more than one case been led to doubt whether the patient would not have had a better prospect by entire abstinence from all antiphlogistic treatment, so called, and the use of stimuli alone *ab initio*. Ample and efficient blisters, with wine, beef-tea, and ammonia, are, I believe, the remedies on which most reliance should be placed in the most aggravated and most characteristic examples of the disease. But such remedies must be used cautiously. Many cases will do well with little or no treatment, even though, as ascertained by the physical signs, there be a considerable amount of inflammation; a simple diet, not too antiphlogistic, rest, and an open state of the bowels, with some counter-irritation, being all that are requisite; anything beyond this, like the homœopathic globules, serving only to amuse the patient, and gain for physic a degree of credit which is not its due.

Since writing the preceding observations, the substance of which was read before the Medical Society of London on the 16th Oct., I have been not a little gratified by finding how much they coincide with those made by my honoured preceptor, Dr. Alison, in the November number of the *Edinburgh Monthly Journal*. I cannot but feel additional confidence in the accuracy of my own observations, when thus confirmed, and I have the less hesitation in pressing on the attention of the profession the views here set forth. But, how long the present type of pneumonia may continue to prevail cannot be predicted, and we must not therefore repeat the errors on which we have ventured to comment, but recall the words of our own Hippocrates. "*Hoc saltem pro comperto habeo ex multiplici accuratissimarum observationum fide, prædictas morborum species, præsertim febres continuas, ita toto celo differre, ut, qua methodo currente anno ægrotos liberaveris, eadem ipsa anno jam vertente forsitan e medio tolles; quodque, ubi semel in genuinam medendi rationem, quam hæc vel illa febris species sibi vendicat, auspicio incidierim, ad eundem scopum colligans (favente, ut fit, Optimo Numine), metam quasi semper attingam, respectu ad temperamentum, ætatem, et reliqua ejusmodi usquequaque habito; donec, extincta illa specie novoque gliscente malo, anceps rursus hæreo qua mihi via insistendum ut ægris subveniam; ac proinde, nisi ingenti adhibita cautela intentisque omnibus animi nervis, vix ac ne vix quidem possum efficere ne unus aut alter eorum qui se primi meæ cure commiserint, vita periclitetur; donec, investigato jugiter tandemque perspecto morbi genio, ad eundem perdomandum recto pede et intrepidus denuo procedam.*" (Sydenham, *Observat. Medic.*, sect. i, cap. ii, p. 29. Sydenham Society's Edition.)

15 Finsbury Square, 13th Dec. 1852.

MECHANICAL SUPPORT IN PROLAPSUS ANI.

By JAMES DEANE, Esq., M.R.C.S.Eng.

It is singular that we hear so little of mechanical support in cases of prolapsus ani. Authors will puzzle us with minute and unpractical distinctions about the first element in different cases of hæmorrhoids, and will then forget that, whether it be their cause or not, the prolapsus, so generally accompanying them, will always aggravate them, and even form fresh piles, however carefully an excision may have been effected. Their treatment, also, has been well discussed; but have we been as attentive to their prevention? Have any means been devised for supporting the perinæum during the expulsion of the fæces? What else can remedy

the relaxed state of the levatores ani? and, if constantly supported during the evacuation of the rectum, what more reasonable than to suppose that the hæmorrhoids will yield to some mild treatment? or, if requiring removal, that their re-formation will be prevented?

Such was the train of reasoning that occurred to me in the after-treatment of the following case.

CASE. A. B., aged 25, had suffered from prolapsus for sixteen years. During that time, the bowel had descended invariably with each act of defæcation. On examination, I found external piles, large and numerous, and a loose fold of integument completely encircling the anus, and projecting about three-eighths of an inch. I found also internal piles, two in number, about half the size of a filbert, and a general abnormal rugosity of the mucous membrane, the nidus of future hæmorrhoids in the present state of parts. She suffered great pain at stool, and the faeces were generally more or less marked with blood. Her general health was very delicate: she had a pale, exsanguined appearance. I removed the external piles by incision, and the loose fold of (skin?) projecting from and encircling the anus. This was, no doubt, originally mucous membrane, which had been protruded, and had taken the form of skin from atmospheric exposure. The wound soon healed, but the prolapsus was as bad as ever, and, if not remedied (the parts being protruded and strangulated for a time daily), would soon have brought on fresh piles, and my patient would soon have receded into her former state.

What was to be done? Here the records of surgery give us no information. I adopted the following plan, with the utmost possible success. My patient has now used it for upwards of two years, and has not during that time had the bowel down, except when accidentally requiring an evacuation when from home. This occasional exception, so certain when the board was not used, distinctly proves that the non-prolapsus, at other times, was due to the contrivance, as truly as the non-descent of the bowel is due to a truss. It is probable that the analogy may go further, and that the *absolute constancy* of this support during defæcation will cure prolapsus, just as a truss cures hernia, when used for years without the permission of one descent of the bowel. I may also add, that, since the use of the board, I have heard no more in this case of piles, either internal or external.

The plan may be described thus:—Cover a chamber utensil with a strong flat board, taking a circular form, and projecting one inch; let into the edge of the wood three or four pieces of wood, to keep it fixed on the chamber-pot; make an opening on the surface of the board, a little behind its centre, just large enough for the anus to act; and surround this opening with an elevated ridge, to support the entire perinæum and sides of the anus during evacuation. The anal foramen which I have used, has been oval, its long diameter being antero-posterior, and four inches in length, and its transverse diameter being two inches and a half. It has been abrupt, not bevelled; the sharpness of the edge being removed. The ridge has been half an inch high, and has extended all round the anal foramen, and has been less than half an inch from it. The great object is to keep it within the ischia. Some of my patients have felt more secure and more comfortable *with*, and some *without* this ridge. For females, I have had a large anterior urinary fissure, half an inch wide towards the middle of the board, and opening into the foramen, and gradually enlarging forwards to the size of three inches in front. With males, I have recommended another utensil in front.

As "there is nothing new under the sun", some such apparatus may possibly have been already used without my knowledge; and I have no doubt that, if the attention of the profession be directed to the subject of this paper, some better plan may be discovered than that which I have suggested.

Chatteris, Cambridgeshire, Dec. 10th, 1852.

AMPUTATION OF THE ARM IMMEDIATELY BELOW THE SHOULDER JOINT, FOLLOWED BY RECOVERY, IN A CHILD AGED FIFTEEN MONTHS.

By ROBERT EDWARDS JONES, M.R.C.S.

On the sixth of September, in consequence of the absence of Mr. Mason, of Sudbury, I was hastily summoned to visit Eliza Harrington, aged 15 months, the child of poor parents, residing at Wickham St. Paul's, Essex, who had been knocked down by a waggon, the wheel of which had passed over her left arm. On examination, I found a fracture of the humerus at its lower third, the bone protruding denuded of its integuments, and the soft parts lacerated nearly to the shoulder joint: the little patient was much exhausted from loss of blood. I administered small quantities of brandy and water, which had the effect of producing reaction; and feeling there was no time to be lost, I at once amputated just below the shoulder-joint, being just able to save sufficient integument to form a good stump. I was ably assisted by Mr. Mason, jun., and very little blood was lost during the operation, and but one artery required a ligature. The child had severe convulsions the following day, which were relieved by small doses of tinctura opii; after which it progressed most favourably, and at the end of a month had perfectly recovered.

I am induced to send this case for publication, because I believe that it is rare for a child of such a tender age to recover from so severe a proceeding.

Long Melford, Suffolk, Nov. 1852.

FATAL DISEASES OF THE STOMACH.

By JOHN CHARLES HALL, M.D., Fellow of the Royal College of Physicians of Edinburgh, etc.

THE most important truths may be derived from a faithful record of those cases in which, notwithstanding all our efforts, the disease we are called upon to treat proves fatal. However pleasing and flattering to our vanity it may be, to publish those cases only in which we have been successful,—fatal cases, at least, as much as those which have had a favourable termination, convey important lessons.

CASE 1. On November 1st, 1852, I was requested to see, in consultation with my friend, Mr. Henry Julian Hunter, surgeon, Mr. F——, of Sheffield, aged 46, a brass-turner. From Mr. Hunter I learnt that he became his patient on the 20th of May, 1852, with symptoms of dyspepsia, under which he had then laboured for some time. He prescribed mild laxatives, with infusion of calumba, and prussic acid, and, on the 25th of May, the nitrate of bismuth and rhubarb. Under this plan of treatment, and a mild diet, the patient became much better. Mr. Hunter did not see him again until the 27th of August, when he called upon him, having, meantime, been under the care of Dr. ———, who had ordered him to be rubbed with the linimentum hydrargyri, etc. He had also been to Dr. M——, a physician, residing some distance from Sheffield, who prescribed calomel and the compound powder of antimony, and a simple saline mixture. He had now the same symptoms as when last attended by Mr. Hunter, and the same plan of treatment was again adopted.

Sept. 19th. He was much worse; had obstinate constipation; almost constant vomiting; and severe pain in the region of the stomach. Small doses of castor oil, and clysters of warm water, were given. An ointment of tartarised antimony was applied over the region of the stomach, which gave some little relief. He continued much the same until Oct. 7th, when all the symptoms were aggravated; the pain and sickness were constant, and the food rejected almost as soon as taken. Sometimes, however, it remained some hours, and was then vomited. The bowels were kept open with clysters; leeches, and a plaster of belladonna, were applied to the epigastrium.