ORIGINAL COMMUNICATIONS.

ON INTESTINAL OBSTRUCTIONS.

By BENJAMIN WEBSTER BROWN, M.R.C.S.Eng.

(Read before a Meeting of the Midland Branch of the Provincial Medical and Surgical Association, at Nottingham, December 2nd, 1852.)*

It is my intention to attempt to point out, first, the causes; secondly, the diagnostic signs; thirdly, the treatment of obstructions of the intestines; and lastly, to illustrate my paper by the history of some of the most interesting cases which have fallen under my own observation.

CAUSES. We are aware that the intestines are endowed with a peculiar property, viz., the peristaltic motion; and it would appear they are assisted in this by a fatty network substance—the omentum,—formed by several duplicatures of the peritoneum, and suspended in front of the alimentary canal.

It seems requisite for the purposes of life, that this peristaltic action should be preserved uninterrupted, to propel the fecal contents onwards; and hence we might at once be led to conclude that some serious derangement to the animal economy would arise from its suspension, whether the cause be mechanical or otherwise.

Experience teaches us there are several causes which have a tendency to retard or suspend the peristaltic action, and entirely obstruct the intestines; amongst these we may class obstinate constipation; ileus; intussusceptio; stricture of the canal; tumours; hernia, etc.

Before we proceed to obstruction of a more formidable character, we will take its most common cause—constipation of a more or less aggravated kind. This may exist *per se*, and yield to simple treatment; sometimes it is constitutional, and no particular ill consequences ensue primarily; some people, however, suffer considerably from an idiopathic condition, and if remedial means have for a length of time been neglected, a morbid state is very frequently set up in the intestinal tube, and the foundation of more serious mischief is laid, as producing an enlarged colon and rectum; this happens, however, in females more than in males.

But constipation may be symptomatic, as in enteritis, peritonitis, intussusceptio, stricture of the rectum, or of any other part of the intestines, hernia, etc. It is sometimes the direct cause of some of these maladies, and becomes the harbinger of a more alarming crisis, if not relieved.

Persons who addict themselves to the taking of large quantities of opium, or laudanum, most generally have a constipated state of the bowels, which under peculiar circumstances, or certain stages of disease, may become obstinate, and finally lead to real obstruction. Obstruction in the intestines will frequently be produced by taking into the stomach hard and indigestible substances; as improper food; the swallowing fruit-stones, or other like bodies, which are apt to lodge in the fecal passages, and produce irritation. I have seen a kidney-bean, on a *post mortem* examination, incarcerated in the appendix cæci; in another instance a plum-stone; each of these was the cause of permanent obstruction, and death. Permanent obstruction, accompanied with very acute suffering, occurred in the case of a young lady who had swallowed a large quantity of cherry-stones.

Constipation generally accompanies scirrhus of the pylorus, as well as stricture of the œsophagus, from the small quantity of nourishment which is received into the intestines.

Impacted scybala are a frequent cause of obstruction to the passage of the upper portion in the canal. These will most generally be situated in the rectum; but this unnatural condition of the alvine contents will also take place in other parts of the large intestines, as in the colon, which

has become enlarged sometimes by long neglected constipation, and will lead to alarming results if not dislodged by remedial measures. This is more likely to occur here, from the sacculated structure of this part of the canal.

Spasmodic colic, or constriction of some part of the intestinal canal, is an occasion of obstruction. This may be brought on from constipation, the taking of improper nourishment, or the immoderate eating of acid substances or unripe fruit; drinks too, which are of an acrid character will be a source of it. Instances, indeed, occur in the practice of every medical man, in which, on one or more of these causes, invincible obstruction and inflammation supervene.

Obstruction very frequently arises from intussusceptio, or the inversion of some part of the intestines; and this may be *progressive* or downwards, or *retrograde* and upwards. This preternatural position of parts is liable to occur in any portion of the canal. From the construction of the parts, the most probable place for such a derangement is at the point where the ileum terminates in the cæcum. Instances, however, are on record in which portions of the colon have descended into itself, and even through the rectum, and *per anum*, so as to represent the appearance of a procidentia.

Intussusceptio most commonly takes place in the small intestines, in which there may be more than one volvulus. Children who have died from teething and convulsions, have been found on *post mortem* examination, with several volvuli in this portion of the tube, which in some instances, apparently, during life, produced no great inconvenience, and consequently were scarcely suspected. When this has occurred, it has been imagined by M. Louis, that owing to the peristaltic action of the bowels, the parts may again have resumed their normal condition.

Whilst at St. Bartholomew's Hospital, I remember a female subject, apparently about twelve years of age, being brought into the theatre for demonstration before the class. Mr. Abernethy had commenced his lecture on the abdominal viscera, and the small intestines nearly throughout their whole length were found in a state of intussusception, yet none of the usual appearances of inflammation were present. I saw Mr. A. draw out or unfold at least a dozen volvuli; and from the absence of inflammation and adhesions, the intussuscepted parts were drawn out without the least difficulty. What the symptoms or history of the case were during life, it is impossible to say; but Mr. A. at the time delivered some useful remarks.

It has sometimes been questioned how intussusception of the intestines can occur? Mr. Hunter affirmed that the abdominal muscles possessed no influence in forming a volvulus, although a prolapsus ani might be produced by their united agency, as in hard straining, etc. I think, from my own experience, and from the numerous instances on record, that it is probably produced by spasm, the result of cold, or as we have before mentioned, of improper aliments and drinks, or the sudden cessation of diarrhœa of a peculiar character, as well as from sudden emotions of the mind. I have had cases under my own care wherein intussusception has been immediately preceded by a severe attack of diarrhœa, accompanied with considerable twistings across the umbilical region. Others have come under my notice which have been the result of some sudden surprise and alarm, thus giving a shock to the nervous system.

Falling, or jumping from a great height has brought on obstruction in the intestines of an invincible kind; a portion of the ileum has been found to have passed through an opening in the diaphragm into the pleural cavity, and strangulation has ensued. From some such cause, or external violence, or even spasm, a rent has been produced in the mesentery, and a portion of intestine has passed through and become strangulated, and thus obstruction has taken place. A part of the colon has been found in a state of strangulation, having passed through a small opening in the diaphragm several inches into the cavity above.

From a morbid condition of the parts, adhesive bands of organized lymph will frequently be found in the abdomen,

* From the pressure of matter, we have been reluctantly compelled to omit some of the introductory paragraphs of Mr. Brown's paper.

which lead to a permanent obstruction. The intestines have on examination been found glued together in several parts by bands of adhesive lymph; consequently, there has frequently been a contracted condition of the larger portion of the tube. Amongst the French archives, we find an account of a youth of fifteen, who, having neglected a call of nature for several hours, in consequence of not being able to find a place of convenience, besides being exposed to a cold wind, became ill; he felt a degree of tenesmus, but no effort could procure an evacuation. Warm fomentations were applied over the abdomen, which had become very painful and tumid; enemata were administered, and other means were made use of; yet, although some hardened faeces were brought away, there was no abatement in the severity of the symptoms, which presented every feature of internal strangulation. Death ensued in about ten days from the commencement of the attack. A *post mortem* examination discovered a high degree of peritonitis; the intestines were glued together by the effused lymph, which had formed adhesive bands. The tumid state of the upper portion of the abdomen was owing to a distended condition of the bowels, but this did not exist below the ileum. At this point there was a loop of intestine, which was crossed with numerous bands, and where, consequently, constriction had occurred. The larger portion of the tube was so constricted as to be very unnatural in appearance, and was more like cords; there was also a rent in the mesentery beyond the caecum, and through this opening a large portion of intestine had passed, and became strangulated.

The next morbid affection to which the alimentary tube is liable, frequently preventing the proper passage of the faeces, and ultimately in most instances terminating in a permanent obstruction, is a scirrhus-contracted state of the rectum. This generally occurs in people, and more frequently in women, who are somewhat advanced in life; it may be confined to a particular part of the bowel, or nearly its whole length may become affected. Sometimes there is more than one constricted portion of the gut, and the diseased part is most frequently found in a gristly condition, accompanied with ulcerated portions of the bowel beyond the constriction. I have found the caecum near its caput in a scirrhus state, which ultimately produced invincible obstruction, and death under very acute suffering.

This morbid condition of the parts appears to be the result of a slow, chronic kind of inflammation which has been set up, producing, as already observed, a thickening and disorganized state of the bowel, which progresses more or less rapidly: permanent obstruction, which no remedial art can cure, takes place ultimately. A case of this kind will sometimes carry the patient off in a few hours, after intense suffering: or it may be protracted to a much longer period, unaccompanied, apparently, by any severe pain; and the patient will die of starvation.

Occasionally, cases occur where there is obstruction arising from congenital contraction of the intestines. The infant shortly after birth will suffer from vomiting and constipation, which are unrelieved by any of the means usually adopted. A *post mortem* examination reveals the cause of all such suffering; there is an impervious condition of, perhaps, the greater portion of the rectum, or of some other part of the alvine canal.

Then, again, we have hernia under its several and direful forms, cases of which present themselves to the notice of almost every practitioner, whether of a simple kind, or those of a more formidable nature. On these, however, I need not here further enlarge.

Thus I have endeavoured to point out the principal causes which give rise to obstruction in the alimentary tube. There may be a few others, as malignant or other tumours, etc., but it is not necessary that I should dwell longer on these.

DIAGNOSIS. We have already remarked that constipation is frequently constitutional as well as symptomatic. Without waiting to describe the former condition further, it will be well for us to endeavour to point out the *diagnostic* signs which appear to present themselves in the

several causes of intestinal obstruction of a more serious character; which, we have said, may be *mechanical*, or otherwise. And here considerable acumen and experience are requisite. In the former case especially, every practitioner knows the difficulty with which he has to contend, before he can in any way satisfy himself with the precise nature of the affection, or seat of the obstruction; and, after all, he is more generally left in extreme doubt, so true is the remark of Hippocrates, "*judicium difficile*." The general history of a case of this kind, the character of the symptoms, the suddenness of the attack or otherwise, connected with other circumstances, will, in some instances, enable him to draw a just conclusion; but, in others, everything leaves him in uncertainty. The perusal of cases, or learning their history, symptoms, and the *post mortem* appearances, and comparing these with his own experience, relieves him but little of his perplexity; so that it appears there is no class of symptoms so characteristic or constant, as will enable him at all times to come to a correct diagnosis.

A scirrhus, or diseased state of the pylorus is usually distinguished by a dense, heavy, yet circumscribed pain in the right hypochondrium. Pain is experienced after each meal, accompanied with a rejection of the food which is taken; constipation, frequently of an obstinate character, invariably accompanies this form of disease. In some cases, where the parietes of the abdomen are thin, the indurated part may be felt.

In a case of intussusceptio, there is mostly considerable difficulty experienced in distinguishing its real character, so varying is it in its attacks. It is, however, generally preceded by an uneasy sensation across the umbilical region, with a dragging down sort of sensation in the intestines, and the pain, which occurs at intervals, in some instances is severe. Sometimes the affection comes on by degrees, and is accompanied with invincible constipation; or it immediately succeeds a severe attack of spasmodic diarrhoea. A gurgling noise is sometimes heard in the abdominal cavity, as if nature made an effort by the peristaltic action to remove the obstruction; and this is marked with a severe increase of pain. Frequently the patient will express himself as having his "bowels fast and tied together." There is, usually, considerable nausea and vomiting of bilious matter; this, however, at first, is not always present; but, as the case becomes aggravated, it is ever an attendant symptom, and it is not unusual to have all the characteristic features of a case of strangulated hernia present; but the absence of any tumour or hardness in the abdominal openings will at once relieve our suspicions on this point, especially when other circumstances are considered. At the onset, the state of the pulse is frequently little affected; sometimes, however, it is full and hard, with pyrexia, and the tongue is generally much coated. Tenesmus prevails without the power of defaecation.

As the symptoms continue, and no evacuations follow, the abdomen becomes considerably enlarged and resonant. The pain, which is fixed and local generally, is little increased on pressure, unless inflammatory action have commenced; during the severe paroxysms, however, if the case last some days, the patient will exhibit an occasional heavy and suppressed moan; frequently the agony is so great, that he will cry out aloud, no relief being obtained by the various postures he may adopt. The respiration, generally, is little affected. As the obstruction remains invincible, the abdomen becomes tympanitic; and, with the vomiting faecal matter, hiccup occurs. The respiration becomes hurried and short; the pulse sinks, and cold sweats supervene, as the sure preludes of death.

Intussusceptio will be distinguished from peritonitis, enteritis, or hepatitis, by the absence of the usual symptoms of pyrexia in the one, and by the pain and tenderness in the other being much increased on pressure. In peritonitis, the patient can scarcely bear the bedclothes to come in contact with the abdomen; and the position chosen is to lie with the body bent considerably forward, the legs being drawn upwards, in order to relax as much as possible the abdo-

minal muscles. Besides, in these inflammatory affections the respiration is quick and short, and carried on as much as possible by the ribs, so as to ease the diaphragm; considerable pyrexia is always attendant; the pulse beats from 120 to 130 or more in a minute; it is generally feeble, but frequently hard and wiry. In peritonitis or enteritis there is grating of the teeth, as a sign of intense pain of which there is little intermission: whereas, in intussusception, there are intervals of fifteen or twenty minutes between the paroxysms of pain, until the symptoms become more aggravated as the case advances. The skin in pyrexia is usually hot and dry; in intussusception it is cold and moist. These, I believe, are the usual characteristics attendant upon obstruction in the intestines, arising from intussusception: but, of course, as we have remarked, they are more or less variable.

A scirrhus or constricted state of the rectum is usually preceded and accompanied by constipation, and very generally terminates in permanent obstruction of the intestines. It may, sometimes, be known by the pain which is experienced within the anus, or along the sacral and lumbar regions, increased at each evacuation, which, as the disease makes progress, seldom occurs more than once in five or six days. If the constricted portion have existed some time, we may generally form a tolerably correct diagnosis of the case, if we bear in mind other circumstances. The stools are frequently streaked with blood and mucus; and are unnatural in their shape, being more or less flattened.

As the disease advances, tenesmus is frequent, and each scanty stool now is not only streaked with blood and mucus, but discharges of matter likewise take place, when ulcerations have commenced. As there is generally no pyrexia, the pulse is little affected; the tongue is much furred but moist; there is occasionally a hectic flush on the cheeks, the countenance being little disturbed; the patient complains only of occasional pain in the abdomen, which he describes as being of a heavy and forcing character; nausea and vomiting generally, but not always, accompany these cases. The abdomen becomes enlarged and resonant, but without any preternatural heat of the skin. Pressure made by the hand does not increase the pain; enemata are mostly returned on each endeavour to throw them up the bowel. The patient at first will not appear to suffer much in health; but as no evacuations follow the means which have been adopted, an examination *per anum* becomes necessary, and if we judge from the symptoms collectively, little doubt will remain on our minds as to the nature and seat of the occlusion. If the constricted part be situated beyond the reach of the finger, the cautious introduction of moderately sized bougies will aid in forming a right conclusion. Of course, as the constriction increases, and an invincible obstruction to the passage of the fecal contents from above supervenes, the features of the case assume a far different form, and the patient ultimately dies from exhaustion.

I have stated above that organised adhesive bands will frequently occur near the caput cæci and colon. These may have existed for several years without producing much inconvenience; but, continuing to increase in substance by the addition of fresh morbid matter, they induce a constricted state of the part, which has resulted in a permanent obstruction to the passage of the feces; and here all the symptoms of internal strangulation, or spasmodic ileus, may present themselves. The attack will sometimes come on almost suddenly; the pain across the umbilicus, and in the lower part of the abdomen, will appear to be most excruciating; the patient will writhe under it, and put himself in almost every attitude, in the hope of finding some mitigation of his sufferings, and die in a few hours.

Impacted scybala, if seated in the transverse arch of the colon, may be frequently detected through the walls of the abdomen, by pressure with the hand; and, as the symptoms of obstruction become more urgent, there will be considerable enlargement above, in consequence of flatus having collected in the small intestines. When scybala are seated in the rectum, and cause obstruction, they are usually discovered by an examination *per anum*.

Cancerous and other tumours of a malignant character will sometimes form amongst the intestines, and give rise to obstruction. These, if situated in the rectum, are readily discovered by digital examination; and, if higher up, as in the cæcum, or colon, they may occasionally be felt through the abdominal parietes. However, to form our diagnosis aright, we must be guided by the symptoms which have been present throughout, as well as by the history of the case.

When the obstruction is dependent upon incarcerated or strangulated hernia, the symptoms and appearances are so well known to us all, that it is almost impossible we should be at a loss to arrive at a correct diagnosis; so that it would be a waste of time to attempt to describe these. I will therefore pass on to consider the third part of my paper, viz., the treatment demanded in each case.

TREATMENT. It is well known to every practitioner, that there are no symptoms so perplexing as those which result from a constipated condition of the intestines. It would be impossible, however, in the present essay, to do more than give a general outline of the chief causes which lead to this condition, bringing on disease, and so frequently terminating in permanent obstruction. These I have endeavoured to point out as fully as a paper like the present will allow, as well as to describe the symptoms whereby we may arrive at a just diagnosis; but, as those causes and symptoms are constantly varying, much judgment is required in the treatment; for every case of this kind is itself a subject for special study.

Without dwelling on those cases of simple constipation, which almost daily come before us, I will proceed to the treatment of those which are of a more obstinate and too commonly invincible character.

The symptoms may indicate that the passage of the alvine contents is obstructed by a distended bladder, an uterus enlarged from pregnancy, or by other causes which produce pressure upon the rectum. These cases will frequently yield to ordinary treatment; but, if the obstruction be referable to other causes, such as spasmodic constriction, the lodgment of hard and indigestible substances, the collection of hardened feces in the colon or rectum, intussusception, or the strangulation of some portion of the intestinal tube, stricture of the rectum, or to other mechanical causes of occlusion, the result of our treatment most frequently becomes exceedingly doubtful, and on many occasions it is unavailable. This is shown to be the case, by the mortality which occurs from this class of diseases.

If the obstruction be dependent on scybala which block up the rectum, they should at once be carefully removed *digitis et aliis instrumentis*, and the bowel washed out with an injection of warm water. When the colon becomes enlarged from long and habitual costiveness, and filled with a collection of indurated feces, etc., which prevent the passage of the alvine contents from above, enemata of turpentine, or olive oil, and warm water, or linseed infusion, should be repeatedly injected through a tube carried high up the bowel. The introduction of O'Beirne's tube as far as into the colon, has been proposed, and warm water has been thrown with some force; but, in doing this, some caution is necessary, lest the bowel should be ruptured. Calomel and colocynth, and castor oil in small and repeated doses, should be given. Should the case prove obstinate, and there are severe tormina, attended with nausea and vomiting of bilious matter, hot fomentations should be applied over the abdominal region. An effervescent saline draught may be given to allay the sickness. The administration of an emetic, after the rectum has been emptied, has, in several instances, removed the obstruction, after other means had been tried in vain. If, however, inflammation threaten, the application of leeches to the abdomen, or the use of the lancet, must be resorted to. Very often a case of this kind will continue obstinate for many days, unaccompanied with inflammation.

When the obstruction arises from spasmodic colic, or intussusception in some part of the intestinal tube, our object should be the prevention of inflammation. If symptoms of an inflammatory nature show themselves, and the

pulse be full and hard, accompanied with tenderness of the abdomen, venesection should be adopted, as well as the application of leeches to the part, together with hot fomentations. Where practicable, it is advisable to place the patient in a warm bath of 95° to 100°; emollient clysters are to be administered; and five grains of chloride of mercury, combined with ten grains of the compound extract of colocynth, should be given, and followed by repeated doses of castor oil. If there are violent tormina and spasm, it will be very proper to give a full dose of opium or laudanum; the former in a pill of two grains and a half, combined with three of calomel; the latter in a dose of sixty drops, in any agreeable liquor. This plan will frequently act as a charm. In some cases of obstruction, the persisted use of strong purgatives is absolutely mischievous. If intussusceptio be suspected, half a drachm of tobacco should be infused in half a pint of warm water, and the liquor thrown up as an enema. Emetics have been recommended, and quicksilver has been given in large quantities, with the belief that by its mechanical property the volvulus would be destroyed, and the parts restored to their normal condition; but some judgment is necessary in its administration, lest the mischief be increased instead of relieved. If, however, active purgatives have been administered for several hours, and no evacuations follow, they should be discontinued, and only those of the mildest nature given.

In most cases of obstruction in the intestines, a similar plan of treatment appears to be the one best calculated to relieve the mischief in a *medicinal* point of view. When the obstruction does not yield to this, chloroform has been recommended; and when symptoms of a severer character supervene, which threaten to extinguish life, it becomes a question how far relief can be obtained by resorting to surgical aid. We are aware, however, there are very few cases of this character which demand operative interference, and those are indeed exceedingly difficult to distinguish.

After a judicious *medical* treatment has been persevered in for several days, in certain forms of obstruction, without the desired relief being obtained, and symptoms supervene, accompanied with stercoraceous vomiting, etc., are we justified in having recourse to an operation, and exploring the cavity of the peritoneum? We are convinced that the patient will die, if no other means are resorted to than those we have already adopted: shall we cut through the walls of the abdomen, search for the cause of obstruction, and endeavour to remove it? This question is a very important one, and requires much serious consideration. The fact, however, that the abdominal parietes, in consequence of intestinal obstruction resulting from some mechanical occlusion (as well as for the removal of diseased ovaries, etc.), have been cut through, when other means have failed in affording relief, and in several instances with success, might sometimes tempt us to an operation; but, unfortunately, before such a procedure can be justly determined upon, the case has generally been too far advanced. It should be remembered, moreover, that there have been instances of obstruction which have resisted the most judicious medical treatment for two or three weeks together, and the patients have, at length, recovered without an operation. On the other hand, where there was neither vomiting nor inflammation present, they have been carried off within a few hours.

In certain cases of intussusceptio and strangulation, the surgeon, by an operation, might be enabled to relieve the abnormal state in the one, and divide or break down the bands which produce constriction in the other; but, in most cases of internal strangulation, as well as in a scirrhus-contracted condition of the tube, the operation would be wholly inexpedient. It is just to observe, however, that whenever such an operation has been determined upon, it should be undertaken in a warm atmosphere; and the less manipulation is used, the better. Of course, no one would be bold or mad enough to attempt an operation of this kind without a consultation.

It is a question to be decided, whether the place of constriction be situated in the small or large intestines. After

witnessing the several volvuli amongst the small intestines, in the instance where Mr. Abernethy with such ease unfolded the intussuscepted parts, I have sometimes been tempted to inquire, in certain cases of suspected intussusceptio, etc., whether an incision through the *linea alba* might not with safety be made. We know, however, that frequently the invaginated portion, in consequence of an inflammatory process having been set up, becomes firmly adherent; thus proving that, in a case of intestinal obstruction dependent on intussusceptio, internal strangulation, or constriction from adventitious bands, the attempt to overcome the evil by a surgical operation, in the manner we have proposed, can only be looked upon as an extremely hazardous one; and the operator might even be stamped as the murderer of his patient. Indeed, the operation of gastrotomy has been severely condemned by some, amongst whom is M. Hevin, formerly a celebrated surgeon on the continent,* as being exceedingly futile. Encouraged, however, by the marked success which has attended the practice of several others, if I felt *convinced* that a case of obstruction was dependent on intussusceptio, or strangulation from adventitious bands of recent formation, which threatened to extinguish my patient's life; and if I had ascertained the whole history of the case, besides having apprised the patient and his friends of the grave alternative, and having received the sanction of some skilful and experienced surgeon, I doubt whether I should hesitate to explore the peritoneal cavity by an operation, with the design of restoring the abnormal condition of the tube, and thus removing the cause of obstruction. Judging from the French writings, a celebrated continental surgeon has lately operated with distinguished success in some cases of this character; and it is recorded, that two cases out of three recovered.

The great danger we have to encounter in the operation of cutting through the walls of the abdomen, in certain forms of intestinal obstruction, is the wounding, to a considerable extent, the peritoneum; but is it not wounded in the operation of strangulated hernia? in that of paracentesis abdominis? or in the extirpation of diseased ovaries? In the operation of tying the external iliac artery, we are aware, the peritoneum is carefully pushed aside, and left intact; so, in some cases of intestinal obstructions, it has been proposed to make an opening into the colon from the lumbar region, where the bowel can be reached behind the peritoneum; and here the membrane is preserved. But I much question the propriety of such an operation, if an artificial anus is intended, since, if the patient survives the operation, he really must become an object afterwards loathsome to himself and to those around him.

When the obstruction proceeds from a scirrhus-contracted condition of the rectum, we can seldom do more than adopt palliative means. The forcible introduction of clysters, or of bougies of any kind, cannot be too severely reprobated; by such practice, the bowel is likely to be ruptured, and a false passage effected.

In the treatment of incipient stricture of the rectum, the most simple yet nutritious diet should be adopted, as milk, sago, beef-tea, jellies, etc., and the bowels kept relaxed. The cautious introduction of bougies of various sizes will sometimes be of advantage to the patient, by their gentle dilatation of the constricted part. Various have been the means adopted, or proposed, for the cure, or mitigation of the disease. Desault employed flexible rolls of lint coated with some ointment, and these he gradually enlarged. Morgagni adopted a mercurial plan of treatment. But when there is complete occlusion of the rectum, thus forming a barrier to the further passage of the faeces, medical and surgical treatment will generally prove equally unavailing. However, when the abdomen becomes tumid from the obstruction, with intervals of occasional severe pain, etc., hot fomentations may be applied, and enemata of various kinds, as the infusion of linseed with laudanum, etc., should be administered; but, although this plan may serve to lessen

* Recherches Historiques sur la Gastrotomie dans le cas de Volvulus, par M. Hevin.

the sufferings of our patient, he will ultimately die of starvation. The same result will occur when the disease is situated in other portions of the intestinal tube. Malignant tumours, likewise, which tend to cause an obstruction, can only be dealt with by treatment of the mildest character. Tumours of this kind, seated in the lower part of the rectum, with a broad basis, when attempted to be removed by the knife or ligature, usually terminate very unfavourably.

Should the obstruction be dependent on hernia, of course, when it is strangulated or incarcerated, the sooner reduction is effected the better. If alarming symptoms threaten, the measures which we are called upon to adopt must be prompt; the taxis, the warm bath of 95° to 100°, or the injection of a tobacco enema; or what is now preferred, the use of chloroform; each, or all of these should be employed without loss of time. Opium may be given, and venesection sometimes employed; but if these fail to allow us to return the protruded parts into the abdomen, I would, without hesitation, proceed to an operation. The operation for the relief of strangulated hernia has too frequently been considered a cruel and dangerous one. I do not, however, exactly coincide with this opinion, when the operation is *early* performed by one who knows his anatomy well: the many deaths which *have* followed the operation have been mainly owing to its having been too long delayed, and indeed until fatal symptoms had already commenced. I think I am borne out in this by the sad mortality resulting from cases of this kind, which have been brought into the wards of our great hospitals in London. I have been particularly struck with this: it is owing, as I have said, chiefly to the operation having been too long delayed; the cases have generally been trifled with before the sufferers are brought into the hospital. If I may be allowed to allude to my own experience, I have resolved to operate *early* in the cases which have fallen into my own hands, when other means of reduction had failed; and I am happy to say, with complete success. I am pleased to acknowledge having done so on one occasion in the presence of my friend in the chair, under circumstances too of peculiar difficulty; my patient was seventy-two years of age, and yet lives in the enjoyment of excellent health; there was a large protrusion of omentum and intestine (inguinal hernia), which had become entangled, besides which there were strong adhesions, not of recent formation. Now, if we consult the writings of the Parisian surgeons, we shall generally find that their practice is an *early operation*, and we know with what success.

Intestinal obstruction arising from an imperforate anus, or a partially impervious rectum, must be treated in the manner the case usually demands.

CASE I. I remember the case of a young man, which came under my late father's care, shortly after I had commenced my apprenticeship; but, from the long distance of time, five-and-thirty years, I am unable to do more than give a general outline of the case. The patient, a farming servant, took cold, and in a day or two afterwards complained of severe cutting pains across the umbilical region which extended downwards. Obstinate constipation ensued, and the pains occurred at intervals of every ten or fifteen minutes. Vomiting of bilious matter frequently took place. The usual plan of treatment had been pursued for some days: purgatives had been given—opiates and enemata administered; venesection was had recourse to: and fomentations were applied to the abdomen; yet no evacuations had been procured, even on the seventh day from the commencement. The pain became now more excruciating, and the general symptoms appeared more urgent. The abdomen became very resonant. Quicksilver was next given, but with no improvement. At length, it was decided to have the opinion of Dr. Alexander, then residing at Leicester. The case was concluded to be one of intussusception; and Dr. A. now directed a tablespoonful of quicksilver to be given every half-hour; adding, that this was the only chance left of saving the patient's life. At first, there was no remission of the symptoms; but, after more than seven pounds of mercury had been administered,

the obstruction gave way, and the patient's life was saved, by the bowels being well emptied.

REMARKS. The administration of quicksilver in these cases has been condemned by some. Mr. Hunter thought it might with propriety be given where the intussusception was *retrograde*; and that by the peristaltic action, the parts might become normal. However, he holds out a caution, that portions of the quicksilver may find their way between the containing and inverted bowel into the angle of reflection, and by pushing it further on, increase the disease it is intended to cure. But here we have an instance where the patient's life appears to have been saved by the use of quicksilver alone; but pushed, certainly, to an amount larger than I have found on record.

CASE II. Thomas Cluloe, a labourer, aged 54, was suddenly seized with severe gripings across the umbilicus, which, increasing, caused him to send for me. He appeared to be suffering from some severe obstruction in the intestines. There was continued urgent tenesmus, without evacuations; there were considerable fulness and resonance of the abdomen. The pain was severe at intervals, but not increased on pressure. Pulse 78, and feeble. The man had now been two days without an evacuation. I gave him calomel combined with extract of colocynth, the black draught, castor oil, etc. Enemata were administered; hot fomentations with turpentine were applied to the abdomen. No evacuations followed on the tenth day, and the symptoms becoming more urgent, with vomiting of bilious matter, I gave my patient two pounds of quicksilver in four doses, but without avail. On the fourteenth day, he took a large spoonful of *newly-fermented yeast* every half-hour; and before he had taken a pint, there was a plentiful evacuation. I have seen great benefit result in several cases of intestinal obstruction from the administration of newly-fermented yeast: this has appeared to overcome the obstruction, when other means have proved abortive.

CASE III. Miss K., a remarkably fine-looking young lady, about 23 years of age, had joined an evening party of friends. She appeared to enjoy excellent health, and exhibited all the vivacity natural to her age. After, however, she had retired to her chamber, the inmates of the house were alarmed during the night with hearing her screams. She was writhing with agony. There was intense pain over the umbilical region, which was increased on pressure. Constant nausea, without vomiting, was also attendant, as well as urgent tenesmus. Hot fomentations were applied over the abdomen; calomel combined with opium, and castor-oil were given in repeated doses; and enemata were administered, yet without any evacuations being procured, or remission of the symptoms. Leeches were applied over the abdomen, and venesection was had recourse to. Notwithstanding, however, the prompt treatment which had been adopted, the young lady's sufferings continued to increase, and she expired, in little more than twelve hours from the commencement of the attack.

As there was considerable mystery hanging about this case, as to the cause of the symptoms, their severity, and the suddenness of the attack, it was thought more satisfactory to make a *post mortem* examination. The abdomen was exceedingly tumid and resonant. On laying it open, the stomach and intestines appeared distended with flatus, and presented all the signs of incipient enteritis. The stomach felt somewhat hard, and its larger curvature was considerably discoloured. An opening made into its cavity at once explained the cause of all the symptoms during life. There was a mass of *unbroken* cherry-stones, which seemed, as it were, cemented together with other matter, completely blocking up any outlet by the pyloric portion. It was then recollected that Miss K. had eaten very heartily of some ripe cherries, with other *bonbons*, during the evening, and had foolishly neglected to cast away the stones. There was a mass of nearly a pound of these indigestible substances. It is difficult to learn whether anything more could have been done to save this lady's life; but a question suggests itself,—Would a powerful emetic have dislodged the foreign mass, had we been aware of its existence?

CASE IV. Elizabeth Barnett, a healthy-looking child, ten years of age, began to complain of great pain in her bowels, particularly at the lower part of the abdomen, on the right side. There were all the symptoms of intestinal obstruction present: obstinate constipation, with nausea, and vomiting of bilious matter. Pyrexia prevailed, the pulse being 120 and upwards. The pain at first appeared to be circumscribed; but, as the case advanced, the abdomen became exceedingly tumid, and was tender to the touch. Calomel, mixed with sugar, was given in small doses, with castor oil, etc. Leeches were applied over the seat of pain, with hot fomentations, etc.: but everything proved abortive, either to procure evacuations, or relieve the child's sufferings. She died on the third day. Permission was granted to examine the body. On reflecting back the abdominal parietes, the small intestines were found to be greatly distended with flatus, and exhibited patches of inflammation. A further search discovered that a *kidney-bean* had slipped into the appendix cæci: it had swollen considerably, in consequence of its locality, and caused inflammation of the appendix. This had extended to other portions of the intestines, and invincible obstruction was the direct consequence. The appendix had lost its usual curved form, from the bean having got into it as far as half its length.

REMARKS. In this case, it would appear, no medicinal treatment could be of any avail, and the diagnosis would necessarily be difficult and uncertain. The question, however, here suggests itself,—Suppose the cause of the obstruction could have been distinctly traced during life, would it have been safe, and should we have gained anything, by making an *early* opening through the *linea semilunaris*, and, by gentle manipulation, trying to dislodge the foreign body, and place it in the axis of the canal?

CASE V. Mr. David Ferriman, of Wymeswold, aged 45, a painter and stone engraver, having been out on business, in the month of January, 1829, called, late in the evening, on his return, at an inn for a glass of ale. He took out his purse to pay for it, and it appears that there were some loose characters sitting in the bar, who, observing Mr. F. had a sum of money in his possession, waylaid him after he had left the house. He was robbed of his money, and also his watch. The night was dark. The alarm created, however, was so great, that Mr. F. had no power to offer the slightest resistance; and such was the effect produced on his mind, that although he met two or three persons shortly afterwards, and even passed close by the residence of a magistrate, he could make no complaint. He was so confounded, that after he reached home, a length of time elapsed before he could be induced to give any account of what had happened. The shock, however, was too much,—so great as to interrupt some of the functions which are necessary for the preservation of life. There were no alvine evacuations afterwards. Intussusceptio had evidently taken place. There was, however, no real amount of positive suffering, except at long intervals. As the case advanced, the abdomen became exceedingly tumid, and there was occasional vomiting of bilious matter. The pulse continued steady nearly throughout: the tongue was thickly coated. Hot fomentations were repeatedly applied to the abdomen. Nearly all the list of purgatives had been tried, and quicksilver, with the administration of enemata, simple, as well as of tobacco, but without avail. The abdomen became tympanitic; hiccup supervened, with cold sweats, and death closed the scene on the fifth or sixth day. No *post mortem* examination was permitted.

CASE VI. Mr. William Pent, aged 50 years, a farmer, who resided about two miles distant, had always enjoyed excellent health. He rose as usual in the month of January, three or four years ago; and being a man of active habits, began to sweep away the snow which had fallen heavily during the night. The weather had been excessively cold, and a pond was frozen over where the cattle were wont to drink. Mr. P. was in the act of breaking the ice with a crew-bar, when he was suddenly seized with a severe attack of diarrhoea, attended with considerable

gripping. In a short time afterwards, I was summoned to attend Mr. P., and when I entered his room, he was writhing with agony. He bent his body forward; his hands were placed on the lower part of the abdomen, on the right side. He described the pain as being just above the inguinal opening, and as if his "*bowels were being torn asunder*." The symptoms were such as usually accompany a case of strangulated hernia; but a careful examination showed that the abdominal openings were quite free. There was urgent tenesmus, attended with nausea, but without vomiting; the pulse beating about 80, and feeble. Fearing inflammation, I took away fourteen ounces of blood from the arm, gave four grains of the chloride of mercury and ten grains of extract of colocynth, with repeated doses of castor-oil; but it was clear I had to contend with an invincible obstruction in the intestinal tube. Fomentations to the abdomen were constantly applied, and enemata were administered. The severity of the symptoms and pain continued unmitigated; no evacuations took place; and the patient expired at about nine o'clock the same evening, being about thirteen hours only from the commencement of the attack.

REMARKS. No examination was here permitted after death; but the fact of the patient having risen in his usual state of good health,—the sudden, yet severe attack of diarrhoea, succeeded immediately by intense pain in the lower part of the abdomen, on the right side, and circumscribed, without the least trace of hernia, convinced me that the case was one of intussusceptio, or internal strangulation, certainly terminating in an extraordinarily rapid manner. That the whole of the symptoms arose from cold, if the case was one of intussusceptio, there appears little doubt, the patient having, on the day previous, being very thinly clad, been exposed to a bitter cold north wind. It is well known that cold will bring on diarrhoea; and, spasm ensuing, intussusceptio will frequently follow.

CASE VII. About thirteen or fourteen months ago, I was requested to attend on Ann Kinton, aged 44 years, and the mother of a numerous family. I found her in bed on my arrival. She complained of great pain and fulness in the abdomen. There was frequent tenesmus, but it appeared there had been no evacuation during several days past, except a very small one, which was scybalated, and effected by very hard straining, attended with great pain. I learned there had been a constipated condition of the bowels for about two years, and that defæcation had never been effected without considerable suffering, and only once in seven or eight days; that latterly each stool had been of a very flattened shape, and frequently streaked with blood. Before this woman sent for me, she had taken a large dose of Epsom salts and infusion of senna, as well as some castor oil, but without the slightest relief, since no evacuations were procured, and the pain became severe. There was now constant nausea, without vomiting. The skin was cool, and the pulse beat 73, full, yet soft; the tongue was much coated, but moist. The abdomen was very large and resonant, without tenderness on making gentle pressure. The pain in the abdominal region was not now very distressing, and only occurred at long intervals. Still there were great restlessness and uneasiness, attended with an aching sensation across the lumbar region, and extending along the course of the sacrum. I prescribed chloride of mercury, combined with extract of colocynth and croton oil, besides the black draught. The vomiting, however, became more urgent, and thinking this might arise from the croton oil, I now omitted this, and the patient was directed to take no aperient except small and repeated doses of castor oil. Enemata were repeatedly administered: and two grains of calomel, combined with one grain and a half of opium, were given at bed-time. As no evacuations had been procured on the tenth day, I wished to try the effect of a tobacco enema, and I waited some hours to watch the result. There was the usual symptom of great prostration of the vital powers produced by this enema, but nothing passed off by the bowels except what had been injected *per anum*. It was now evident that some mechanical obstruction existed; but a digital examination detected nothing in

the rectum. If there existed a constriction in this part, it was higher than the finger could reach. When a small-sized bougie was attempted to be introduced, it met with resistance about six or seven inches high. Different sized bougies were carefully tried, but with the like result,—each became coiled, if much pressure was used. The patient's friends now wished to have the opinion of Dr. J. C. Williams, of Nottingham. This gentleman saw the woman on the evening of the 21st of August, 1851; there were no very urgent symptoms. All attempts, however, to give an enema proved ineffectual; it passed no further than the constriction. On the following morning, after Dr. Williams had seen the patient, I was resolved to try the introduction of O'Beirne's tube; it, however, met with the same resistance the other instruments had done, and returned in a coiled form. When I repeated my visit in the evening of the same day, to my great surprise, the poor woman stood looking out of her window, and said, "*I am taking another look round before I go, for I am dying in health.*" She then sat up, and took some tea with her family. As Dr. Williams had seen her, she now expressed a wish to try whether anything more could be done for her if she went to the Nottingham General Hospital, and asked me if I should object to this, and whether it would be safe for her to risk the journey, a distance of ten miles. Of course, to gratify her wish, I assented, provided she travelled in a covered conveyance, on a bed or mattress; at the same time, I assured her that little more could be done by medicine. On the following day she was no worse, and her wish was gratified by her being admitted into the Nottingham General Hospital. A consultation took place, and I believe attempts were made to pass some bougies, etc. Unexpectedly, however, on the following day, the poor woman ceased to breathe. No evacuation had been procured. Hearing of her death, I felt naturally anxious to know whether a *post mortem* examination had taken place, and if so, what were the appearances of the intestinal tube. Accordingly, I wrote to Dr. Williams, and received his reply, with an inclosed description of the morbid appearances. The following is a copy of a letter written by Joseph White, Esq., the resident surgeon of the hospital:—

"In the case of Mrs. Kinton the principal seat of stricture was in the rectum, about seven inches from the anal extremity; a smaller amount of contraction existed lower down, though this was easily passable. The colon was ulcerated in patches throughout the greater part of its length; though most extensively in the transverse portion. The other viscera were all healthy. The stricture would with difficulty admit the point of the little finger; and the bowel immediately above was thickened and corrugated, so as to form a valve, preventing the passage of any matter downwards."

CASE VIII. Samuel Marshall, aged 60 years, a bricklayer by trade, was suddenly seized with all the severe symptoms which accompany a case of strangulated hernia. His sufferings appeared horrible. He threw himself on the floor, and rolled about in agony; then he would as quickly rise, with his hands firmly pressed on the lower part of the abdomen; but no position or place gave him the slightest relief. He never had had hernia, but had frequently felt some severe cutting pains along the abdomen. Calomel, castor oil, and calomel combined with a full dose of opium, were given. Enemata were administered; no relief, however, was obtained; the symptoms presented a darker aspect, and the man died in less than twenty hours of intense suffering. An examination after death showed a constricted condition of the intestinal tube near the caput cæcum coli. This, it appeared, had been progressing for some years past, and fresh morbid matter being continually laid down, a tough and constricted band had been formed, which gradually becoming closer, strangulation of the gut ensued.

CASE IX. A few months ago, I attended a case of intestinal obstruction, arising from internal strangulation. A young woman had for several months before complained of cutting pains across the umbilical region. When I was summoned to visit her, there was considerable nausea, accompanied with

occasional severe pain in the abdominal region, and which extended downwards. I learned that she had been confined to her bed for more than a week before I was requested to see her; she had taken castor oil of her own accord; a few days before she took the oil, she had a severe attack of diarrhoea attended with much pain. She informed me that she had frequently felt a dense pain in the lower part of the abdomen, on the right side, and that constipation had prevailed to a considerable extent. There was now considerable tenderness from the right hypochondrium downwards, but without the least tumidity of the abdomen. I could distinctly feel a circumscribed hardness nearly opposite the cæcum, or descending portion of the colon; and it was evident that the locality of the obstruction was in or near to these parts. It was, however, with the greatest difficulty that an enema pipe could be introduced up the rectum, and very little injection would pass, being quickly returned. The mildest aperients, and calomel combined with opium, were given. To attempt an operation, as suggested in another part of my paper, even if in other respects it might appear practicable, in this instance was forbidden by the symptoms. Death occurred early on the third day after I first saw the patient. A *post mortem* examination proved that a portion of the descending colon was constricted by a firm band of lymph not of recent formation, and the appendix cæci had become attached at its extremity to the descending portion of the colon, and formed a ring, through which a portion of the ileum had passed and became strangulated. In most of these cases the abdomen is considerably resonant and tympanitic, but in this instance it was perfectly flat.

Wymeswold, Leicestershire, February 1853.

THE CLIMATES OF THE WORLD, IN REFERENCE TO THEIR EFFECTS ON MAN'S GENERAL WELFARE AND DESTINY.

By T. G. HAKE, M.D., Physician to the Suffolk General Hospital.

At the present hour, when millions of human beings are leaving their homes in Europe, or contemplate doing so, for countries which may or may not possess a climate similar to that which gave them birth, it cannot fail to be perceived that many interesting inquiries are thereby laid open to science, on which none but the physiologist can enter with advantage.

If the race of man, then, in all its varieties, affords a general expression of the existing climates of the world, in reference to the circumstances which contribute to healthy organisation; if one climate maintains, while another modifies more or less the characteristics of human beings, as is undoubtedly the case, it cannot be out of season to make this investigation: With what result to his own physical conformation and moral character, and more especially to that of his descendants does man, born in a congenial climate, emigrate at random into different zones?

Having examined this subject in its many bearings, both by collating the written observations of travellers made cursorily without any immediate scientific object, and by questioning intelligent persons from our colonies, and other newly formed states, I premise with confidence that it is one which is most important and curious. To take a common example;—What a diversity of effects must obviously arise from one member of a family taking up his permanent abode in a portion of the American, and another in a part of the Australasian continent. Both regions afford climates which are called delightful and healthy; but their separate influence on the organisation of the European is already perceptible. Indeed, it has been forcibly argued that man cannot travel out of his own zone without entailing extinction on his race. Be this as it may, an inquiry into the subject of isogenetic*

* I venture to introduce this term in order to mark the distinction between zones of equal geniality of climate, and those of like temperature, or isothermal zones, both of which differ from each other, and from geographical zones, as much as the magnetic differs from the common equator.