thyroid. The isthmus is usually prominent, thickened, and broadened. One lobe is commonly more enlarged than the other; occasionally the enlargement is unilateral (Figs. 5 and 6).

Microscopically the goitres have presented in detail the appearances described by Williamson and Pearse as characteristic of the progressive stage of lymphadenoid goitre in the human subject. It is exceptional to find pronounced retrogressive changes—fibrosis and atrophy—in the rat goitres, although these have occasionally been encountered in relatively mild degree. Lymphadenoid changes have also been observed in glands presenting no obvious signs of enlargement; they were confined to certain areas in an otherwise apparently normal organ.^{9 10 11}

Having had an opportunity recently to examine under the microscope sections of lymphadenoid goitre in man, side by side with others of experimentally produced lymphadenoid goitre in rats, the identity of the pathological state of the thyroid gland in man and in rats was apparent.

To the physician or surgeon who has to deal with goitre in the human subject the points of interest arising from these observations are: (1) lymphadenoid goitre occurs in man; it can be produced under experimental conditions in rats. (2) It arises in approximately 25 per cent. of young rats despite the adequate ingestion of iodine. (3) Deficiency of iodine in the food is certainly not the cause of lymphadenoid goitre as it occurs in rats; it is unlikely, therefore, to be the cause of lymphadenoid goitre as it occurs in man. (4) The basal factor in the causation of lymphadenoid goitre in rats is a dietetic one; it may reasonably be expected that the basal factor in its causation in man is likewise dietetic and of a like order. (5) Judging from experimental experience in rats the diets which are likely to be associated with lymphadenoid goitre in man are those composed largely of white flour or other vitamin-poor carbohydrate, vitamin-poor protein, and vitamin-poor fats and in which there is a paucity of fresh fruit and green leafy vegetables. In view of the almost universal use of white flour as the main staple of the dietary in Western countries, of the widespread use of vitamin-poor carbohydrate foods, of substitutes for butter which are relatively poor in vitamin A, of the use also of frozen and tinned meats, and of the small proportion of fresh fruit and green leafy vegetables in many people's dietaries, the etiological significance of the observations made on rats appears to me to be great.

REFERENCES.

- ¹ Williamson and Pearse: *Journ. Path. and Bact.*, 1923, xxvi, p. 459.
- ² Williamson, Pearse, and Cunningham: *Journ. Path. and Bact.*, 1928, xxxi, p. 255.
- ³ Williamson and Pearse: *Quart. Journ. Med.*, 1928, xxii, No. 85, p. 21.
- ⁴ Idem: *Journ. Path. and Bact.*, 1926, xxix, p. 167.
- ⁵ Idem: *Journ. Path. and Bact.*, 1925, xxviii, p. 361.
- ⁶ Simmonds: *Virchow's Archiv*, 1923, B. 245-246.
- ⁷ Tebbutt: *Supplem., Med. Journ. Australia*, August 20th, 1927, p. 54.
- ⁸ McCarrison: *Studies in Deficiency Diseases*, London, 1921.
- ⁹ Idem: *Indian Journ. Med. Research*, 1927, xv, No. 1, 247.
- ¹⁰ Idem: *Indian Journ. Med. Research*, 1927, xv, No. 4, 909.
- ¹¹ Idem: *Simple Goitres*, London, 1928.

THE PREVENTION AND EARLY DIAGNOSIS OF THE ACUTE ABDOMEN.*

BY

ZACHARY COPE, M.D., M.S., F.R.C.S.,

SURGEON TO ST. MARY'S HOSPITAL, AND TO THE BOLINGBROKE
HOSPITAL, LONDON.

THOUGH the title of my paper includes the early diagnosis of acute abdominal disease I wish to deal chiefly with the prevention of the acute abdomen. This convenient but slang term is generally taken to cover those acute conditions which come without warning, like a bolt from the blue, and it is seldom that a word is heard about measures for preventing such catastrophes. It is easy to see how this fatalistic attitude has arisen. In the course of practice the surgeon—who is largely responsible for current teaching—

sees the patient in the throes of an acute calamity, the slighter premonitory features have been overshadowed by the catastrophic culmination, and there is frequently no strong inclination to prolong the time between diagnosis and operation in eliciting a meticulously accurate history of preliminary symptoms. As an example of this attitude I could cite a reference to a book published in 1914 by a famous London surgeon who, in describing a case of appendicitis, said "no surgeon of experience pays much regard to history." Until this attitude is replaced by one of very careful study of all the antecedents of the more acute conditions we cannot hope to diagnose early, much less prevent the disease.

I will first consider the extent to which it is possible to prevent the onset of any acute abdominal disease, and secondly how to prevent any acute condition which has actually started from reaching a catastrophic stage.

Until the exact pathology of the various abdominal conditions is known it is, of course, impossible to take exact precautions to prevent, but by a careful attention to symptoms it is occasionally possible to detect serious disease before it is acute. The different diseases vary greatly in their preventability, but it is worth while considering the question in each group of cases.

Take first the question of perforated gastric and duodenal ulcer. Is it possible to prevent perforation? If we only consider the acute ulcer which develops within a few days and precipitately perforates, the answer must be No; but when we recollect that the majority of ulcers which perforate are those of long standing the reply should be a modified affirmative. Chronic ulcers can now be diagnosed by means of the x rays in the great majority of cases, and if the correct treatment is adopted it should be possible to prevent perforation. If I am asked, What is the correct treatment—is it medical or surgical?—I cannot tell unless the type of ulcer is specified, and who is to carry out the treatment. Some ulcers are, I think, curable by the alkali method or by means of the duodenal tube; others hold out no hope of cure other than by surgery. The patient should be kept under treatment and supervision until the radiological evidence of ulcer has disappeared and the symptoms have subsided.

An ulcer may continue for many years without preventing the patient carrying out his duties, so constant supervision is necessary. A medical friend of mine, when a student with me, had a very severe melaena and all the symptoms of duodenal ulcer. He was advised to have an operation, but refused, and for twenty years managed a busy practice, though constantly ailing with dyspepsia and latterly having to give up from time to time. He was supposed to be rather neurotic by those who had not seen the melaena and did not know his character. Eighteen months ago (and twenty-two years after the melaena), while on a motor tour in Scotland, he had all the symptoms of perforation—pain, collapse, etc.—though fortunately the symptoms subsided with rest. I saw him a little later. An x-ray examination revealed very extreme pyloric stenosis and an ulcer crater in the duodenum. At the operation I found a mass of fibrous tissue all round the first part of the duodenum, with a recent perforation sealed over by omentum. Now this patient, who was known to have an ulcer, waited twenty years for his perforation, and but for the kindly offices of a bit of omentum might easily have succumbed. Correct treatment of the ulcer at an earlier date should have prevented this.

It is a common occurrence for there to be an exacerbation of symptoms for a few days before the perforation of a chronic ulcer, and this fact should be pointed out to all those who are known to have an ulcer and yet will not submit to give up the time necessary for the correct treatment to be carried out. Rest should be insisted upon in times of acute exacerbation of the symptoms.

The most common acute condition in the abdomen is appendicitis, and it is no exaggeration to say that any method of preventing that disease would not only save two thousand lives annually in this country, but would also more than halve the number of acute cases upon which the surgeon would need to operate. Appendicitis is no new disease, for in the College of Surgeons can be seen an

* An address delivered to the Croydon Division of the British Medical Association, October 16th, 1928.

account, written by John Hunter's own hand, of a *post-mortem* examination performed by him on the body of a colonel who had died with symptoms of intestinal obstruction, but in whose abdomen Hunter found only a perforated appendix with peritonitis. Yet it cannot be doubted, either on looking into old *post-mortem* records or examining the statistics of cases of appendicitis and peritonitis in later years, that the disease has become much more frequent in the last three decades. If we could but find the reason of this, prevention should be within our power. An exhaustive inquiry into the possible causes of this increase was made comparatively recently by Mr. Rendle Short. He came to the conclusion that the increase in the incidence of appendicitis was most likely due to some change in the dietary of the nation, but his evidence was against the common assumption that increased meat-eating had anything to do with the increase of the disease. By comparing the common articles of diet at the present time with those most in use before the great increase of appendicitis Short found that the chief change was a gradual diminution in the amount of cellulose-containing foods. He suggests that herein lies the cause of the greater predisposition to, and incidence of, appendicitis.

I think it will be allowed that appendicitis is a microbial disease which attacks certain people who are at the moment of attack in a state of diminished resistance to that infection. It will hardly be questioned that stagnation of the intestinal contents and the presence of much decomposing meat in the intestines are factors which would favour the growth of the infective organisms. It may be that cellulose-containing foods help to prevent stagnation of the intestinal contents, and they may possibly supply some vitamin which maintains resistance against the organisms. The moral to be deduced from this is that, so far as our knowledge goes at present, a predisposition to appendicitis would be lessened by (1) preventing constipation, (2) minimizing the amount of meat eaten, and (3) increasing the amount of food containing cellulose.

A fourth suggestion is to diminish the total amount of food taken. It is generally agreed that most people in this country eat more than is necessary, and often more than is good for them. Now it has been shown by Rendle Short that in institutions such as prisons and some children's homes, in which a low diet is given, the incidence of appendicitis is much lessened; while Horsley Gantt, in his very interesting account of the classes of disease which attacked the starved and neglected Bolshevik masses after the revolution in Russia, pointed out that appendicitis, cholecystitis, and similar inflammatory diseases of the intestinal tract almost disappeared, although most varieties of infective disease were rife at the time. I believe that appendicitis will once more diminish in frequency if some of the suggestions here given are followed up.

In considering diseases of the gall-bladder it is wise to deal separately with gall-stones and cholecystitis. It is not at all uncommon to find an inflamed gall-bladder without any gall-stones, and though gall-stones are usually accompanied by inflammation, that is not necessarily the case.

Cholecystitis apart from gall-stones has a close relationship with appendicitis. The two conditions are sometimes present together, and may occur at different times in the same person. It is quite likely that the same causes will tend to produce both diseases, and this is supported by the remarkable and parallel diminution in their incidence in Russia after the revolution. It has lately been shown that the infection of the gall-bladder may be primary in the wall of the viscus, a fact which testifies to the route of the infection being by way of the blood stream. In the prevention of cholecystitis and appendicitis, therefore, the importance of dealing with any focus of infection which might furnish a source of origin of organisms which could travel by the blood stream to the abdominal viscera must be remembered. The throat, mouth, and accessory sinuses should particularly be attended to.

The regular emptying of the gall-bladder is important if infection is to be prevented from obtaining a hold. Though the exact mechanism of the emptying of the viscus is not yet determined with certainty, there is strong evidence, furnished by Lyon and Meltzer, to show that the lower end of the common bile-duct is relaxed under the influence of

magnesium sulphate applied round the orifice in the second part of the duodenum. Though some of the magnesium sulphate taken by the mouth may be altered in the stomach, yet it would appear rational to recommend the occasional taking of Epsom salts by those who are thought to be predisposed to disease of the gall-bladder.

The actual prevention of gall-stones depends not only upon the prevention of infection, but also on the prevention of hypercholesterinaemia and the avoidance of too much inspissation of the bile. Our knowledge of the formation and metabolism of cholesterol is at present too vague for any definite dietetic precautions to be taken with any chance of success of avoiding the formation of gall-stones. The encouragement of the flow of bile has been mentioned above.

Of the other common intra-abdominal inflammations salpingitis due to infection with the gonococcus is a preventable infection in theory, but practically the question is not such an easy one. The female is not usually infected from an acutely inflamed male urethra—for obvious reasons—but the trouble arises from the awakening of a chronic infection which has been latent in the prostate or seminal vesicles, or possibly in the crypts of the urethra itself. It is of the utmost importance, therefore, that all infections of the male genital tract should be as far as possible eradicated, and permission to marry should not be given unless and until the infection has been destroyed. I am aware that this is a counsel of perfection, for the channels of the genital tract are devious and labyrinthine, but at all events a series of stringent tests should be made to determine whether the infection due to the gonococcus is still present or not. Only in this way shall we be able to diminish the number of those cases of acute salpingitis following soon after marriage—cases of truly tragic import. It is not sufficient to examine the discharge obtained from the urethra after prostatic massage. An artificial congestion must be produced by passing a full-sized bougie, or by injecting into the prostatic urethra a few minims of a solution of silver nitrate, 5 grains to the ounce, and then examining the discharge. A big dose of vaccine should be given to induce a local reaction, and some have even recommended the administration of an amount of alcohol sufficient to produce urethral congestion. One examination should never suffice; several examinations at spaced intervals are necessary, and even when every examination has been made there can be no absolute certainty as to the absence of the organism, which may lurk in so many crannies in the tract. It is sometimes possible to destroy local foci of infection by means of the operating urethro-scope.

Another form of acute infection which invades the female genital tract is that which leads to pneumococcal peritonitis. The work of Fraser and others makes it fairly clear that the pneumococcus gains entry through the vagina and Fallopian tubes, and since the condition chiefly attacks girls between 5 and 10 years of age we have a clear indication for prevention—that is, to promote local cleansing and hygienic measures so as to prevent the entry of the organism. The carrying out of these measures would devolve upon those who have the care of young children, and the medical man's part is to instruct those concerned.

The prevention of intestinal obstruction is impossible in some cases, problematical in others, but easily attainable in the majority of instances. This statement may cause surprise until it is recollected that the majority of cases of intestinal obstruction are due to the strangulation of external herniae. It is clearly easy to prevent strangulation of an external hernia by undertaking a radical cure of the condition. There are three or four factors which have a bearing on the question of operation in such cases—age, reducibility, control by a truss, the position of the hernia, and the general constitutional condition of the patient.

When a hernia is strangulated operation is undertaken whatever may be the age of the patient, and in any hernia which is irreducible, painful, or prone to attacks of sub-acute obstruction, operation should in general be recommended irrespective of age. It can be performed under gas and oxygen combined with local anaesthesia with very

little risk to the patient. There is no definite age at which it can dogmatically be said that the sufferer is too old for operation. If a hernia is easily reducible, and can readily be controlled by a truss, the danger of strangulation is greatly reduced, and there is no need to urge operation for such herniae in people over middle age who have no strenuous work to do. If the truss does not control, then the risk of operation is no greater than the risk of strangulation.

Of the three common kinds of hernia—inguinal, umbilical, and femoral—the first-mentioned is not quite so dangerous as the other two. The explanation of this depends probably on two factors. In the first place, umbilical and femoral herniae issue through rings with hard, fibrous margins, which have no “give” in them and make reduction more difficult; secondly, there is a greater nerve supply in the inguinal region, and strangulated herniae there are as a rule much more painful than those in the other situations. It is seldom possible to overlook a strangled inguinal hernia, but not so difficult for the patient and any unobservant person to miss a small strangled femoral hernia. Again, umbilical herniae usually occur in very fat people, and sometimes a small hernia may be embedded in the fat and hardly visible on the surface. There is all the more reason, therefore, to urge operation on even small herniae in the umbilical and femoral regions so as to prevent a strangulation which might easily be overlooked until too late. There is one exception to this statement—that is, the case of any enormous irreducible umbilical hernia of long standing occurring in a feeble patient. By operating and forcing the herniated mass into the general abdominal cavity the intra-abdominal pressure may be unduly increased, with considerable shock, embarrassment of respiration, and possibly fatal result.

Herniation through a weak scar of the abdominal wall is another preventable condition. It is not always possible to prevent the occurrence of a hernia in an operation scar—though much may be done by careful suturing and the limitation of drainage—but it is certainly possible to keep an eye on the scar for some time after operation, and to detect any weakness which may be remedied in time to prevent any risk of strangulation of the abdominal contents.

Intussusception is a disease of healthy babies who are attacked about weaning time, when the change from liquid to more solid food is taking place. Such strong babies are more likely to be favoured with “a little bit of what we have” at an earlier date than the more weakly, and this may sometimes account for its occurrence even from the fourth to the sixth month. Given an ileo-caecal region which is furnished with a longer mesentery than usual, an infant whose intestinal muscle is strong and equal to vigorous peristalsis, and a large food bolus sufficient to stimulate excessively the as yet uneducated ileum, and it is not difficult to see how an intussusception is produced. Indeed, it is strange that the condition is not more common. The only factor which can in any way be controlled is that of diet. Too sudden a change should not be made in the diet of a child aged from 6 to 18 months; let the transition from liquid to solid be graduated; also, do not let solid food be given too early simply because the child is stronger than usual. See, also, that the milk-clots in the child's stomach and intestine are made small by administering some sodium citrate by the mouth. Keep the bowels open so as to avoid constipation and straining.

Many forms of intestinal obstruction are almost impossible to prevent because the preliminary symptoms are so very vague and the physical changes so slight that nothing except actual vision can determine their importance. Cancerous stricture of the gut, peritoneal pouches causing intraperitoneal herniae, strangulation by Meckel's diverticulum, and a puckering of the mesentery and intestine by the attachment of a calcified lymph gland can hardly be prevented, though intra-abdominal adhesions following operations can, to a limited extent, be prevented by care in manipulation of the intestines, reduction of drainage to the minimum, peritonizing any raw surfaces, and attending to the details of suturing the abdominal wall.

In the case of recurring abdominal pain of a subacute colicky type, however, we have the barium x-ray examination to help in detecting any kink or stricture in the intestines, and though this method is very much less helpful than some enthusiasts would have us believe, yet sometimes it may lead us to operate and detect obstruction before it becomes acute.

Having considered a few measures which can be taken in order to prevent the onset of acute abdominal disease, I will now briefly comment on the early diagnosis of an acute condition which has started. I could not, even if I wished, do more than skim the surface of this very important problem, but I can best provoke discussion by making an extreme statement of the surgical attitude. *There should be a time limit for the making of a diagnosis in a case of acute abdominal disease.* I do not mean to suggest that it is always possible to diagnose a case in a few hours, but I do assert that any case of severe abdominal pain which lasts for more than six hours is serious enough to merit the calling in of a surgeon—not necessarily with a view to operation, but to exclude those forms of acute abdominal disease which need prompt operative intervention if they are to be successfully treated. I particularly put in the proviso about operation, for I believe the wise surgeon is recognized as much by his knowing when to operate as by his operative skill. I hope I am not presuming when I venture to think that the surgeon sees a much greater number of acute abdominal cases than any other branch of the profession, and if he is worth his salt he should be able to give a sound opinion on those cases.

I have often been astonished by the stoicism of patients, who will bear the most agonizing pain for many hours in the hope that it will subside spontaneously, and I have occasionally been surprised at the vicariously stoical attitude of some doctors when faced by a patient in pain. There are many acute abdominal pains for which a dose of morphine is the correct treatment—such, for example, as renal and biliary colic, gastric crises, diaphragmatic pleurisy with pain referred to the abdomen—but there are other conditions of an apparently similar nature for which to give a dose of morphine is, to say the least, an unwise, and, to say the most, a possibly fatal, procedure. It is generally possible to distinguish between these groups, and such a distinction should be made at as early a stage as may be done.

I have taken the time limit with reference to pain because that is the only constant symptom, but it is rare for that to be the only symptom. What I would insist upon is that any other common symptom is not to be regarded as essential. I have frequently had the statement made to me, “I did not think it could be serious because of the absence of a certain symptom”—the absent symptom varying according to the case. Now the common symptoms found in acute abdominal disease are: pain; nausea or vomiting; local tenderness; local muscular rigidity; distension; hyperaesthesia; shock; alteration of pulse, respiration, or temperature; furred tongue; constipation or obstipation; altered facies. There is hardly any absolutely constant symptom, but it may be safely asserted that pain is by far the most common, and is present in the vast majority of all acute abdominal conditions. All the others are most variable.

I think if I were asked which was the most important help in diagnosis after pain I should say altered facial appearance. I cannot describe the exact changes which the countenance undergoes, for this can only be learnt by experience, and some have great difficulty in acquiring this knowledge. To learn when a patient is really ill should be the first step in the diagnosis of acute abdominal disease, but it is astonishing how seldom this knowledge is acquired. Ward sisters often have more knowledge of this than the resident officer. There is one time, however, when the face does not help quite so much—that is, when the latent stage has supervened a few hours after the start of the disease.

Vomiting is frequently absent in appendicitis, often occurs only once immediately after the perforation of an ulcer, is not to be expected in obstruction of the large

bowel, and may even be absent in acute obstruction of the small intestine when gangrene quickly follows obstruction; it is not a constant symptom in the rupture of an ectopic gestation. Nausea is more constant, but not so striking a symptom.

Distension must never be waited for, and should not be present in the early stages except in obstruction of the large intestine. I well remember a man being sent to St. Mary's Hospital some years ago by a doctor who wrote: "Last night I thought this man very likely had a perforated ulcer, but he was not distended, so I left him alone; this morning he is distended, so I send him up to you."

Alteration of the pulse rate is of value when noted, but nearly every kind of acute abdominal disorder may be present without an appreciable alteration of the pulse. In some of the acute colics the pulse may be slowed; in perforated gastric or duodenal ulcer the pulse after two or three hours generally becomes quite normal; in intestinal obstruction of either the small or large intestine the pulse is frequently normal or slowed until toxæmia begins or unless there is severe shock on account of strangulation. In appendicitis I expect to find a normal pulse, unless (a) there is commencing peritonitis, or (b) the appendix is distended with toxic material. I have even known a normal pulse with acute pancreatitis, though I must admit that this is very exceptional.

I need hardly dwell on the *variability of temperature* in acute abdominal disease. A drop of several degrees accompanied by acute pain in the abdomen at once proclaims the occurrence of a catastrophe, however quickly the patient may regain a normal appearance. But many conditions cause no fall in temperature. Fever is absent in the acute obstructions, is moderate in amount and rather slow in onset in the peritoneal inflammations, and is high only in those conditions where infection is occurring in small cavities such as the pelvis of the kidney and the biliary ducts. I make it a rule in any case of acute abdomen with a temperature of 104° or 105° to turn first to the kidneys and then to the lungs and pleura.

Constipation is too vague a symptom to be of much use in early diagnosis. The same remark applies to furred tongue, though it is often possible to learn a lot by looking at the tongue.

An *increase in respiratory rate* must make one examine the lungs very carefully, but this increase quite often occurs in acute abdominal disease, so it is not of such great help.

Local muscular rigidity, though frequently referred to in the textbooks as a regular sign of appendicitis and peritonitis, should never be waited for. It is seldom present when the inflammation has not pierced through the walls of the appendix, and even when peritonitis has commenced muscular rigidity is absent when the inflammation affects a silent or undemonstrative part of the peritoneal cavity—for example, the pelvis or central part of the abdominal cavity. I wonder how many lives have been lost by ignorance of this simple fact.

We come, then, to the conclusion that the diagnosis of the presence of acute and serious abdominal disease is decided chiefly by the occurrence of pain—spontaneously complained of by the patient or elicited by local pressure on certain parts of the abdomen. The pain must be investigated from every standpoint—time of onset, character, duration, severity, position, variation in position, whether it occurs on breathing, passing urine, or defaecation. Finally, the practitioner should ask whether it is referred to any part other than the primary part, and observe how the patient turns over on to his side. Local tenderness must also be carefully sought; this examination is assisted by a careful study of the face. All other symptoms are observed, but it is generally possible to say whether the condition is serious or not by the simple observations I have described.

The doctor must make up his mind within six hours or less as to the seriousness of the condition. If he is doubtful he should not be ashamed to get a colleague to help in diagnosis. He must never procrastinate, for procrastination is the thief of lives.

TREATMENT OF ASTHMA BY RADIATION.*

BY

S. GILBERT SCOTT, M.R.C.S., L.R.C.P.,

D.M.R. AND E.CAMB.,

DIRECTOR OF THE RADIOLOGICAL DEPARTMENT OF THE LONDON HOSPITAL.

THIS disease, or rather symptom of disease, must still be regarded as an unsolved medical problem. As Professor Wild states: "Asthma is one of the most fascinating diseases for everyone except the patient." Theories have been advanced from time to time as to possible causes. The numerous so-called cures that are advocated simply indicate that no one "cure" is effective.

While not proposing to discuss the many different theories or even to advance a new one as the possible cause or causes of asthma, I will briefly outline one that is at present gaining ground, as it is helpful in explaining the effects obtained by radiation (x rays), administered by the method about to be described.

It is now generally accepted that asthma is not primarily a lung disease, but that some condition outside the lung is responsible for the spasm of the bronchial tubes. Abnormalities of the upper respiratory tract, such as nasal polypi; infection, such as sinusitis; chronic appendicitis; ovarian irritation; intestinal worms; the presence of cats, dogs, horses, feather cushions; the intake of certain foods, and many other agents, fantastic and otherwise, are known to give rise to an attack of asthma in different individuals. But should not these be looked upon rather as *triggers* that let off the asthmatical gun than as the primary causes of asthma? In more scientific language, do they not start a reflex, peculiar to asthmatical patients, in some organ or organs, which upsets or disturbs the metabolic or endocrine balance?—a somewhat nebulous term, but indicating a complex condition, which is best described as a loss of balance between the various ductless glands. The investigations of many workers tend to prove that one gland is, in some indefinable way, dependent on the quantity and quality of the secretion of another for its proper functioning. If a break occurs in this cycle, either a nerve reflex or otherwise, the whole balance is upset, and this is made manifest in many and various ways.

It is conceivable that each gland, when at fault, gives rise to its own particular set of clinical manifestations. Is it possible that in asthma the secretion of the suprarenal glands is defective? This somewhat speculative theory is mentioned here, for I believe that, in some manner, general radiation influences the secretion of the ductless glands. In other words, the beneficial results may be achieved by restoring the metabolic balance.

X rays have been used by many workers as a therapeutic agent in asthma, but the available literature on the subject shows that the results are, on the whole, disappointing. I believe two technical faults are responsible for this failure—*too small* a radiation field and *too large* a dose. The majority of authors treat only the thorax, using single front and back areas of about 10 by 15 cm. and the short wave (high voltage) therapy.¹ Others have found that some cases respond better if the spleen alone be radiated in the same way.² Even the skull has been radiated for this disease.

The important factors in the method of radiation about to be described are *the use of a large field and small dosage*. I wish to emphasize this combination, for, so far as I can find out by examining the literature on the subject, it has not been previously used in asthma.

As the beneficial results were discovered accidentally, it is difficult to say with certainty upon which organ or organs influence is being exercised. The whole trunk, back and front, is therefore irradiated, using the long wave therapy (low voltage). It is interesting to note that results have been obtained by radiation of the *abdomen* alone, but not by the radiation of the *thorax* only. The possible biological effects of a combined large field and small dosage have yet to be investigated, and the budding research worker should

* Read at the Second International Radiological Congress at Stockholm, 1928.