ABBREVIATED BIBLIOGRAPHY.

- ¹ On the Artificial Ripening of Cataracts. *Trans. Ophthal. Soc. U.K.*, vol. 10, p. 24 *et seq.*
- ² Jessop, W. H.: The Prognosis in Incipient Cortical Opacities of the Lens. *Trans. Ophthal. Soc. U.K.*, vol. 34, p. 151 *et seq.*
- ³ Duke-Elder: *Recent Advances in Ophthalmology*.
- ⁴ Hopkins, Gowland: *Biochem. Journ.*, 1921, 1923, 1925.
- ⁵ Friedenwald, J. S.: Permeability of the Lens Capsule. *Arch. Ophthal.*, 1930, vol. 3, p. 182.
- ⁶ Tassman and Karr: Glutathione in the Crystalline Lens. *Arch. Ophthal.*, 1929, vol. 2, p. 431.
- ⁷ Adams, Dorothy R.: The Nature of the Ophthalmic Lesions produced Experimentally by Naphthalene. *Brit. Journ. Ophthal.*, vol. 14, p. 49.
- ⁸ Burdon Cooper: Aetiology of Cataract. *Brit. Journ. Ophthal.*, vol. 6, p. 437.
- ⁹ Dor, L.: Le traitement médical de la cataracte. *La Clinique Ophthal.*, June, 1919.
- ¹⁰ Cantonet, A.: *L'ionisation de l'oeil*, Paris, 1927.
- ¹¹ Smith, Henry: *The Treatment of Cataract*, 1928, and *Trans. Ophthal. Soc. U.K.*, vol. 48, p. 89.
- ¹² Ellis, Z. H.: Non-operative Treatment of Cataract, with Report on Lens Antigen Treatment. *Arch. Ophthal.*, 1928, vol. 57, p. 46.

THE DIAGNOSIS OF CANCER OF THE LARYNX.*

BY

EDWARD D. D. DAVIS, F.R.C.S.,

SURGEON, NOSE, THROAT, AND EAR DEPARTMENT, CHARING CROSS HOSPITAL.

EXTRINSIC laryngeal cancer or cancer of the pharynx which has extended to the larynx usually comes under observation so late that the diagnosis is obvious. These cases are generally seen by the laryngologist much too late for successful treatment either by operation or by radium, and earlier diagnosis is urgently needed. The problem of early diagnosis is intensified by the fact that cancer in itself does not produce any specific signs or symptoms, and it is only when the cancer cells form a tumour or ulcer that a series of mechanical symptoms and signs appear. These symptoms also develop slowly and insidiously, and it is unusual for the patient to consult a doctor until such symptoms are well marked and until it is often too late for a successful removal of the growth. Moreover, in a hidden cavity like the pharynx where there is room for the growth to expand unseen, the first symptoms are trivial and may be neglected until secondary growths are manifest in the cervical glands.

The intrinsic cancers of the larynx are seen at an earlier stage, as the early symptom of hoarseness sends the patient to the doctor much sooner, but cancer seen in its earliest stage is more difficult to diagnose accurately. Hence cases of intrinsic laryngeal cancer present more frequent and more difficult problems in accurate diagnosis. I propose, therefore, to spend most of the time available in discussing the diagnosis of intrinsic cancer of the larynx. It is to be hoped that the more frequent use of the modern directoscope and direct laryngoscope, and suitable cancer propaganda, will produce the much needed earlier diagnosis of extrinsic cancer. The earlier stages of these cases may then present as many problems in diagnosis as the intrinsic cases.

Extrinsic Cancer.

The characteristic and offensive epitheliomatous ulcer commencing just below the aryteno-epiglottidean fold and rapidly involving that side of the larynx, produces fixation of the vocal cord with its consequent hoarseness. The first symptoms are sore throat and pain radiating to the ear, with a definitely tender spot near the greater cornu of the thyroid cartilage; then follows the characteristic hoarseness or roughness of the voice, with a copious frothy blood-stained expectoration. The top edge of the growth can be seen with the laryngoscope and palpation with the finger reveals the stony hard ulcer.

A post-cricoid carcinoma, which commences as a superficial growth on the posterior aspect of the cricoid cartilage in women, soon surrounds the lumen of the pharynx and spreads upwards to form the whitish warty upper edge seen in the later stages behind the oedematous arytenoids. These patients first complain of a soreness or pricking sensation with pain in the ears during swallowing. Pressure on the cricoid backwards against the spine or lateral movement of

the larynx produces pain in practically all cases. Dysphagia occurs early and when these symptoms are present without a satisfactory explanation the patient should be examined with a directoscope or oesophagoscope. The laryngoscopic mirror will not show any signs in the early stage, and when the upper edge of the growth is visible the disease is advanced, and matters are still more hopeless when a vocal cord is paralysed. Neither is the growth within reach of the longest finger, and the passage of a bougie is valueless and even harmful. In the absence of physical signs, the dangerous diagnosis of neurasthenia is still made; but such a diagnosis should never be concluded until the presence of a growth has been eliminated by a direct examination.

Carcinoma in the vallecula, either on the base of the tongue or on the anterior surface of the epiglottis, excavates the base of the tongue and rapidly extends to the surrounding structures, and the submaxillary and deep cervical lymphatic glands of both sides are soon involved. The only symptoms are persistent severe pain and a catch on swallowing located by the patient to the base of the tongue, with blood-stained salivary expectoration. Occasionally a chronic paroxysmal cough is the first symptom. Later, the painful and limited protrusion of the tongue with the accompanying trismus impedes laryngoscopic examination, but the growth is within easy reach of the palpating finger and does not require a skilled laryngologist or endoscopist to detect and gauge the extent of the growth. The situation, the characteristic appearance, the odour, and the history of these growths make the diagnosis more or less obvious; but the differential diagnosis should always be considered, as I have seen two cases of syphilis in the post-cricoid region which have been mistaken for carcinoma.

Intrinsic Cancer.

A rapid and accurate diagnosis of a typical case of intrinsic cancer of the larynx is easily made. The characteristic miniature epitheliomatous wart or ulcer situated on the middle or anterior third of the vocal cord in a patient of cancer age, in which the differential diagnosis of chronic laryngitis with pachydermia, tuberculosis, syphilis, and a simple tumour can be undoubtedly excluded, presents no difficulty. Such a growth has been and is frequently removed by operation without a pre-operative diagnostic section. On the other hand, the atypical case or the hidden subglottic growth requires considerable ingenuity in diagnosis. The differential diagnosis may be obscure, and tuberculosis is more often mistaken for cancer than any other condition. A chronic tuberculous ulcer of the vocal cord in a man of cancer age, or a healed scarred tuberculous lesion which has broken down with a recrudescence of the disease, closely resembles cancer.

Lupus, owing to its distribution in the soft tissues, is more readily distinguished; but in both cases tubercle bacilli cannot always be found in the sputum. In a few cases of tuberculosis it is almost impossible to make an accurate diagnosis without microscopy. The guinea-pig test by inoculation from the piece removed unfortunately takes six weeks to mature, but it is so conclusive that it might be useful. Chronic laryngitis with localized pachydermia renders diagnosis difficult, but the long history, the type of patient, the slow progress with a tendency to a wide distribution, gives the clue.

Syphilis is readily differentiated from cancer. The presence of other syphilitic lesions, the history, and the Wassermann reaction indicate the correct diagnosis; but when cancer is grafted on an old syphilitic lesion, the diagnosis may be difficult and can only be settled by the aid of the microscope.

Benign or simple tumours are recognized by their appearance, their situation, and slow growth. These tumours are usually removed by intra-laryngeal methods, and if the section for microscopy in any way suggests malignancy a more extensive operation can be done. Occasionally a malignant tumour has been labelled by the section to be benign, or a papilloma has recurred and has become locally malignant.

Sarcoma of the larynx is fortunately very rare. I have only seen two cases, both in young adults. The tumours were smooth, sessile, globular, and vascular, and occupied

* Read in opening a discussion in the Section of Laryngology and Otology at the Annual Meeting of the British Medical Association, Winnipeg, 1930.

the inner surface of the aryteno-epiglottidean fold and ventricular band. The tumours bulged into the pyriform fossa of the pharynx and rapidly increased in size. Chevalier Jackson states that not one true sarcoma was found among 643 neoplasms of the larynx seen at his clinic, and there are only three specimens of sarcoma of the larynx to be found in all the museums of the London medical schools. The diagnosis of sarcoma is completed by exploration and a diagnostic section, and a survey of the literature of such cases shows the tumours to be either round-celled or spindle-celled sarcomata.

The subglottic growth, often found below the anterior third of the vocal cord, cannot be clearly seen by the direct or indirect method of examination. A heaviness in movement or fixation of the vocal cord occurs comparatively early in subglottic growths, but our President, Sir St. Clair Thomson, has shown that it is undoubtedly a late and unfavourable sign in the manifest cancer. The diagnosis of such cases of cancer should be made before fixation of a vocal cord is evident. The subglottic growth is more malignant and is quickly disseminated to the cervical lymphatic glands—therefore an early diagnosis is urgent and the "wait and see" policy cannot be adopted.

The directoscope and Jackson's anterior commissure laryngoscope with retraction of the vocal cord outwards now enable us to obtain a better view of the subglottic growth and the subglottic extension of a more manifest growth.

There is still some controversy as to the value of the removal of a portion of the growth for diagnostic purposes. Our President in July, 1922, recorded that in 37 out of 51 of his cases, the diagnosis was completed and proved to be accurate without the removal of a piece for diagnosis. In the remaining 14 cases the diagnosis obtained by a diagnostic section was misleading in two cases. A diagnostic section is a valuable but not an infallible aid to diagnosis, and it should be emphasized that a negative section should be ignored and considered to be useless. The value of a diagnostic section is increased if a sufficiently large and deep portion from the centre of the growth is removed. The whole portion should be carefully sectioned and searched from the surface to the depths and a number of sections made. For mechanical and other reasons, it is frequently difficult to obtain a satisfactory or sufficiently deep piece for section; but we are now better equipped with Jackson's tissue forceps and improved direct laryngoscope. There is an impression that the direct method of examination should be more frequently employed and every opportunity should be taken to make use of it. For example, when an anaesthetic is given for the extraction of teeth as a preparation for operation, information can be obtained by a direct examination.

If the diagnosis is in any way doubtful, a diagnostic section is imperative and hastens the diagnosis. If the wound produced by the removal of a portion does not heal, or if the tumour increases in size, it is strong evidence in favour of cancer. I have never seen any harm arise after the removal of a diagnostic piece, but the picture may be entirely altered and diagnosis made more uncertain in the case of a negative section. No doubt in suitable cases time will settle the diagnosis, and judicious waiting, with an examination once a week or at even longer intervals, will finally reveal the accurate diagnosis; but patients and their friends are apt to become restive and demand to know whether it is cancer or not. It is not always expedient to delay by waiting, but there are cases in which it is advantageous to wait and to be cautious. The delay, which should be as short as possible, gives an opportunity to have the teeth and mouth cleaned up. It serves to get the patient in training for an operation and a second opinion can be obtained if desired. It is often useful and expedient to have a second opinion; it increases the patient's confidence in his surgeon and may be helpful.

If these methods of diagnosis fail in a case of suspected cancer, it is justifiable to perform an exploratory laryngofissure with full permission to proceed to a radical operation. Where the growth is obscured by oedema of the mucosa or by inflammatory changes, such an exploratory operation is a necessity. It has been done, and no harm to the patient

has accrued, particularly in those cases in which tuberculosis has been excluded. Exploratory laryngofissure has of course been frequently done to ascertain the extent of the growth and to decide whether a partial or complete laryngectomy is necessary.

Conclusions.

1. An earlier diagnosis of extrinsic cancer of the larynx is urgently needed.
2. Time will settle the diagnosis in some cases of intrinsic cancer, and it gives an opportunity to prepare the patient for operation, etc.
3. The direct method of examination is an aid to diagnosis and will ascertain the extent of the growth.
4. A diagnostic section of an intrinsic cancer is a valuable but not infallible aid to diagnosis. Such a section should be made in all doubtful cases.
5. An exploratory laryngofissure is justifiable when all other methods have failed.

Memoranda:

MEDICAL, SURGICAL, OBSTETRICAL.

AN UNUSUAL CASE OF INTESTINAL OBSTRUCTION.

The following case we believe to be unusual as we cannot find reference to similar cases in the literature, at our disposal.

F. K., aged 27, a reservist ambulance orderly, was admitted to hospital at 10.30 a.m. on January 16th, 1930. He gave the history that the previous evening at about 7 p.m. he had partaken of a heavy meal, almost immediately afterwards being seized with acute abdominal pain. He vomited at the time and continued to do so throughout the night. The following morning he reported sick at his medical inspection room and was at once sent to hospital for admission.

On arrival the patient was markedly collapsed, with a pinched face and a pulse uncountable and imperceptible at the wrist. The respiration rate was 22 and the temperature 97.7° F. The abdomen was distended, rigid, and tender, especially in the epigastrium; movements were practically nil. The patient had not passed urine and the bowels had not been open since before the onset of the attack; a gentle enema gave no result. He was at once given brandy 1/2 oz. and pituitrin 1/2 c.cm., and prepared for operation.

The abdomen was opened by a mid-line subumbilical incision. Loops of slightly distended but much engorged small gut presented, and the abdominal cavity contained a considerable amount of blood-stained fluid. The trouble was eventually traced to the sigmoid loop, which was long and mobile, and which had swung across the abdomen to the right side, where it had herniated into a large retrocaecal fossa. It was tightly gripped in this position by a tense peritoneal band extending from the abdominal wall behind the caecum to the lower end of the attachment of the mesentery of the ileum. The sigmoid loop was tightly stretched but was not twisted. It was found that in taking up this position it had come to lie across the mesentery, obstructing the blood supply of the ileum for six or eight feet to within one foot of the ileo-caecal junction, where the structures, including the appendix, were quite normal.

The constricting band of mesentery was divided and nine inches of strangulated, distended, and engorged sigmoid colon withdrawn; this might have recovered, but the viability of the small gut was very doubtful. The ileum above the obstruction was turgid but not greatly distended; the portion involved was blown out, but the bowel from below the obstruction round to the sigmoid was nearly empty and normal in every way. Continuous intravenous saline was given throughout the operation, but at this stage life was almost extinct and it was considered advisable to close the abdomen. The patient died about one hour later.

The points of interest in this case seem to be the sudden onset, the absence of any marked distension, and the nature of the double obstruction. There were no adhesions to suggest a congenital malposition of the sigmoid. A possible explanation is that the small intestine was the first to suffer, allowing the contents of the large bowel to travel on till they reached the sigmoid loop, where they were held up, and so led to the second strangulation.

B. K. SIKAND,
Captain I.M.S. (T.C.).
E. VENN CLAYDON,
Lieutenant I.M.S.

Rawalpindi.