CASE VII.—*Type V Bacteraemia with Recovery.*

This case belonged to Type V; the blood culture was positive on the eighth day, though the temperature was below 102°. This patient either sterilized his own blood, or there was sufficient protection for Type V associated with the Type II serum. There were probably only 400 units of Type V protection given in the serum. (See Chart 3.)

CASE VIII.—*Type V Bacteraemia mistaken for Type II: Fatal.*

This patient was a married woman who, on the third day of her illness, had a positive blood culture and sputum; the organism was reported from another laboratory as Type II. On the fourth day she had a miscarriage at eight months. She received large doses of Type I and Type II serum. On the fifth day there was ample antibody for Types I and II. Her blood culture continued positive; the serum was unavailing. The original blood culture, which had been reported Type II, was then re-typed, as well as a blood culture after ample antibody for Type I and Type II had been administered and was demonstrable in the blood, and was found to be actually Type V.

CASE IX.—*Type II Bacteraemia: Fatal—"Influenza Pneumonia."*

This patient came in on the influenza service of the hospital. When a consolidation was discovered on the sixth day a blood culture was taken. He was given 15,000 units of Type II serum, and was placed in the oxygen chamber. Serum was administered more actively—as much as 60,000 units, 80,000 units, and 60,000 units being given. In spite of this intensive treatment antibody was not detectable in his blood. The colonies in his plate did not increase, but he continued to become more poisoned, and finally died on the twelfth day of his disease. It is probable that, in this case, if more intensive treatment had

been given at the very onset, sufficient to produce a positive agglutination test and maintain it, the result might have been different.

It will be seen from these cases that the problems connected with the serum treatment of pneumonia are numerous. Many of them are unsolved. Those of us who are working in hospitals, with ample laboratory assistance, have much for which to be grateful. I have shown the difficulties so that it should not be assumed that the serum treatment of pneumonia is a finished chapter; but I have also shown the triumphs, so that practitioners may be impelled to use serum in the treatment of their patients. If it is employed generally, so that the toll of pneumonia is lessened, as I believe it will be, the purpose of the Littauer Pneumonia Research Fund of New York University will be accomplished, and those who have devoted themselves to these problems will know that they have not worked in vain.

BIBLIOGRAPHY.

- Jesse G. M. Bullowa: Use of Antipneumococcic Refined Serum in Lobar Pneumonia. Milton B. Rosenbluth: Relation of Bacteremia in Lobar Pneumonia to Prognosis and Therapy. William H. Park and Georgia Cooper: Antipneumococcus Serum in Lobar Pneumonia. *Journ. Amer. Med. Assoc.*, April 28th, 1928, pp. 1349-1358.
William H. Park, Jesse G. M. Bullowa, and Milton B. Rosenbluth: The Treatment of Lobar Pneumonia with Refined Specific Antibacterial Serum. *Journ. Amer. Med. Assoc.*, November 17th, 1928, pp. 1503-1507.
Jesse G. M. Bullowa: Lobar Pneumonia and its Treatment with Refined Sera. *Colloid Chemistry*, by Jerome Alexander, vol. 2.

THE CARE OF RHEUMATIC CHILDREN.*

BY

A. DINGWALL FORDYCE, M.D., F.R.C.P.Ed.,

HONORARY PHYSICIAN, ROYAL LIVERPOOL CHILDREN'S HOSPITAL, AND TO THE BIRKENHEAD AND WIRRAL CHILDREN'S HOSPITAL.

For the past three years I have held, both at the Liverpool and the Birkenhead Children's Hospital, a special out-patient clinic for rheumatic children; to this clinic school medical officers have sent considerable numbers of cases, and practically all children have been recommended by doctors. Through the co-operation of the respective child welfare associations home conditions have been investigated and reported upon, and regular supervision of the children obtained. Most of the children have passed through one or more of the in-patient branches of the Liverpool or Birkenhead hospitals, and their supervision continued thereafter. Acute cases have mainly been dealt with in the city hospitals, while long-continued residential care has been carried out in the Heswall and West Kirby institutions. Most of these latter cases have been at the Heswall Hospital, some for many months, and all have had the benefits of special schooling and graduated exercises in addition to medical and nursing care. The physical condition of the children has been investigated as fully as possible, and attention paid both to general treatment and as regards points of focal infection. Reports have been sent to the respective chief school medical officers dealing with the actual physical condition—and particularly the condition of the heart—and recommendations made as to the desirable nature of school attendance or type of school.

Up till March 31st of this year 540 children had attended, a few of them since before the commencement of a special clinic and the others for varying periods since. At this date (March 31st) these cases could be classified thus: 321 children with apparently no permanent cardiac lesion, 119 children with an apparently established cardiac lesion but with satisfactory myocardial response, 25 children in whose cases I was still uncertain as to whether cardiac damage was permanent or not, 41 children with cardiac lesions and failing heart, 9 cases of death from rheumatic carditis, 9 children suffering from acute rheumatic fever, and 16 cases of congenital heart deficiency.

Excluding cases of congenital heart deficiency and all cases suffering at the date (March 31st) from acute rheumatic fever, the outcome of whose illness hung in the balance, and giving the benefit of the doubt in prognosis to all cases of doubtfully permanent cardiac lesion, I find

that of 515 cases medically recommended to the clinic as rheumatic, and accepted by me as such, 346 could be regarded as suffering from no permanent heart lesion, 119 definitely had a permanent lesion but showed good response, 41 showed evidence of failing heart muscle, and 9 had died. In other words, 50 children—10 per cent. of cases—had died or were unlikely to survive childhood, and another 119 children—25 per cent. of cases—were permanently crippled to a greater or less extent.

I put forward these figures with no further object than to show that rheumatic infection in childhood is a common and dangerous disease on Merseyside, and I cannot be accused of making a case for the rheumatic child, as these children were all sent to me as rheumatic or suspected rheumatic. During the first three months of this year—to March 31st—273 children attended the clinic on one or more occasions, and the average attendance was about 25.

In all, approximately 400 reports have been sent to the chief school medical officers, the form of each report being as follows:

RHEUMATISM AND ACQUIRED HEART DISEASE.

Name..... Sex..... Date of Birth.....

Address..... School.....

Place of Examination.....

History, symptoms, and physical signs.....

CLASSIFICATION OF CASE:

A.—Acute pyrexial rheumatism.

B.—Acute chorea, i.e., chorea of less than two months' duration.

C.—(1) Rheumatism or chorea other than that included in A or B; with no heart lesion, or apparently curable heart lesion.

(2) Apparently permanent acquired heart lesion—good prognosis.

(3) Apparently permanent acquired heart lesion—bad prognosis.

RECOMMENDATIONS:

1.—Fit for Public Elementary School.....

A.—No restrictions.

B.—No violent games.

C.—No drill or gymnastics.

2.—Fit for Special P.D. School.....

A.—Day School.

B.—Residential School.

3.—Not fit for any School.....Period of exclusion.....

Signature of Examiner.....

Date of Examination.....

* A paper read at a meeting of the Liverpool Division of the British Medical Association.

These recommendations are sometimes not acted upon by the authorities because the facilities do not at present exist, but I hope they may assist ultimately in the framing of schemes for administrative advance. The number of cases is large, and their continued supervision a matter of very considerable difficulty.

The serious figures I have given, showing that in 35 per cent. of the cases death or permanent cardiac crippling supervened, may be subjected to criticism from two aspects. In the first place, it can be urged that chiefly serious cases were sent to the clinics, and this is no doubt true to some extent. But it is not altogether so, because school medical officers were informed that it was *rheumatic* children with whom it was desired to deal, and they knew that the underlying effort was the prevention of heart disease. The other criticism that can be put forward is that these children were not efficiently cared for. This criticism I also, regretfully, accept in part, and shall deal with some details in a moment; but this must now be said—that exceptional care and attention have been paid to these children, no pains have been spared in their treatment, and in many instances the degree of their improvement has been remarkable. I am definitely of opinion that the development of specific rheumatic clinics with associated home visitation and regular supervision has improved the prognosis.

There are some cases in which we are forced, in spite of every conceivable form of treatment and all facilities, to watch the child passing from the early infected period to that of grave cardiac disease and ultimately death. But such cases should be rare, and are rare if the early signs of the first active infection are noted and dealt with. When these early signs are noted close care for a prolonged period is essential. But in a series of 52 consecutive cases of acute rheumatism recently admitted to the Children's Hospital, Liverpool, in only 18 instances was it possible for direct transference to another institution—country hospital or convalescent home—to be carried out, while of the total 52 cases 75 per cent. were retained in the City Hospital for only a month or less. Efficient therapeutic prophylaxis was here evidently not carried out; but, owing to the need for beds in town for acute cases, it is impossible to retain these children in city hospitals, and sufficient accommodation is not provided for them in country hospitals. Many patients have to wait months for admission to the Heswall Hospital, and frequently such waiting period leaves marks of serious damage.

The new Local Government Act will bring about many changes and open up many possibilities. I am sure that no more important development can occur than the setting aside of specific beds or wards for the prolonged care of rheumatic children.

As regards the active care of the children, I would lay great stress on the removal of infected tonsils. Of the 531 rheumatic children alluded to in this paper, 130 underwent tonsillectomy in hospital and 22 at a school clinic; in 4 cases parents refused permission for operation, and in 67 cases operation is desirable and probable. Thus in 213 out of 531 children—or about 40 per cent. of cases—I have considered tonsillectomy advisable.

Contrary to what some may think, I consider that in being responsible for these figures I have acted in a conservative manner. I have considered as indications for operation:

1. Obviously unhealthy tonsils—irrespective of size.
2. The presence of rheumatism with a history of tonsillitis as initial symptom.
3. The presence of rheumatism with recurring attacks of nasopharyngeal catarrh associated with exacerbations of active rheumatism.
4. Cases of active rheumatism running a prolonged course resistant to all medical measures, whatever the appearance of the tonsils.

In none of the cases operated on did unfavourable symptoms develop from the operation, however active the rheumatic symptoms at the time; while in most cases obvious and marked benefit accrued. It is, however, impossible to consider the operation as an isolated factor in treatment, as good results from it can only be justly

expected when enucleation is complete, when the immediate after-care of the child is as thorough as after a major operation—which this is—and when, after suitable institutional care, home supervision is so prolonged as to tend to prevent or at least deal early with subsequent intercurrent infections.

I cannot but feel that the term "acute rheumatism" in childhood has been stressed unduly in the effort to obtain accurate delimitation. Our foe is *active* rheumatism, and, perhaps unfortunately, its symptoms are not always acute. It is usually easy for the clinician to make a diagnosis between rheumatic infection and the varied rheumatoid conditions; and syphilitic cardiac conditions are rare at this age. It is the early diagnosis of active rheumatic infection which is so important, which is sometimes difficult, and which, I think, is sometimes delayed because of the common use of the term "acute" rather than "active."

The early diagnosis and suitable care of active rheumatism give good results. Fortunately for the children of the well-to-do, frequent medical supervision and constant educated home care serve to prevent acute ebullitions and disastrous end-results even without the need of accurate diagnosis. For, however, the children of the hospital class, once they show symptoms of active rheumatism, and whether they have been previously classed as "dyspeptic," "hypoglycaemic," "pre-acidotic," "intestinal," "toxic debilitated," or simply "pre-rheumatic," I consider the value has been fully demonstrated of specific hospital clinics and associated home visitation and prolonged supervision by private practitioner or through the clinic.

I also consider that tonsillectomy, efficiently carried out and succeeded by suitable and prolonged care, takes a very important place in treatment, and that the need of adequate institutional beds for prolonged care is clamant.

HYSTERO-SALPINGOGRAPHY IN THE DIAGNOSIS AND TREATMENT OF STERILITY AND PELVIC DISEASE.*

BY

R. S. STATHAM, O.B.E., M.D., CH.M.,

HONORARY GYNAECOLOGIST, BRISTOL ROYAL INFIRMARY; LECTURER IN OBSTETRICS, UNIVERSITY OF BRISTOL.

In a previous paper on the subject of sterility¹⁷ I emphasized the great importance, in all cases where no definite cause could be found for lack of conception, of ascertaining the patency of the Fallopian tubes before treatment was begun. At that time I was using the Provis-Rubin method of CO₂ insufflation, and I instanced two cases of pregnancy following on the clearing of the so-called "sticky tube"; since that date I have met with four further successful cases, in each of which the patient became pregnant within six weeks of the insufflation. The object of this paper is to further the claims of a method of diagnosis and treatment which is, in my opinion, far superior to the gas insufflation, good though that is. This procedure consists of injecting lipiodol into the uterus and tubes under the control of an x-ray screen, and then the taking of a series of skiagrams. It is equally valuable in diagnosis and treatment.

Before coming to the technical details it may be of interest to give a short history of this method of diagnosis. As far back as 1914 Carey⁸ in the United States attempted to pass collargol through the tubes, but got no practical negatives, while Dimier in Paris recorded a death while using the same procedure. These results caused the method to be abandoned. In 1915 Rubin¹⁵ published a paper on collargol injections. In 1923 Kennedy¹³ made use of sodium bromide, and in the same year we also tried it at the Bristol Royal Infirmary; but these methods gave poor and unreliable results, and a few cases of peritonitis were recorded.

The discovery of the suitability of lipiodol for the injection material was the means of establishing this technique on a sound basis. Carlos Hueser¹¹ and Carelli,⁵ during

* From a lecture with lantern slides delivered to the Bath and Bristol Branch, February 27th, 1929.