tolerated, and in milder conditions the bulk of the carbohydrate may be taken as bread. The following skeleton diet, which may be modified to suit the particular case, is in use at the Royal Hospital.

Meat, fish, poultry, eggs, clear soup, bovril, broth, cabbage, cauliflower, lettuce, endive, tomato, grape fruit, asparagus, mushrooms, sea-kale, cucumber, chicory, spinach, celery (*as desired*).
Tea, coffee, and unsweetened lemon juice in water (*as desired*).

Compulsory Ration.

Benger's food made with milk half pints during the day.
..... units of insulin are to be taken before each half-pint of *Benger's food*.

Bread: ounces during the day, with meals. Butter as desired.

(If specially ordered) *Jam or preserve*—half an ounce with bread, at a meal.

A further units of insulin is to be taken before of the daily meals, one of which must be the one at which the jam is taken (if ordered).

Total daily insulin dosage units.

Carbohydrate approximately grams.

CARE OF DIABETICS BEFORE AND AFTER OPERATION.

In operations upon diabetic patients it is a matter of the utmost importance to avoid ketosis. For a clean operation, such as an interval appendix or radical cure of a hernia in a well-controlled diabetic, few special precautions are necessary. Where, however, there is sepsis, as in the case of amputation of an extremity for wet gangrene or the excision of a carbuncle, it is well, if possible, to delay the operation until the diabetes has to some extent been controlled. The following procedure, subject of course to modification to suit the individual case, is the routine at the Royal Hospital.

1. Three days prior to operation increase the carbohydrate content of the diet to at least 100 grams.
2. Operate early in day.
3. Three hours before operation give carbohydrate content of usual breakfast *plus* 10 grams of glucose. Give usual morning insulin at same time.
4. Simple enema one and a half hours before operation.
5. Immediately before operation give 10 to 20 units of insulin.
6. On returning to ward give 30 to 40 grams of glucose as 10 per cent. solution in saline, rectally, with 20 to 30 units of insulin.
7. Two hours after operation take blood specimen and give glucose or insulin according to result.

The diet in early days after operation should be liquid and soft, and should largely consist of carbohydrate until the temperature is normal and manifestations of toxæmia have disappeared. The total calories should not exceed maintenance diet during the period of active healing.

In conclusion, I would insist that the successful treatment of diabetes demands the whole-hearted co-operation of the patient. He must have an intelligent appreciation of the condition from which he is suffering, and time spent in explaining this to him is time well spent by his medical attendant.

COLLES'S FRACTURE:

AN EFFICIENT AND ECONOMICAL METHOD OF SPLINTING.

BY

W. MITCHELL SMITH, M.D., C.M.,

HONORARY ORTHOPAEDIC SURGEON, CRIPPLES' HOSPITAL, STOKE-ON-TRENT.

THE cost of after-care (massage, physiotherapy, and electrotherapy) in the treatment of fractures has been mounting up in recent years to an alarming extent. The time has come for the surgeon who prescribes this treatment—often as a matter of routine—to consider whether this expense is justified, or whether equally good, or even better, results might not be obtained by well-considered splinting.

Efficient splinting should ensure complete immobilization of the fragments after reduction, and, in the case of the Colles's fracture, fixation of the hand in such a position that the patient is able to use it as soon as the damage to the soft tissues begins to subside. Jones's and Carr's splints are the two most commonly used. The former provides the more perfect fixation of the fragments; the latter the better position from which to start re-education. I find that the

best features of both splints can be combined in a carefully applied plaster case.

The treatment of a Colles's fracture may be divided into three stages: (a) reduction of the fracture; (b) splinting; (c) after-care.

A. Reduction of the Fracture.

Reduction of the fracture under an anaesthetic can generally be easily obtained by steady traction on the hand and manipulation of the fragments into position by the thumb and forefinger. If more force is required the well-known manipulation of Sir Robert Jones provides all that is necessary; but as little force should be used as possible. Complete reduction is indicated by the restoration of the radial curve, as described by Sir Robert Jones, and of the correct relationship of the styloid processes.

B. Splinting.

A roll of stockinette is run up the arm, extending from the knuckles to the elbow. Over this a light plaster case is applied. With the thumb and forefinger of one hand the surgeon holds the fragments of the radius in the corrected position, and with the other hand holds the wrist in about 15 degrees of dorsiflexion and slight ulnar deviation while the plaster sets. It is essential that the plaster should be closely moulded to the dorsum of the wrist and that the angle of dorsiflexion should be exactly opposite the joint. To prevent rubbing the skin, the plaster should extend, glove fashion, half-way up the metacarpals, and should allow full palmar flexion of the metacarpo-phalangeal joints. (Figs. 1 and 2.)

FIG. 1.

FIG. 2.

FIG. 3.—Showing strip cut from plaster.

The occurrence of pain and swelling of the fingers may be avoided by cutting out from the back of the plaster, before it sets, and from the underlying stockinette, a strip about half an inch wide, leaving a bridge at the top and the bottom (Fig. 3). A radiogram to confirm the correction should be taken through the plaster.

C. After-care.

As soon as he recovers from the anaesthetic the patient is encouraged to use his fingers and gradually to develop the use of his hand. In a week or ten days he is able to do anything within the limits imposed by a cock-up splint. The plaster case is removed at the end of a month, by which time bony union has generally taken place, and as

the patient has already been using his hand freely for two or three weeks no further treatment is required.

I have been treating Colles's fractures by this method for the past three or four years with uniformly good results and with considerable saving in the cost of treatment.

HYPERSPLENISM AND HYPOSPLENISM AND SPLENECTOMY.

BY

F. PARKES WEBER, M.D., F.R.C.P.

APART from the results of traumatism and the exceedingly rare primary malignant neoplasms and some of the primary (or "predominant") tuberculous and other chronic infectious (parasitic) conditions of the spleen, the indications and counterindications for splenectomy will doubtless be more definitely formulated when more is known about the various functions of the organ in question. In regard to the functional activity of the spleen it will probably become more and more convenient to speak of "hypersplenism" and "hyposplenism"—by analogy with the terms "hyperthyroidism" and "hypothyroidism," "hyperpituitarism" and "hypopituitarism." It will probably also be found that there are various forms of hypersplenism and hyposplenism corresponding to involvement of the various functions of the spleen (and reticulo-endothelial system), apart from the now well-recognized function of the spleen as a blood reservoir (Barcroft, etc.).

In cases of harmful (not compensatory) hypersplenism splenectomy or some effective depressor of splenic function (radium, x rays, and other as yet undiscovered means) will be indicated. At present splenectomy is not rarely (sometimes with excellent result) performed for conditions, such as congenital haemolytic jaundice, thrombocytopenic purpura haemorrhagica, and the Gaucher type of splenomegaly, where there is more or less clinical evidence of hypersplenism. For cases of hyposplenism some effective activator of the spleen and reticulo-endothelial system will have to be searched out.

In regard to splenic infections it should be remembered that, though one of the functions of the spleen, as of the tonsils, is to resist and "hold up" infections, the spleen sometimes, like the tonsils, may become so overwhelmed as itself to constitute a harmful focus of infection—a state of affairs for which splenectomy may constitute a rational part of the treatment. In some such cases the bulk and altered consistence of the enlarged spleen may themselves become a source of danger for the individual (possibility of rupture from relatively slight traumata).

Under infectious and parasitic splenomegalies, besides the relatively transitory enlargements due to the ordinary acute febrile infections, one would place the more chronic ones due to malaria, syphilis, tuberculosis, hydatid cysts, kala-azar, schistosomiasis (one form of "Egyptian splenomegaly"), the problematic splenic "mycoses," and perhaps certain other cases that have been clinically labelled as "Banti's disease."

Memoranda:

MEDICAL, SURGICAL, OBSTETRICAL.

THE ANATOMICAL METHOD OF OPENING QUINCY.

Not everyone has been satisfied with the old method of opening quincy by cutting into the soft palate. The oedema present often makes it difficult to find the pus, and more than one cut may be necessary. On the other hand, immediate enucleation is so drastic a form of treatment as to appear unnecessary as well as risky. Both of these extremes can be avoided by opening the abscess in the right way.

A quincy is a collection of pus in the supratonsillar fossa. It forms because of inflammatory adhesions at the normal

orifice of the fossa. As pus accumulates the palate bulges and is further swollen by oedema. In order to evacuate the pus it is only necessary to break down the adhesions by introducing a pair of sinus forceps with closed blades into the supratonsillar fossa in the direction shown by the arrow in the diagram. The point of insertion is immediately

under the arch formed by the anterior and posterior faucial pillars on the same side. A slight push is needed. As soon as they have penetrated the abscess the blades are widely opened, the adhesions are broken down, and pus pours out.

There are many advantages in this method:

1. It is simple, certain, easy, and only one instrument is required.
2. Trismus is no obstacle.
3. There is no pain beyond pressure from the forceps points.
4. There is hardly any bleeding because nothing is cut.
5. The abscess is freely drained by its natural outlet.
6. The tonsils can be enucleated without risk at a later date.
7. There is no unnecessary mutilation of mobile parts.
8. No anaesthetic is necessary, either local or general.

London, W.

H. MORTIMER WHARRY, F.R.C.S.

HEREDITARY ANGIO-NEUROTIC OEDEMA.

THAT heredity plays an important part in the causation of Quincke's disease is well established, and numerous authors have reported genealogies. The hereditary form would seem to be of a much more severe type than the comparatively common acquired cases, and this is illustrated by the cases of Griffith,¹ where both father and daughter died of acute oedema of the larynx. J. R. and T. R. Crowder² report five generations of a family, and of 63 known descendants 27 inherited it; of these, 14 died as a direct result. Although a number of cases are on record, the hereditary form would appear to be of sufficient rarity in the individual experience to justify a case report.

A woman, aged 55, was referred to me on account of angio-neurotic oedema of lifelong duration; the disease was of the cutaneous type, and there was a complete absence of any of the puzzling attacks of abdominal colic due to intestinal oedema. The attacks occurred every two or three weeks, and the reason that advice was sought was the alarming increase in severity and frequency of the laryngeal involvement; the areas of oedema were peculiar in the absence of itching. Physical examination revealed little apart from a slight degree of dermatographism—much less than is usually seen in the common variety of case.

The patient was considerably apprehensive as to the outcome of the disease, as her mother and her only two brothers died of it; in an effort to avert the family tragedy the patient was desensitized by a course of peptone in small doses, and she was instructed to ingest half a grain of ephedrine sulphate at the first warning of an attack. Up to the time of writing the results have been most happy.

The following is the genealogical tree.

An asterisk indicates that the individual was affected. D = Death from oedema of the larynx. M = Male. F = Female.

ERNEST BULMER, M.D.,

Assistant Physician, General Hospital, Birmingham.

¹ Griffith: *British Medical Journal*, 1902, i, 1470.

² Crowder, J. R. and T. R.: *Arch. Int. Med.*, 1917, 20, 840.