had developed subsequently to the zoster eruption. The question naturally arose whether the encephalomyelitis was due to the zoster virus or not, but since both the vesicle fluid collected from the eruption and the cerebro-spinal fluid fixed complement in the presence of known zoster convalescent scrums, there was little doubt that the case was in fact one of zoster encephalomyelitis. It is true that the reaction given by the cerebro-spinal fluid was not as strong as that obtained with the vesicle fluid, but it was nevertheless quite definite, and cerebro-spinal fluids from cases of encephalomyelitis complicating diphtheria and scarlet fever gave entirely negative results.

Recently I have had the opportunity of examining the serum of individuals who had recovered from psittacosis for the presence of specific complement-fixing antibodies, and here again I have obtained evidence of a specific reaction. Psittacosis may not be so rare a condition as is imagined, and apart from extensive outbreaks like the recent one, where the number and severity of the cases inevitably attracted attention, sporadic cases probably occur from time to time and pass unnoticed. The inoculation of animals is too slow a method to be of practical use as a diagnostic procedure, and a specific serum reaction would be a great help in clearing up the diagnosis.

It may be said that the examples given are both of rather rare conditions, but the principle is nevertheless of general application. In this brief and rather sketchy survey of the present position of research in the realm of filterable viruses an attempt has been made to show that something of practical import has been achieved. Undoubtedly an enormous amount remains to be done, but the number of workers in this field is steadily increasing, and on every hand a determined effort is being made to gain a clearer understanding of this group of viruses. Some of the problems seem almost insoluble, but it must not be forgotten that the commonplace in bacteriology to-day must have appeared equally obscure to those who preceded Pasteur.

REFERENCES.

- ¹ Olitsky, P. K., and Gates, P. L.: *Journ. Exper. Med.*, 1922, 36, 501.
- ² Mills, K. C., Shibley, G. S., and Dochez, A. R.: *Ibid.*, 1928, 47, 193.
- ³ Steinhard, E., and Lambert, R. A.: *Journ. Infect. Dis.*, 1914, 14, 87.
- ⁴ Parker, F., and Nye, R. N.: *Amer. Journ. Path.*, 1925, 1, 325.
- ⁵ Carrel, A., and Rivers, T. M.: *C. R. Soc. de Biol.*, 1925, 65, 848.
- ⁶ Maitland, H. B., and Maitland, M. C.: *Lancet*, 1928, ii, 526.
- ⁷ Muckenfuss, R. S., and Rivers, T. M.: *Journ. Exper. Med.*, 1930, 51, 149.
- ⁸ Dochez, A. R., Shibley, G. S., and Mills, K. C.: *Proc. Soc. Exper. Biol. and Med.*, 1929, 26, 562.
- ⁹ Vallée, H., Carré, H., and Rinjard, P.: *Rev. gén. de méd. vet.*, 1926, 35, 129.
- ¹⁰ Bedson, S. P., Maitland, H. B., and Burbury, T. M.: *Journ. Comp. Path. and Ther.*, 1927, 40, 1.
- ¹¹ Laidlaw, P. P., and Dunkin, G. W.: *Ibid.*, 1928, 41, 1 and 209.
- ¹² Fairbrother, R. W.: *Brit. Journ. Exp. Path.*, 1930, 11, 43.
- ¹³ Gordon, M. H.: Medical Research Council, Special Report Series, No. 88, 1925.
- ¹⁴ Burgess, W. L., Craigie, J., and Tulloch, W. J.: *Ibid.*, 1928, No. 143.
- ¹⁵ Bedson, S. P., and Bland, J. O. W.: *Brit. Journ. Exp. Path.*, 1929, 10, 393.

MIGRATION AND HEALTH.*

BY

J. S. PURDY, D.S.O., M.D., C.M., D.P.H.,
F.R.S.Ed., F.R.G.S.,METROPOLITAN MEDICAL OFFICER OF HEALTH, SYDNEY, NEW
SOUTH WALES.

TO-DAY the distribution of the white population throughout the British Empire is so important that it warrants consideration not only from an economic and social point of view, but also from the point of view of health.

It is perhaps as necessary for Great Britain as for any country other than Japan to organize emigration on an extensive scale, on account of over-population. After or during every great war, there appears to have been a "stock-taking." Thus, after the South African war, the English Inter-Departmental Committee in 1904 showed that sections of the people, more especially in the more congested centres of population, were liable to suffer from the ill effects of overcrowding, or an artificial, unnatural environment. As a matter of fact, there was no evidence of actual physical deterioration, since no previous standard for comparison existed.

Actually the average span of life has advanced ten years during the present generation; and man to-day has excelled

* A paper contributed to the Section of Medical Sociology and History of Medicine at the Annual Meeting of the British Medical Association, Winnipeg, 1930.

in feats of physical endurance and established records surpassing those of earlier periods. We do know, however, that a considerable proportion of people are below an optimum of health and efficiency, physical, mental, and moral. Description in Great Britain, the United States, and New Zealand, during the later stages of the war, brought to light some startling statistics. Although it is possible to prove anything by statistics—even the truth—yet it is certainly disquieting to learn that "in the average, out of every nine men of military age in Great Britain, only three were found fit; two were upon a definitely infirm plane of health; three were incapable of undergoing any extra exertion, being practically physical wrecks; and the remaining man was a hopelessly chronic invalid." In Great Britain alone, a million men examined at the recruiting depots were found to be not sufficiently fit to be worth training for war service owing to structural defects or organic disease.

In regard to conditions to-day, the following passage occurs in Mr. and Mrs. Sidney Webb's monumental work, *The History of the Poor Law*:

"Is there any condition more sordidly tragic than the lot of the youthful unemployed—to-day counted by the hundred thousand, under-nourished, physically undeveloped, empty-minded, lounging in disconsolate groups at street corners, or aimlessly tramping from one casual ward to another—a prey to all evil influences?"

In Australia and New Zealand, although the conditions under which the masses live are vastly better than in Great Britain, still it is comforting to read the statement of the late Professor Cunningham, the distinguished anthropologist:

"In spite of the marked variations which are seen in the physique of the different classes, anthropologists believe, with good reason, that there is a mean physical standard, which is the inheritance of the people as a whole, and that no matter how far certain sections of the people may deviate from this by deterioration, the tendency of the race as a whole will always be to maintain the inherited mean. In other words, those inferior bodily characteristics which are the result of poverty (and not vice, such as syphilis and alcoholism), and which are therefore acquired during the lifetime of the individual, are not transmissible from one generation to another. To restore, therefore, the classes in which this inferiority exists to the mean standard of national physique, all that is required is to improve the conditions of living, and in one or more generations all the ground that has been lost will be recovered."

The Anzacs.

Together with the sister overseas Dominions of Canada and South Africa, Australia and New Zealand offer a field for studying the effects of what is or ought to be generally a healthy and natural environment on a people, for the most part at least, only one or more generations removed from the parent stock of the British Isles, or, in the case of parts of Canada and South Africa, people from France and Holland. Sufficient data have already been collected with regard to school children to show that the improved environment has had a beneficial effect. Records collected by the late Mr. Symons and myself for the New Zealand Health Department in 1908 and 1909—quite as extensive as to numbers examined as the data available to witnesses at the time of the Inter-Departmental Committee—were especially interesting in showing that, while the average stature at the ages of 11 to 12 of boys in the public schools in England in 1883 was 54.98 inches, that of the cadets of the Auckland Grammar School and King's College was 56 inches at the same age. Contrasting the heights of boys at Marlborough College, England, with those of King's College, Auckland, and Wanganui College, we found that between the ages of 14 and 15 the figures were 61.96 for the famous English public school and 63 for the New Zealand schools. Even more interesting was the fact that a comparison between the boys of the Chapel Street School, Auckland (probably the children of the poorest parents of the largest New Zealand city), with the boys of Remuera (the wealthiest suburb), showed less disparity than the measurements and weights of children in the poorer districts of English cities as compared with the children of the more prosperous parents residing in the suburban centres and the boys of the English public schools. Such a result was only to be expected in a country where there is not the severe handicap of poverty—such a dead weight in some of the congested centres of Great Britain. Moreover, the prowess of the young Australian and New

Zealander in the field of war or of sport need not make him ashamed to meet the stranger either within or without his gates.

Dr. Harvey Sutton, Director of the School of Hygiene and Tropical Medicine, Sydney, late Principal Medical Officer to the Education Department, New South Wales, has shown by statistical tables and graphs that the individual Australian boy or girl is physically superior to the Australian born with one British parent, and the latter to the child with both parents born overseas, judging by height and weight. Whether the comparison is made in the metropolis, town, or country, the more Australian the child, the better the physique. The rural child of whatever "Australianship" has been shown to outstrip the child of the large country town and of the metropolis; between these the difference is much slighter. Although the Australian figures for recruiting examinations have not been properly analysed, where three out of nine were classified A1 in England, about four in every nine were up to the standard in Australia. The C3 class was certainly smaller. The predictions of school hygienists in regard to the findings of from 50 per cent. to 60 per cent. of defectives in schools were thus verified. Dr. Sutton points out that the defects in the years of school life prove to be the basis of the later physical deterioration.

Thus all is not well, even in Australia. Lieut.-General the Honourable Sir Neville Howse, V.C., K.C.B., the distinguished Director-General of Medical Supplies of the Australian Imperial Force, later Australian Minister of Defence and Minister of Health, expressed grave doubts in 1921 whether 50 per cent. of Australia's young men were fit for active service. In a note on the health of the young male Australian, written in reply to a request by myself, as to the percentage of those volunteering for service who were fit, he stated:

"The necessary information is not available to make an authentic statement, but a certain amount of reliable evidence does exist which lends support to my belief:

- (a) Out of 21,000 senior cadets, born in 1905, who were examined, 13.6 per cent. were found unfit for transfer to the Citizen Forces, notwithstanding that many medical officers passed a number whom they recognized would never be fit for service, but who would, in their opinion, be benefited by a few years of physical training;
- (b) Dental examination of Citizen Forces at camps of training shows that a large proportion suffer from dental defects unfitting them for active service."

These facts, together with the appalling number of men who during the war were found, on examination, unfit not only for active service but for strenuous civil employment, are a travesty on public health. Speaking generally and comparatively, however, the good physique of the Australian Imperial Force was conspicuous—as might be expected in a voluntary army from a country blessed with a good climate, good stock, and with a people given to outdoor life and sport—even in our cities which possess a healthful open-air environment, where slum conditions are few, and overcrowding has not existed long enough to cause any deterioration of type.

The analysis of the present population of Australia shows that 97 per cent. are of British origin, and actually 99 per cent. are British subjects. Although Australia and New Zealand have the greatest natural increase of population in the world except Quebec, at the present rate of natural increase—the difference between births and deaths being only 2 per cent. per annum—we would have 12 million people in thirty years, even if we did not get a single immigrant, and at least 50 million in a century, through natural growth, actually as many people as there were in the United States of America at the end of its first century.

Ellsworth Huntington, in *Civilization and Climate*, makes out a convincing claim that climate is the chief factor in the development of a nation. A mean temperature of 64° F., a mean humidity of 80 per cent., and frequent changes of temperature, are the best conditions for physical work. He shows that people who, like the Scotch, live under conditions of relative storminess, become industrious, capable, and successful.

McCollum, on the other hand, in *The Newer Knowledge of Nutrition*, emphasizes diet as the prime factor in the develop-

ment of a people. He ascribes, for instance, the virility and progress of the Canadians in contrast with the sloth, improvidence, and retrogression of the people of the Bahamas—both descended from a common loyalist stock who emigrated from the Northern and Southern States, rather than live under the Stars and Stripes—not so much to the highly stimulating climate of Lower Canada in contrast to the mild and uniform climate of the Bahamas, as to the difference in diet. The Bahamas produce tamarinds, olives, lemons, oranges, limes, citrons, pineapples, pomegranates, figs, bananas, melons, yams, potatoes, cucumbers, sugar, maize, and peas, or fruits and vegetables not unlike those of Queensland and the Northern Rivers district of New South Wales, but their bread and meat are imported. The people of Lower Canada subsist mainly on cereals, peas, beans, potatoes, meat, milk, cream, butter and cheese—a diet similar to that of the people of Australia.

With regard to the Australian national diet, D. T. Sawkins, in a memorandum to the New South Wales Board of Trade, states:

"The average food supplies per head of population remain the most important factor in determining national physique. It is hard to avoid the conclusion that the comparatively superior physique of the Australian soldiers was built up on the liberal average diet of the decades before the war. Then, the average gross supply of meat in New South Wales was 240 lb. to 290 lb., in contrast to the present [1922] consumption of 170 lb. to 200 lb. per man per annum. If this great change in the national diet is to stand for any considerable period, then a great change may be expected in the national physique."

Equally true with the dictum that an army fights on its stomach is it that "the fate of nations depends upon how they are fed," and, as my old teacher, Professor E. J. Hamilton, used to say: "It is better for a man to have a good stomach than a good head."

As far as the health and longevity of the individual is concerned, however, second only in importance to being begotten of a healthy long-lived stock or parentage, is the food of the mother during his pre-natal life, and his diet in infancy. The first fundamental requisite to rear a healthy race is of course that children should be fed, as Nature intended them to be, at the breast of mothers who have themselves been well nourished by a well-balanced vitamin and mineral-containing diet of which fresh milk, salads, and fruit should be the chief constituents.

The superior physique of the pastoral peoples of the past, as of the present, is chiefly the result of a generous milk ration. One cannot visit the Hawkesbury District of New South Wales without commenting on the fine physique of the people, undoubtedly due, as far as the earlier generation is concerned, to a diet of milk, maize, meat, fruit, and other fresh foods, rich in vitamins and mineral constituents, combined with a life in the glorious sunshine and fresh air. It is regrettable that resort to the grocery store rather than to local farm produce, and the feeding of whole maize to the fowls rather than to the children, is not maintaining the fine physique of the Hawkesbury native. The "cornstalk," a New South Wales type with a reputation for lankiness and wiriness, is the immediate successor of the Hawkesbury native. A visit to a Hawkesbury race meeting shows that the older generation consists of tall, brawny, well-made, big men. Mr. J. H. M. Abbott, hailing from the Hawkesbury River Flats, writes:

"The first of the free settlers who commenced the march westwards 'squatted' there, and owned the land by right of occupation. If they could obtain them, they took wives unto themselves, and reared up families. And the families subsisted principally on pumpkin and ground maize, and wore no boots in their childhood, and led a free, wild, untrammelled sort of life. So they grew into tall, clean-limbed, deep-chested men, and sturdy, comely women, and spread north and west and south over the land—and their offspring were the fathers and mothers of the Cornstalks of to-day."

The Analogy of the Merino.

The people of Australia to-day owe a great debt to Captain John MacArthur of Camden Park, who imported the first merino sheep in 1796. The pioneers were told that the climate was unsuitable and that the merino wool would deteriorate as the sheep were taken nearer the Equator. At the beginning of last century the heaviest fleece grown weighed 3½ lb., whereas actually a fleece of 35 lb. was recently

cut from a stud sheep. The merino sheep to-day, however, is very different from the original stock. Some of the direct descendants are still to be seen at Camden Park. Various strains have been crossed with the original blood—Saxon, Negretti Infantado, and, above all, the French Rambouillet. Thousands of the last strain were brought to Australia in the first half of last century. So readily did they adapt themselves to their new environment that it was speedily recognized that this type was ideal for producing fine wool under our special conditions of sunshine, climate, and green pasturage. Since then the great sheep breeders have evolved a truly national Australian merino, quite distinct from all the older strains of Europe—greater in every respect, greater, indeed, than any other variety of the breed in any other country in the world.

Two young Englishmen of French extraction—the Pepins—must be credited with the actual development of the merino as it is known to-day. In 1864 they began to build up a stud flock at Wanganella, by purchasing a ram, now famous under the name of "Emperor," a big well-proportioned, plain-bodied, outstanding animal with strong head and horns. So well did he nick with the Wanganella ewes that he became leader of the stud. The progeny of the other imported animals was gradually eliminated, until the whole flock became inbred to him.

Various sheep breeders since the day of the Pepins, like the Falkiners, the Austins, the Mills, the Millars, the McKaugheys, and others, by careful selection, judicious mating, and heavy culling, have improved each generation until now our large flocks are bred to type with variations according to climate and soil: Stamina above everything else is the quality demanded. Pampering is anathema to the big breeders. Science has come to the assistance of the stud masters. For instance, Professor Brailsford Robertson says that "if any one of the amino-acids that build the tissues of the sheep is lacking in the food supply of the animal, the whole of the rest of the food becomes valueless as far as building up the tissue is concerned."

The industry of sheep breeding is now being conducted on a more scientific basis. Who can say whether sheep at the end of the present century will not be carrying twice the fleece they carry to-day, when they average 8 lb. per sheep, instead of the average of 3 lb. half a century ago? If we could apply the same system to the human element in Australia—that is to say, cull by the elimination of the unfit, especially the mentally unfit, and select the best type of immigrant, physically, mentally, and morally—we would still further improve the present breed. Certainly the fine agricultural country type of Englishman, Irishman, and Scotsman, predominating before 1820, which may have deteriorated to a certain extent by the industrialism of the later years of last century, is being reproduced in Australia, South Africa, Canada, and New Zealand by a type which bids fair to be even better in physique.

The Growth of Population in Australia and Canada.

The causes of emigration are stated to be adversity, economic distress, and the attraction of the prospect of wealth and improved conditions, apart from the earlier causes, such as political or religious oppression. The great migration between 1841 and 1847 when 2½ million people left Ireland, mainly for the United States, was caused by the years of potato famine. As far as Australia, including Tasmania, is concerned, although in the early days a certain number of people are said to have left their country for their country's good, undoubtedly many of them made good in the new country, as was the case also with Virginia and Maryland in America, where those transported became in many cases industrious and exemplary citizens.

Australia to-day is peopled by immigrants or the descendants of immigrants. There is no one living in Australia (other than the aborigines) who is a descendant of the fourth generation of Australians. We have actually an area of 3,000,000 square miles, nearly equal to that of the United States of America, four-fifths of that of Canada, over one-fifth of the area of the whole of the British Empire—that is, more than three-quarters of the whole area of Europe, and actually twenty-five times as large as Great Britain and Ireland. We have a population of only 6½ million, while

Great Britain and Ireland have a population of roughly 50 million. As a matter of fact, the Dominions overseas, embracing Canada, Australia, New Zealand, and South Africa, have only about 18 million whites. It is to be remembered, however, that five-thirteenths of Australia is within the Tropics.

In spite of the distance from the Motherland, the population of Australia since Governor Phillip landed 141 years ago has shown phenomenal growth, especially in the last hundred years. During this period it has grown to the same size as that reached in Canada in three centuries. The most conservative estimate of the population that Australia could accommodate has been stated to be 40 million. The Right Honourable William Morris Hughes estimated that Australia could maintain a population of 100 million, while some have put the figure as high as 200 million.

While Canada occupies one-sixteenth of the world's area, it has only one two-hundredth of its population. The first settlement in Canada was at Port Royal or Annapolis, Nova Scotia, in 1605. By 1763 there were 90,000 people, in 1800 a quarter of a million, whereas now there are 10 million. In Canada to-day 78 per cent. of the people are native born, while only 14 per cent. are British born.

As showing how history repeats itself, the Napoleonic wars, just like the last war, created tastes, habits, and conditions incompatible with the peaceful activities of the new era that followed. The total population of the world in 1800 was 700 million, whereas it is 1,900 million to-day. During that time the population of England and Wales increased from 12 million to 38 million.

Immigration Problems.

The problem of the distribution of population throughout the British Empire is not so simple as it might appear. The mere transfer of the surplus population from one part to another which is sparsely populated, so far from solving the problem of unemployment, may tend to aggravate it. In Australia it is extremely difficult to absorb some immigrants, especially those whose ways and ideas have been set before they arrive; youths under 20 years, provided they are of reasonably good stock, can be made into good Australians. Older people find the conditions so different from their previous environment that they have difficulty in accommodating themselves.

One method of settling immigrants has been that of the group settlements in the south-west of Western Australia, where immigrants receive 160 acres of land and 10s. a day for two and a half years. These settlements are well described in *Amateur Settlers* by Lord Apsley, who found that one-third of the settlers within an area abandon their holdings. He ascribed seven failures out of ten to ill health. Mr. W. Banks Amery, British Dominions Office Representative in Australia, found that 32.5 per cent. of British settlers either abandoned their holdings or were dismissed. Of 1,105 Australian settlers, 45 per cent. left their holdings or were dismissed. Thus actually among "stickers" 68 per cent. were recent British arrivals and 59 per cent. native-born Australians. A Royal Commission found that one of the difficulties was the dislike of the women for dairying, and the preference of numbers of men to work for wages rather than accept the responsibility of fending for themselves. It was also found—as in the case of the returned soldiers from the war for whom many millions were spent in settlement on the land in Australia—that poor and unsuitable land had sometimes been selected. The opinion now prevailing is that these settlers should be placed only on first-class land, that payment should be by piece-work, and that they should be drafted into preliminary training camps. Altogether there have been 135 groups of settlers, which have provided a population of 9,580 at an expenditure of 3½ million pounds.

The Big Brother Movement, by which boys are brought out to Australia and placed under the protection of citizens, as the boys brought out under the Dreadnought Fund, has been satisfactory.

The general impression of the people with regard to the immigrants that arrive is that they are less than "fair average quality." Most of the States welcome land workers and

domestics, but do not encourage other immigrants except nominees—that is, those who have friends already in the country prepared to assist them.

Where the medical profession might do great work for the Empire is by encouraging a suitable type of youth to emigrate, one with a good family history, of good physique, ambitious, and above all, with a will to work. Dr. Stewart Mackintosh's observations¹ suggest that readiness of adaptation to change argues an ancestry derived from many and varied stocks, such as are usually bred together by urban influences. One result of the rapid transition from an agricultural to an industrial nation has been the disintegration of rural communities, their aggregation in new centres, and the extraction by a severe process of natural selection of types that could adapt themselves to industrial exigencies, and climatic and other conditions. The New World has drawn some of its most hard-working labourers from limbos of despair, men who develop a surprising physique and energy as soon as they can enter environmental conditions at all conformable to their physical needs. Professor Karl Pearson emphasizes "that it is essentially the man who makes his environment, and not the environment which makes the man." We have chosen to make ourselves an industrial nation and allow our agriculture to decay; we have offered industrial work to those who made their living on the soil, luring them into the cities and into environments differing in many respects from those in which they were brought up.

A few sons of medical men have come out to Australia since the war, received some training at the Hawkesbury Agricultural College, and, in some cases, are doing well on the land to-day. If anyone cared to organize a batch of sons of medical men, it would not be difficult to arrange for them to be met on arrival and have them distributed to sheep and cattle station owners or farmers in Australia and New Zealand on similar lines to the Big Brotherhood Movement. Certainly if they came out with the intention of settling and marrying, the Australian girls, who are even a finer type than the men, would give them a hearty welcome.

REFERENCE.

¹Mackintosh, J. S.: The Effects of Intra-Migration on National Health, *British Medical Journal*, 1912, ii, p. 367.

RADIUM TREATMENT OF EPITHELIOMATA OF THE SKIN.*

BY

ROY WARD, M.B., B.S.,
THE RADIUM INSTITUTE, LONDON.

In this opening paper I propose to deal mainly with the subject of basal-celled epithelioma or rodent ulcer, and to touch briefly on the question of squamous epithelioma, believing, as I do, that this disease has a surgical rather than a radiological aspect.

RODENT ULCER.

Radium is undoubtedly the best treatment in most cases of rodent ulcer; it is in this condition that the most striking and permanent results of radium therapy are obtained. Excision is equally effective in some cases, but, owing to the difficulty in excising completely even a small growth from the situations most commonly affected, radium is to be preferred. Cosmetically it gives results which cannot be obtained by any other method, the resulting scar being supple and the skin almost normal in appearance. X rays will cure a number of these lesions, but the treatment is tedious, and the action of the rays cannot be limited with such great accuracy. Recurrences are more frequent, and permanent healing is not so certain.

Clinical Varieties.—A rodent ulcer usually begins as a smooth, waxy, semitranslucent pearly nodule, which grows very slowly, often assuming the form of a rosette of nodules with a raised margin and depressed centre. A growth such as this belongs to the early hypertrophic variety, and frequently shows cystic degeneration. More nodules may

appear and coalesce, thus forming a larger hypertrophic growth, or, owing to some slight trauma, the growth may break down, with the formation of a brownish scab. Gradually there develops a superficial ulcer with a gelatinous base, and the raised pearly margin which is so characteristic of the disease. The ulcerated area may heal over in one part, leaving a smooth atrophic scarred area, while in another part the disease continues to advance. This is known as the superficial flattened cicatrizing type, which has a characteristic braided margin. Both the hypertrophic and superficial ulcerative types may reach a considerable size, but generally remain limited to the surface. The disease may, however, extend deeply to reach the underlying tissues, penetrating muscle and bone, thus giving rise to deep irregular cavities. Such lesions of the deeply ulcerative variety seldom have the typical rolled edge; the margins are usually indurated and sharply cut, with a well-marked inflammatory zone surrounding the ulcer. Although there may be extensive local destruction of tissue, these growths never give rise to metastases unless they assume the clinical and histological appearances of a squamous-celled epithelioma. It is very important to recognize this transitional type. The relative frequency of the various types can be shown from an analysis of 1,773 consecutive cases treated at the Radium Institute, London:

Type of lesion.	Relative frequency.
Hypertrophic (small and large)	51 per cent.
Superficial:	
Ulcerative	30 "
Cicatrizing	3 "
Deeply ulcerated	16 "

The following shows the results for each group:

Type.	Apparently cured.
Hypertrophic	91 per cent.
Flat	77 "
Deeply ulcerated	38 "
All cases	77 "

Many of the last group were such that healing was out of the question, owing to extensive involvement of bone and cartilage.

Sites.—Rodent ulcers are most common on the face. The most frequent situations are the nose and cheek, as shown in the following table:

Nose	468	Chin	36
Cheek	377	Mastoid	22
Inner canthus	193	Trunk	21
Lower eyelid	151	Neck	20
Temple	141	Upper eyelid	19
Forehead	98	Lower lip	17
Ear	84	Eyebrow	12
Outer canthus	56	Scalp	10
Upper lip	47	Hand	4
Preauricular	37		

Etiology.

The disease is more common in men than in women—953 patients were males and 820 females. The youngest patient in the series was a girl who was first seen at the Radium Institute at the age of 19. She was sent on account of a deeply ulcerative lesion which affected the inner canthus, and had started as a nodule at the age of 12. It will be seen that the disease is most common between the ages of 60 and 70.

Age.	Frequency.
10-20	1
20-30	11
30-40	63
40-50	218
50-60	394
60-70	484
70-80	334
80-90	82
90-100	2

Multiple lesions are more common than is usually supposed. There were 94 cases in this series; 55 were in men and 39 in women. In one case a woman, aged 28, had as many as nineteen different lesions treated with radium.

Previous Treatment.—A number of patients had been operated on once or several times, and many had received treatment by x rays or radium at other hospitals. The disease had either recurred or had become resistant to

* A paper read in opening a discussion in the Section of Radiology at the Annual Meeting of the British Medical Association, Winnipeg, 1930.