different things, it may be that similar factors are at work in dysenteric infections, and may explain the absence of lasting agglutination in some acute cases and its persistence in those people who can give no history of dysentery.

CONCLUSIONS.

1. Agglutinins to *B. dysenteriae* were present in 6.29 per cent. of 524 serums taken from individuals only one of whom gave a history of dysentery, thirty years previously.

2. The series consisted of an equal number of males and females. The males showed 6.1 per cent. positive and the females 6.48 per cent. The highest titres were found in specimens from males.

3. The males reacted most frequently to Sonne's bacillus, but the highest titres were to Types V, W, and Z of the Flexner group. The females reacted most frequently to Type V Flexner.

4. The presence of these "normal agglutinins" may interfere with serological diagnosis, as they sometimes come within the range of the standards laid down as indicative of active infection; besides, in active infections agglutinins may not always reach these suggested standards.

5. There is sufficient evidence to suggest that these so-called normal agglutinins are the outcome of subclinical or mild infection with homologous or associated micro-organisms, and that the term "normal" when applied to this reaction is a misnomer, and should be avoided.

My thanks are due to Dr. F. M. Rowland of Lichfield for revising the manuscript.

REFERENCES.

- ¹ Report on an Investigation of Dysentery and Diarrhoea in Poona, by Major Manifold, D.S.O., R.A.M.C., and Assistant Surgeon A. J. DeMonte, Indian Medical Department, R.A.M.C. Journal, July, 1928.
- ² Medical Research Committee Special Report Series No. 7.
- ³ Medical Research Council Special Report Series No. 51.

A CASE OF CONJOINED TWINS.

BY

M. WITKIN, M.B., CH.B., L.R.C.P. AND S.ED.,
DURBAN, SOUTH AFRICA.

RARE as are conjoined twins, still rarer are cases in which one twin is normal and the other has a parasitic foetus attached. The case recorded below seems, therefore, worthy of publication.

A multipara, aged 35, living near Durban, Natal, gave birth to male twins on November 12th, 1928. The labour was, on the whole, easy. The first child, a head presentation, was normal in every respect, and weighed 7 lb. The second child, a breech presentation, had a parasitic foetus attached to it. I was called half an hour after the delivery, as the midwife, finding an unusual bulging in the umbilical region of the second twin, was afraid to cut the cord. When I arrived I found the placenta in the bed, and after cutting the cord I noticed a hernia-like protrusion of the bowel in the umbilical area.

Family History.

The father is 38 years of age, a factory employee of poor physique, below the average height and weight, and of intemperate habits. The mother is a healthy woman, and is of average height, weight, and physique. Three other children (two boys and a girl), aged respectively 7, 5, and 3, are normal in every respect. There is no history of miscarriages or of difficult labour.

The Autosite.

The autosite, a small baby weighing 5 lb. 8 oz., was poorly covered, had deficient muscle tone, loose skin, a Hippocratic facies, and a feeble cry. It was alert, cyanotic, especially about the feet, and had normal powers of suction and deglutition. The head was normal, except that the cranial bones were unusually soft. The lungs, the upper and lower extremities, the abdomen, and the anal orifice were normally developed, and hearing, sight, and intelligence appeared to be normal. On the other hand, a bruit was heard in the heart, due, probably, to a patent foramen ovale, and the genito-urinary system was abnormal. The left testis was undescended, and metrorrhagia occurred at independent times from the penis of the autosite and from that of the parasite.

The Parasite.

The parasite was attached to the autosite by a pedicle the circumference of which was about that of half a crown. The pedicle was partly bony and partly muscular. The upper bony part was attached, slightly to the right, to the lower end of the sternum of the autosite; the lower part of the pedicle was muscular, and joined to the upper part of the abdominal wall.

The chest of the parasite was undeveloped, and contained no heart or lungs. Its upper extremities were rudimentary; the

right arm was longer than the left, and each hand had one rudimentary metacarpal, finger, and nail. The lower extremities were well developed and equal in size, but they were flexed to a right angle, and could not be straightened or bent any further. The ankles were stiff; each foot had five well-developed toes. There were no testes in the scrotum, but the penis was well formed, and the parasite passed water over the autosite. There was no umbilicus and no anal orifice, though a distinct dimple marked the place of the latter. The parasite displayed no movement, but when its skin was pinched or pricked with a needle the autosite winced and uttered a feeble cry.

Operation for Umbilical Hernia.

On November 13th, when the child was one day old, I opened the abdomen, removed the hernial sac, returned the intestines to the abdomen, and closed the wound.

The child made an uninterrupted recovery, the cyanosis disappearing almost at once and its general condition improving. Waterproof material was used to protect the wound from the parasite's urine. Healing took place by first intention, leaving a clean scar.

I considered that the child's condition did not justify an attempt to remove the parasite, and decided to wait a little before undertaking this operation. Later, however, when he was 10 weeks old, I found him looking extremely ill, and was able to induce the father to allow me to remove the parasite at once.

Removal of Parasite.

The operation was performed on February 6th under light chloroform anaesthesia. I made an incision round the pedicle, and on opening the abdomen I found that there was free communication between the abdominal cavities and between the bowels themselves. The peritoneum of the autosite was continuous with that of the parasite, but the latter's bowel had an independent mesentery attached to its posterior abdominal wall. The intestines of the parasite were very thin and narrow; they originated from the autosite's small intestine and ended in its lower bowel. The parasite had one small right kidney, but no liver or spleen. Large blood vessels and nerves passed from the autosite to the parasite. I ligated and cut the vessels passing into the parasite, and then decided to cut through the bowel of the parasite, taking care not to endanger the intestines of the host. Keeping as close to the parasite as possible, I clamped the bowel, removed it as one would an appendix, and returned the host's intestines into his abdomen. I closed the abdomen in the usual manner and brought the edges of the skin together, by interrupted silk-worm-gut sutures. The operation lasted fifteen minutes, and except for a little sloughing of the wound the child made an uneventful recovery. The parasite weighed 8 ounces.

The child was breast-fed at three-hourly intervals, and twelve hours after the operation he had a normal stool. The fact that thereafter the stools were of normal colour and consistency assured me that his intestines had not been damaged at the operation. He began to improve steadily and gained in weight, and a month after the operation he was still alive and doing well.

A, The conjoined twins at the age of 2 months. B, The parasite immediately after the operation. C, The autosite at the age of 3½ months.

PROGNOSIS.

The prognosis is grave. The child is undersized, undeveloped, and has a malformed heart. His environment is bad; the parents live under insanitary conditions, and the child does not receive the care essential for his survival.

His case differs from that of Laloo described by Sir John Bland-Sutton, and from that of Karramat Hussam described by D. D. Kapur; both Laloo and Hussam were well developed and healthy in every other respect. My operation differs from that performed by Kapur, who did not cut through the intestines of the parasite, but transferred them complete into the abdomen of the host.

I wish to record my thanks to Dr. H. Mundy for his valuable help and advice.

LITERATURE.

Sir John Bland-Sutton: *Trans. Path. Soc. London*, 1888, xxxix, 427; *Lancet*, 1928, i, 24; *British Medical Journal*, January 5th, 1929, p. 1; *Tumours*, p. 507.

W. Roger Williams: *Medical Journal and Record*, New York, 1927, p. 6.

D. D. Kapur: *Indian Medical Gazette*, 1927, lxii, 1.

Gould and Pyle: *Anomalies and Curiosities of Medicine*, 1897, p. 192.

Memoranda:

MEDICAL, SURGICAL, OBSTETRICAL.

SPONTANEOUS RUPTURE OF THE SPLEEN.

HAVING read the article by Mr. D. H. Patey in the *British Medical Journal* of May 18th (p. 898), I am submitting the following case, as, in view of its similarity, it may be of interest in conjunction with the other cases.

The patient, a cashier aged 50, was admitted to hospital under the care of Mr. Girling Ball with a history of having, six hours before admission, while standing at his work, been seized with sudden acute pain in the upper abdomen. He vomited once (apparently his last meal); the pain was severe enough to double him up, and he collapsed; the pain remained of about the same severity until admission. For the last ten years he had been troubled by flatulence after food, but had had no clear symptoms suggestive of a gastric ulcer. Many years ago he fractured some ribs on the left side, which was followed by pleurisy.

On admission he was very ill, pale, and sweating. The temperature was 96°, pulse 82, respirations 20. The tongue was dry and furred. The chest showed some bronchitis; the heart was normal. The abdomen was not distended and did not move on respiration; there was marked rigidity and tenderness in the upper half of the abdomen. The liver dullness was not diminished. Free fluid was present in the peritoneal cavity. The urine was normal.

A diagnosis of perforated gastric ulcer was made.

Operation.—Under nitrous oxide and oxygen with ether (endotracheal method) a right paramedian incision was made. On opening the abdomen the peritoneal cavity was found full of blood. This was mopped out, and the haemorrhage found to be coming from an irregular tear, 2 inches long, on the anterior surface of the spleen. The spleen itself was markedly adherent to the abdominal parietes posteriorly and at the upper pole. The abdominal incision was enlarged by a transverse incision across the left rectus muscle and splenectomy performed; all haemorrhage was easily controlled, and the abdominal wound was closed, a drainage tube being inserted down to the region of the spleen. At the conclusion of the operation the patient was transfused with 350 c.cm. of blood.

The patient progressed satisfactorily as regards the abdominal condition, but on the seventh day he developed signs of bronchopneumonia at both bases and a left-sided pleural effusion. He died four days later. There was no post-mortem examination.

Macroscopically the spleen was not diseased, but showed old adhesions posteriorly and at its upper pole. Microscopically there was a slight degree of fibrosis present, and a hyaline degeneration of the blood vessels. The sinusoids were devoid of blood vessels; there was some pigmentation present.

I am indebted to Mr. Girling Ball for permission to publish the case.

W. E. UNDERWOOD, M.B., B.Chir.,
Late House-Surgeon, St. Bartholomew's Hospital.

OBSTRUCTED LABOUR DUE TO OVARIAN CYST.

THE following case presented interesting points in diagnosis and treatment.

A married woman, aged 31, who had given birth five years previously to a child after a normal confinement, was admitted to the North Lonsdale Hospital with a report that on examination it had been found impossible to identify the cervix, or to define any presenting part of the foetus, though labour had been in progress for five hours.

General inspection of the abdomen suggested the presence of a relatively large uterus, and by abdominal palpation the vertex

was found to be occupying the lower pole of the uterus. The head was not engaged, and was freely movable above the pelvic brim. There were no signs of hydramnios or of a twin pregnancy. The uterine pains were feeble. The foetal heart was normal in rate. On vaginal examination the fingers entered a cul-de-sac with no outlet superiorly, and no cervix was palpable. The posterior vaginal wall was bulging forwards, and it was evident that a mass occupied the pouch of Douglas. Under anaesthesia a tiny chink was felt high up anteriorly, through which the finger could be passed on gentle pressure; the cervix was felt high up in this position, and was dilated to the size of a florin. The mass in the pelvis was firmly impacted, and it was not possible to effect any upward reduction of the tumour. The diagnosis of an ovarian cyst obstructing labour was made; it was ascertained that the child was alive.

The abdomen was opened and the uterus brought through the wound; it was as yet impossible to define the ovarian tumour or to reach its connexions. Caesarean section was therefore performed, and a live female child delivered. The tumour was now apparent; it was of large size and of hour-glass shape, the larger part being abdominal in position, and the lower portion firmly impacted in the pelvis. The pedicle was defined and ligatured and the tumour was removed. The tumour was cystic in nature, and was large enough to fill almost entirely an ordinary pail. The child weighed 8 lb., and the placenta was heavy, hence the total weight removed from the abdomen was very considerable. The puerperium was uneventful, and the patient and her child were ready to be discharged from hospital in three weeks.

I wish to thank Dr. Allan Rutherford for his co-operation in the management of this case.

HERBERT F. BLACKLEE, M.B., F.R.F.P.S.,
Honorary Assistant Surgeon, North Lonsdale Hospital,
Barrow-in-Furness.

Reports of Societies.

CANCER OF THE LUNGS AND MEDIASTINUM.

At a meeting of the Manchester Medical Society, held at the University, with the president, Dr. R. W. MARSDEN, in the chair, a discussion took place on malignant disease of the lungs and mediastinum. The opening paper, from the clinical point of view, was given by Dr. A. HILLYARD HOLMES; Professor J. SHAW DUNN discussed the pathological aspect, and Dr. E. W. TWINING the radiological side of the subject.

Clinical Aspects.

Dr. HILLYARD HOLMES said that primary malignant disease of the lungs and mediastinum was not rare in that part of England; in the last three years he had collected notes of twenty-five cases in which the possibility of the presence of this condition had to be seriously considered. Of these twenty-five cases, eighteen were finally judged to be malignant, and of these eighteen cases seven were submitted to necropsy and to histological verification of the diagnosis. The diagnosis of malignancy was of vital importance in regard to prognosis, for no form of treatment was known which held out any hope of cure. Since the problem of diagnosis was best illustrated by records of actual cases the speaker cited several in which the diagnosis was established by post-mortem examination.

Case 1.—A man, aged 44, ceased to work in April, 1926, because he had strained his back. Pain in the neck was so severe that his doctor suspected cervical caries, and had an x-ray examination made, with a negative result. He lost weight rapidly. Four months later he attended hospital as an out-patient; cough and hoarseness had then existed for only one month. He was thin and pale. His neck movements were stiff; there were signs in both lungs suggestive of chronic phthisis, especially in the left lower lobe. X-ray examination showed appearances consistent with old-standing tuberculous disease. The Wassermann reaction was negative. His doctor noticed some small, hard lymph glands over the inner end of the left clavicle, and sent him into hospital as an urgent case two weeks later. Within twenty-four hours the man had a profuse haemoptysis and died. Post-mortem examination showed a large carcinoma in the left lung, at its root, with a cavity 5 cm. in diameter in the lower lobe, full of blood clot; the stomach contained 1½ pints of blood; there were secondary malignant deposits in the right suprarenal gland, left kidney, left cerebral hemisphere, and left supraclavicular glands. Histological examination showed all these growths to be of epithelial nature, with a tendency to alveolar arrangement. The primary growth was a carcinoma of the bronchus. There was no evidence, macroscopic or microscopic, of tuberculosis. The key to the solution of this diagnostic problem was the enlargement of the supraclavicular glands and the rapidity of the man's decline.

Case 2.—A man, aged 64, was in hospital in January, 1925, for left pleural effusion and slight haemoptysis. The effusion was aspirated and artificial pneumothorax induced. Examination of the fluid revealed red and white blood corpuscles and endothelial cells; no organisms were found; cultures were sterile. The sputum