Drainage consists in exposure of the cyst, evacuation of the contents, removal of the endocyst, and fixation of the tube. Where possible in these cases I usually suture the parietal peritoneum around the proposed site of drainage as a protection against leakage, but in large or multiple cysts this is not feasible. Preliminary sterilization of the contents by injection of formalin is practised in all methods. The duration of post-operative drainage varies with the size and nature of the cyst. The wound is generally healed in three weeks; in some cases, however, discharge may continue for many weeks, and is often bile-stained; occasionally a troublesome sinus persists, which may require radical treatment.

This is a simple and satisfactory method of dealing with the majority of cysts.

Drainage with Partial Obliteration.—The essential difference between this method and that of simple drainage is that an attempt is made to diminish the size of the ectocyst by approximating its walls. In the three cases thus treated this has been effected by means of superimposed layers of continuous catgut sutures after excision of as much of the membrane as possible. The cavity has not been completely closed, a small rubber drain being left in its upper portion for three days. For large cysts with soft walls, and where the endocyst has been removed *in toto*, I consider that this is a sound method of treatment.

Complete Removal (Exsection).—Cysts of moderate size, especially those near the free border of the liver, may be completely removed by a wedge-shaped exsection which includes a portion of the liver substance. Haemorrhage, as a rule, is not serious, and can be controlled by finger pressure, clamps, or gauze. The resulting wound in the liver is closed by a series of deep mattress sutures and by a superficial running suture. The results of this operation are extremely satisfactory in suitable cases. No drainage is required, and patients make a rapid and uneventful recovery.

Note.—The adventitia should be separated from the liver as far as possible by blunt dissection or gauze, thus reducing the actual incision to the smallest possible dimensions.

TABLE II.

	Cured.	Relieved.	Died.	Total.
Drainage	10	4	4	18
Obliteration	2	1	—	3
Exsection	10	—	—	10
Total	22	5	4	31

From a study of the above table the results of treatment may be easily seen. All cases treated are included. Cases are reported as cured where the wound is soundly healed and where there is no return of symptoms, and relieved where, although the general condition of the patient is good, a sinus persists. These latter have all been large calcareous cysts with suppuration of contents. Of the four deaths, three occurred some months after operation among patients with enormous cysts involving practically the whole liver, and were due to hepatic insufficiency. The fourth died of post-operative shock.

Extrahepatic Cysts.

These, which numbered 9, were situated as follows: rectus femoris 1, ischio-rectal fossa 1, lung 3, omentum 3, uterus 1. The cysts in the rectus and ischio-rectal fossa presented the appearance of chronic abscesses, and were recognized only at operation, which consisted of evacuation, swabbing with formalin (2 per cent.), and packing with bipp.

The uterine case was sent in with severe haemorrhage. This patient was aged 55 and a nullipara. On examination, the uterus was found to be enlarged and soft; the os was patulous and admitted two fingers. The vagina contained several small cysts and others were expressed from the uterus. The condition of the patient was very poor and precluded hysterectomy. The uterus was evacu-

ated with a flushing curette, and a number of small hydatids and a large piece of endocyst removed. Haemorrhage was free, and the cavity was packed with flanne gauze. The patient improved and left hospital, having refused further treatment. She died one month later from what was stated to be a severe haemorrhage.

Of the three lung cases, one patient refused operation; the other two were treated by resection of ribs, suture of pleura, drainage, and removal of endocyst. The operations were done under local anaesthesia, and the wounds healed rapidly. Two of the omental cysts were lying in the pelvis and bound down by adhesions. The third was situated in the epigastrium and was freely movable. They were excised intact together with a portion of omentum.

TABLE III.

Site.	Cured.	Unchanged.	Died.	Total.
Rectus femoris	1	—	—	1
Ischio-rectal fossa	1	—	—	1
Uterus	—	—	1	1
Omentum	3	—	—	3
Lung	2	1	—	3
Total	7	1	1	9

DIAGNOSIS.

In a hydatid country the possibility of a cyst is always in the surgeon's mind, and in the case of the liver there is seldom difficulty in making a correct diagnosis. In other sites the difficulties are greater and the symptoms and signs less definite. Diagnosis is occasionally rendered easy by the presence of portions of hydatid material in the faeces, urine, or sputum. X-ray examinations and laboratory tests are of considerable help in doubtful cases. The three lung cases referred to were all well demonstrated by radiography; all gave a positive Weinberg reaction (complement fixation); two a positive endermic test (Cassoni). Eosinophilia occurs in about 50 per cent. of cases, but the blood film is not of great value, especially when chronic malaria is a concomitant condition.

I have to thank Dr. M. Gosden, bacteriologist, for undertaking the laboratory investigations, and Dr. E. J. Blackaby, district medical officer, for permission to include Case No. 18.

FOREIGN BODIES IN THE UPPER CONJUNCTIVAL FORNIX.

BY

THOMAS COLLEY, F.R.C.S.Ed.,

SURGEON TO THE WEYMOUTH AND DORSET COUNTY ROYAL EYE INFIRMARY.

The following two cases were recalled to my mind by a report of a somewhat similar case published in the March number of the *British Journal of Ophthalmology* by Mr. Sawhney. These cases are of particular interest, since it is quite possible for the presence of such foreign bodies to be overlooked, when abscess formation would undoubtedly follow. Once a case of this type has been recognized there should be no difficulty in making a diagnosis or suspecting the condition in subsequent cases.

First Case.—In November, 1929, a man attended my out-patients' clinic with a muco-purulent discharge from the right eye; he stated that while threshing a week previously something had entered this eye. The appearance was typical of muco-purulent conjunctivitis, the lashes being stuck together with inspissated discharge; the upper lid, however, was somewhat more thickened than one would have expected. On separating the lids the bulbar conjunctiva and that of the lower fornix were relatively clear except for slight injection; eversion of the upper lid showed the conjunctiva to be somewhat injected, but not to the extent that the external appearance suggested. The discharge appeared to come from the inner end of the upper fornix, but even when this area was prolapsed, by pressure on the globe with the upper lid everted, no foreign body could be detected. This part of the conjunctiva was intensely injected and somewhat thickened. The cornea and the eye itself were normal throughout.

Local treatment, consisting of frequent lavage with boracic lotion, drops of colloidal silver, and yellow ointment, was prescribed, and considerable improvement ensued; the oedema of the upper lid gradually disappeared on the outer side, but persisted over the inner end. The conjunctiva cleared rapidly, but considerable discharge was present, which still came from the inner end of the upper fornix. Further examination of this region now showed the presence of a small granulation.

With the conjunctiva anaesthetized by cocaine, and the upper lid doubly everted, this granulation was gently scraped with a sharp spoon, and a yellowish-white point was exposed. This was seized with a pair of forceps, and a piece of macerated straw, almost an inch in length, was withdrawn from the orbital tissue. The condition cleared rapidly in the next three days, and a week afterwards the patient was discharged.

Second Case.—In July, 1930, a man presented himself at my hospital clinic for treatment, giving a history that, while threshing a week previously, a piece of barley straw had blown into his right eye. The upper lid was inflamed and considerably thickened, the man being unable to raise it. The site of inflammation appeared to be in the region of the lacrimal gland, where a definite swelling was present; there was a slight muco-purulent discharge, but no fluctuation of the swelling was obtained by palpation. On raising the lid the appearance of the globe was normal; there was, however, slight limitation of movement upwards and outwards. The cornea was clear, and the conjunctiva mildly injected. The lid was so thickened by oedema—especially the outer part—that it was impossible to evert it.

A hand retractor was employed, and the lid was raised, when the discharge was found to come from the outer part of the upper fornix, where the conjunctiva was raised in folds by oedema. This region was washed out, and a bead of pus was noticed coming from between two folds which, when separated, revealed the end of a piece of barley straw. This was removed with forceps; it appeared to have been embedded in the palpebral portion of the lacrimal gland. The condition resolved satisfactorily in forty-eight hours, and the man was discharged in six days.

It is quite common, especially in rural areas, to see cases where pieces of straw or like material have been retained in the upper fornix for considerable periods. The upper fornix is a cul-de-sac, which is formed by the reflection of the conjunctiva from the upper lid on to the globe. The sensitiveness of the conjunctiva is relatively low and, since the anatomical structure of this area is such that foreign bodies are held well away from the sensitive cornea, their presence is often undetected by patients until a local reaction has been set up.

Foreign bodies in this position produce a clinical picture which is fairly typical. Pain is usually absent, and the discharge, at first watery, quickly becomes muco-purulent. The upper lid becomes oedematous and the superior palpebral fold puffy. The lashes stick together with discharge, but when the lids are separated the conjunctiva appears fairly healthy. The upper fornix, however, is the site of an intense reaction, the intensity being dependent on the length of time during which the foreign body has been retained. In some cases ulceration may take place, and the foreign body be found embedded in granulation tissue. It is also feasible to think that such foreign substances as straw, which are relatively sharp, and which at first lie horizontally along the fornix, may later cause ulceration, and the action of the lids may actually drive them into the tissues of the orbit.

In the first case recorded I believe that the straw actually penetrated the conjunctiva of the upper fornix on entry, since there was no ulceration and only one small granulation locally at the point of entry, but in the second case, where there was much local inflammation and reaction, the straw might have been thrust into the orbital tissues through a softened and ulcerated conjunctiva. In all cases where a muco-purulent discharge is present on one side only, and where lacrimal obstruction and dacryocystitis are absent, the necessity of complete eversion of the upper lid and an exploration of the upper fornix cannot be over-emphasized.

Had these two cases been seen soon after the entry of the foreign bodies it is more than possible that their presence would have been detected, either in the fornix itself or penetrating through a conjunctiva which would at that time have been practically normal. Advice was not sought, however, until some seven days after the entry, when the clinical picture was much more like that of a lid abscess.

Memoranda:

MEDICAL, SURGICAL, OBSTETRICAL.

NICOTINE POISONING BY ABSORPTION THROUGH THE SKIN.

THE causal circumstances in the two cases here reported appear to be unusual.

A. B., aged 19, an under-gardener, was seen at 10.30 a.m. on December 21st, 1929, suffering from prostration, abdominal pain, and violent spasmodic vomiting, sickness having come on about 6 o'clock on the previous evening, continuing at short intervals all night. F. K., aged 34, head gardener, was seen in a similar condition at 3.30 p.m. on December 21st, vomiting in his case having started at 9 p.m. on the 20th, continuing at intervals of about half an hour. The vomitus appeared to be chiefly bile. In both cases the bowels were confined; temperature subnormal; pulse thin and accelerated (slightly irregular in the case of A. B.); tongue dry and yellow-coated; skin on the extremities pale and cold. There was marked muscular relaxation in the lower extremities. The slightest attempt at movement appeared to induce further retching and vomiting. A. B. had cycled home, a distance of two miles, at 5 p.m., and felt ill immediately on taking a cup of tea. F. K., who lived at the gardens, had tea at 6 p.m., and first felt sick two and a half hours later.

Both men were of robust physique, and could offer no explanation of the trouble. Questioning elicited the statement that from 2 p.m. till 4.30 p.m. they had been applying a winter-wash insecticide to trees in a peach-house. The solution, in a strength of one part to sixteen of water, was well worked into trunks and branches, using a motor-car spoke-brush, an instrument calculated to splash liberally the hands, forearms, and face. To reach inaccessible parts and small twigs against the wall, the brush was dispensed with and the bare fingers used. After work they found it difficult to remove the brown stains from the skin. F. K. had had previous experience of this wash, and "seemed to remember that it had made him feel sick before." He usually smoked while at work, but had not done so on this occasion.

A tin produced was labelled "'XL All' Liquid Insecticide. Nicotine 3.7 per cent. Caution: When spraying or dipping for a length of time, rubber gloves should be worn." Though printed in large type, neither caution nor other particulars had been read.

Stimulants were given, heat applied to the extremities, the bowels opened with calomel, and a dessert-spoonful of kaylene administered every fifteen minutes for three hours, by which time in each case vomiting ceased. F. K. was well in two days. A. B., who had done much less brush work and more finger-dipping, had some abdominal discomfort for a week, and was shaky on his legs for ten days.

A wide literature centres round nicotine poisoning in relation to tobacco smoking, the volatile alkaloid, existing as organic salts, being dissociated by heat. Toxic effects from smoking, however, are not referable to nicotine alone, as the volatile chemical compounds pyridine, picoline, and lutidine (all having a similar physiological action to nicotine) are likewise dissociated, while in addition, carbon monoxide in high proportion may be inhaled. Nicotine and the still more powerful oil nicotianin are both soluble in water and alcohol, and are readily absorbed through the skin.

The poison produces a very complex mixture of effects—successive stimulation and paralysis of a variety of central and peripheral structures, among these stimulation and paralysis of the receptive mechanism of striped muscle.¹ When anaesthetics were rudimentary, relaxing rather than anaesthetizing, tobacco was a favourite among analgesics administered per rectum. In recent times a case is recorded² of a convict who secreted an ounce of cut Cavendish tobacco in his rectum and successfully conveyed it past the searchers, but, failing to recover it, was four hours later in a state of collapse, with nausea, vomiting, and paralysis of both legs. He recovered. Toxic effects following the external application of the leaf are on record,³ chopped tobacco leaves being applied under a bandage to a varicose vein which had been accidentally wounded. The patient became prostrated and pulseless, with nausea and vomiting, but recovered after several days.

The nicotine content in the insecticide above mentioned is stated to be 3.7 per cent.; so far as can be ascertained, the proportion in proprietary sprays and washes varies from 2 to 38 per cent. "XL All" Fumigator, used for agricultural and horticultural purposes, has a nicotine

¹ Sollman and Hanzlitz: *Experimental Pathology*, 1928.

² *British Medical Journal*, 1901, vol. i, p. 1544.

³ *Lancet*, 1879, vol. i, p. 296.