healthful as possible, in the interests of the nation, apart from all considerations of the happiness of the individual.

In this health campaign every sort of method is adopted. There are lectures, leaflets, posters, films, and, last but not least, powerful articles from the pens of various discarded leaders of medical thought in the columns of certain organs of the press. The most promising sign about all these activities is the stress that is laid on the education of the young. Whether or no the performance keeps faith with that promise is a question which requires our serious consideration. It should always be borne in mind that health is more likely to be attained when it is not the direct object of pursuit. If we wish to instruct the individual, health education is more valuable if it teaches the main conditions of a healthy life, and forms those habits which are likely to attain it.

Everyone here will admit that the Boy Scouts and the Girl Guides are doing the most valuable work possible for those of school age, while the National Association for the Provision of Playing Fields is carrying on the work for the older children and the adolescents. Their success is due to the fact that health is not the direct object of pursuit, but that they do teach those habits which are likely to attain it. It has been claimed that previous health propaganda dots the i's and crosses the t's for all these movements. One can only insist that in so far as they succeed in making the young think of health with a capital H they are doing the very worst thing possible for the furtherance of their ultimate aim.

With regard to the posters here are three questions for the consideration of the meeting:

1. As they are at present do they do any good?
2. Could they be amended so as to do more good?
3. Are they edited with sufficient care?

The third question I would myself answer with an emphatic No.

One poster was put up in at least one child welfare centre depicting in gay and attractive colours a little girl reaching up to get hold of a kettle boiling on an open stove. The audience to be instructed consisted of children under 5 years old. One must admit that is a perfectly sound way of teaching children by bitter experience the maxim "The burnt child shuns the fire"; but surely if the health societies really think it necessary that this lesson has to be learnt by personal experience, it would have been more humane and probably less costly in child life to have instituted during health week, on the lines of the old vaccination stations, a *burning station* where each child was made to touch a red-hot poker. It would at least have been possible to regulate the dosage.

From the extended use that is made of lectures, articles in the press, etc., it is evident that it is not intended to restrict propaganda to the juveniles. No child could be expected to read and enjoy Mr. Bruce Lane's weekly letter in the *Daily Ailment*. Probably more, and not less, care is needed in the editing of these for the advice of adults. Because men and women have more reason to think of their health it is the more incumbent on us to see that our teaching does not fall into the easy but disastrous error of making them think of their ill health. No poster or lecturer ever reminds people to think of their manifold blessings; and yet all doctors in their daily work achieve more than half of their success by engendering in patients a cheery optimism, either because of the doctor's personality, or because, seeing that his doctor does not look worried, the patient decides for himself that all is well.

There are various bodies and societies which devote themselves to certain special aspects of the social health. For instance, the British Council of Social Hygiene, which runs the anti-venereal disease campaign. Its efforts are worthy of all praise, and it does achieve a large measure of success with regard to men; perhaps to a less extent with women. It does produce a certain amount of syphilophobia; but this is a small matter, since the complaint is so promptly cured as soon as advice is sought. In considering the success of this propaganda it must never be forgotten that it has an enormous advantage, in that it advocates early advice, and can point to an almost certain cure free of all expense if its advice is followed.

The position of the cancer leaflets issued by the Ministry of Health is by no means so satisfactory. It would be very valuable if this meeting would discuss thoroughly the advisability of the distribution of these to the public. The leaflets have not the value noted as appertaining to the anti-venereal disease campaign. We cannot point to any certain cure even in the early stages of the disease. We are not able to tell people how to distinguish those symptoms which may arise early enough in the course of the malady to give treatment every possible chance of success with the one exception of uterine haemorrhage after the menopause. On the other hand, we do know that if we circulate these pamphlets we shall cause an enormous number of the laity to be more terrified of cancer than they already are. My own view is definitely against circulation. There is one great disadvantage that a doctor suffers when ill: he knows too much about disease for his own peace of mind; the qualifying mitigation is that his training enables him to appreciate reasons against his worst fears. Why should we hand our disability to the public in an aggravated form, since we can give them part only of the knowledge and none of the training?

It is very difficult to harm the young by propaganda, since they care for none of these things; but with adults the position is vastly different. The particular class of adult which greedily devours everything about health it can get hold of is exactly that type to whom it is most harmful. This group of introspective people is already far too much occupied with the functioning of the various bodily organs. On all of them the effect is bound to be deleterious, to many it will be disastrous. A genuine hypochondriac is a sick man, and as such deserves our respect and care. But does not propaganda tend to produce his pinchbeck imitation the valetudinarian, who is of all men the most miserable, and of all human objects the most useless? So long as health is regarded as a means to a useful life, it is life's greatest boon; once it is made his aim in life, it is man's greatest curse. This danger is peculiarly inherent in the activities of that society which advocates constant regular medical examinations.

The infinitesimal good that might be gained is far outweighed by the enormous harm that is certain to be caused by making people overanxious. Take, for example, their suggestion that all persons should be examined before taking a holiday in order to see what they were fit for. What sort of a holiday would the golfer get after the doctor had told him to wear gumboots on a wet day, the angler who might go out only when the sun had full power, or the Leicestershire man told that he might not jump any fences? Why, the thing is an absurdity.

It is surely unnecessary to say anything on the subject of health weeks. If a man holds that health can only be taught by teaching healthy habits in youth, what can you expect him to say about a scheme which sets aside some hours of one week out of fifty-two for the consideration of the subject? The very idea of a health week is the complete reversal of all his views on health.

Lest you should say that no constructive suggestion has been put forward, may I give you a quotation from *How to Make the Best of Life*, by Dr. Glanville? It contains a truth much neglected at the present time; it is a better motto for a poster than any yet issued by a health society, and it is no bad creed for any man.

"The best way to live well is to work well. Good work is the daily test and safeguard of personal health."

III.—THE PRESS.*

BY

H. A. GWYNNE,

PRESIDENT OF THE INSTITUTE OF JOURNALISTS.

THE press and the medical profession in some respects are opposed to each other, for the former lives and has its being by purveying publicity, while the latter must by the nature of the functions it performs be restrained and discreet. A common ground of agreement between these opposite aims is somewhat difficult to discover, but I am

* Read in a discussion in the Section of Medical Sociology at the Annual Meeting of the British Medical Association, Manchester, 1929.

inclined to think that health propaganda is perhaps the one subject in which the two can co-operate wholeheartedly. We both ardently desire to improve the health of the nation, and we must frankly admit that neither of us can do much in this direction without the help of the other.

I hope this conference will pardon a personal reminiscence, but I relate it only in order to put myself on good terms at the outset with my audience. I am, in a way, a pioneer, for in the year 1913 I published a letter, courageously signed by some of the greatest men of your profession, drawing attention to the grave consequences of the ignorance of the public on the subject of venereal diseases, especially syphilis. It was urged upon me by many eminent doctors and surgeons that such an indictment of our apathy and indifference was necessary in the interests of the nation. I took what little courage was left me by my friends in both hands, published the letter, and supported it warmly with a leading article. The result was that in a short time a matter of great and even vital importance to health was openly discussed and is now no longer confined to the technical columns of the medical press.

It may therefore be taken for granted that the press is only too anxious to march side by side with medicine in the fight for public health. Yet, in spite of this, there are some obstacles to be overcome before we can act together with complete understanding. The press will help in the good work, but it will always want to do so in its own way. The medical profession, bound and tied by strict rules and regulations, would like it done in its way, and there comes the first difficulty. Every newspaper seeks to have an individuality of its own and to carry this individuality right through its pages. For this reason, the concentrated wisdom of all the doctors, if it be distributed to the press generally, will be set aside in favour of the contribution of a lowly general practitioner with a lively pen. Let not this be counted against my profession. I have known of cases where a good-looking doctor with tenuous qualifications has snatched away the woman's portion of the practice of his colleagues in a few months.

Health is a matter of intense interest to everybody, and journalists, having quickly grasped the fact, are naturally eager to provide their readers with the best information they can. In the search for this they often find themselves face to face with the strict rules and regulations of the medical profession. Like many other institutions, such as the Law Society and the Stock Exchange, members are forbidden to advertise, and it has, I believe, been laid down that a signed article in a lay paper is an infraction of this rule. We are thus brought to an impasse. Newspapers require articles at odd times and on odd subjects. They search out the best specialist on the particular subject required, and in nine cases out of ten meet with a ready, not to say eager, response. Then comes the difficulty. It must not be signed by the author—that would bring the wrath of the gods on his head. The article is therefore published as from “a distinguished Harley Street specialist,” and all we get for our pains is a series of letters from our readers accusing us of inventing the specialist and writing the article in the office.

Any fair-minded journalist will readily grant that it would not be in the interest of medicine generally that doctors should advertise. It certainly would add to the attraction and revenue of a journal that Dr. X. should use the advertising columns to announce that he would “undertake to make the most melancholy sing in his bath,” or that Dr. Y. should say to the world “Don't let them cut you up; let Nature and Dr. Y. put you right without an operation.” One's imagination soars at the thought, and, like Clive, we are amazed at our moderation in not starting a press campaign in favour of the abolition of this rule.

Still, there is the problem. We journalists want authoritative information in a hurry, with a name at the top of it, and you of the medical profession would rightly see in that a dangerous innovation which might do infinite harm. It is to be remembered, however, that several

eminent doctors do contribute to the press under their own names and no protest appears to have been made by the General Medical Council. Doubtless you might argue that this small band of privileged writers should suffice for the needs of newspapers; but in point of fact they are not numerous enough to fulfil the requirements of the newspaper press to-day. It seems to me that the British Medical Association might well examine this matter and see whether it cannot evolve a solution of the difficulty. As things are at present I am convinced that medicine is losing many great opportunities of teaching the people the great lessons of health, so essential for the well-being of the country.

A cynic once remarked that one of the reasons why a doctor is called in is because the patient has the rare and refreshing opportunity of talking about himself and being heard with attention. Whether that is true or not the fact remains that there is nothing quite so interesting to newspaper readers as discussions on health. Rarely a day passes without a letter being received from a reader asking how best he or she can get fatter or thinner; acquire an appetite or lose some of the craving for food; where to find the best brand of vitamins; whether a whisky-and-soda at bedtime is good for insomnia; or which is the best doctor to consult for persistent halitosis. To our credit be it said that our invariable answer is “Consult a doctor,” and in the last case the reply was “Consult any doctor.” Such daily experiences are conclusive proofs of the fact that medical matters are, in the language of our craft, “good copy,” and therefore it needs no stimulus from outside to quicken our interest in them. However pontifically we may speak to the public on a great variety of subjects, we are quick to realize that it is highly dangerous to pronounce opinions on matters that concern the physical health of our readers. To help us we must turn to your profession, and often in vain.

It might well be suggested that, if there exist such an interest in medical matters among the newspaper reading public, it would be to the advantage of journalism to have a doctor on the permanent staff as general adviser and writer on all that concerns medicine. This suggestion may have been put into practice, though personally I cannot recall any instance of its adoption. There are objections to it which are fairly obvious. In the first place it is possible that a qualified doctor who accepted such a position on the staff of a newspaper might be running foul of some of the conventions of the General Medical Council. Even if it was otherwise, I am somewhat doubtful whether such an experiment would succeed. One of two things would happen: either the medical correspondent would absorb the spirit of his environment and become more of a journalist than a sober *terre-à-terre* doctor, or he would retain the restraint, reticence, and admirable dignity of his profession and be useless as a member of the staff of a newspaper. Besides all this, I doubt very much, except in cases of young doctors still in search of a practice, whether the career of journalism would be tempting enough for a man who has spent four to six years to qualify. It would be a bad thing for both medicine and journalism if incompetent men were engaged.

Owing to the restrictions to which I have referred we journalists are limited in our endeavours to act as interpreters for your profession to the people. We are driven, therefore, to the published words of eminent physicians or surgeons over whom the General Medical Council does not seem to exercise control. Here we are met with the difficulties of such a divergence of opinion as to render our efforts nugatory. Take diet for example. Sir A.B., a physician with a most distinguished record, says alcohol is poison. Sir B.C. makes a speech and pronounces it to be a most excellent and salutary stimulant when taken in moderation, and confesses that he himself always takes a nightcap. “Beer,” says Sir D.E., “is both drink and food. Taken in moderation it is an excellent digestive, a magnificent and safe pick-me-up, and should be on every table in the land.” Whereupon Sir F.G. declares that if a man has any sort of regard for his liver, he should not even smell an empty beer-pot, much less drink the vile stuff. If people will drink, let

them drink light wines, whisky, or egg-flip. Who will decide when doctors disagree?

So far, I am afraid that you will think that I have been a little lukewarm in my advocacy of joint action between the doctors and journalists. This, however, is not the case. If I have mentioned particular circumstances under which co-operation between the two has been somewhat difficult, I have no hesitation whatever in declaring that on the broad question of the presentation of the plain facts of health there can be, and there should be, the most hearty and constant collaboration of the two professions.

There will be moments in our national life when disaster can only be averted by the combination of medicine and journalism. Epidemics may rage again, and the people will require daily warnings and advice. When this happens I feel sure that the doctors can call on the press to help, and that call will receive an immediate and enthusiastic response. The seasonal ills of spring, summer, autumn, and winter are topics of interest to the press, and we gladly receive and print the advice so generously placed at our disposal by your great profession. Every day there is made some advance in the great science of medicine, and if, as is right and proper, you are reluctant to raise hopes too soon, or to divulge a secret until it has been thoroughly tried out, we, too, are willing to keep silent until the right time for discovery has come. To enforce the simple rules of health which are approved by your profession, to save the people from expensive and dangerous quacks, to do all that is in our power to make of our people the healthiest, as they are the sanest, in the world—these are noble aims worthy of the great professions to which you and I belong.

The Sections.

SUMMARY OF PROCEEDINGS.

(Continued from page 205.)

SECTION OF MEDICINE.

Thursday, July 25th.

TREATMENT OF PLEURITIC EFFUSION.

DR. H. J. STARLING (Norwich) presided over the second session of this Section, when Sir THOMAS HORDER read a paper on the treatment of pleuritic effusion. First enumerating the various factors of importance that had to be considered in any case, the speaker went on to uphold the traditional division of effusions into serous and purulent, and enumerated their various causes. He emphasized the fact that it was bad practice to remove all effusions at sight, whether they were serous or purulent; before the fluid was tapped the associated condition of the lung must be assessed as carefully as possible, and it must be determined how far the patient was embarrassed by the fluid itself. In a general way the more ill the patient the more carefully and slowly should the fluid be removed. Speaking more particularly of serous effusions, Sir Thomas Horder argued that too early intervention was meddlesome; if postponed till unduly late it might result in permanent collapse of the lung. Strapping of the chest was often advantageous, and so might be the exhibition of iodides and even calcium. Little good could be said of auto-serotherapy. Once tapping had been determined on, slow aspiration was preferable to siphonage, and a portion only of the effusion should be removed. Local anaesthesia and the preliminary use of a sharp scalpel were to be recommended. It must be remembered that most serous effusions were tuberculous in origin, and a prolonged convalescence was indicated. Turning to empyema, the speaker advocated the removal of the pus as soon as it was diagnosed. One or two aspirations might be successful in pneumococcal and in some septicæmic cases (streptococcal), should the patient be too ill to stand the resection of a rib, or open method, that was most generally advisable. But rarely could the septic cavity be opened, emptied, cleaned out, and closed up again with success. It was important to disturb the diseased lung as little as possible;

irrigation of the diseased cavity, though it had been tried again and again, was of but doubtful virtue, whatever the nature of the irrigating fluid. Dr. L. S. T. BURRELL confined his remarks to exudates of doubtful or tuberculous origin. The great majority of pleural exudates were due to tuberculosis, so that, unless there was any definite evidence to the contrary, it was safest to base treatment on the assumption of a tuberculous origin. The patient sought advice in most cases either because of acute pain from pleurisy—the effusion develops later—or because of dyspnoea and general malaise; he was found to have a pleural effusion. In the latter case it might be possible to obtain a history of previous pain for which the patient did not see a doctor, but in many cases there was no history of any pain or illness, and the sole complaint was increasing dyspnoea. In either case the patient should be put to bed and 5 c.cm. of the liquid should be aspirated for examination, because treatment was largely dependent upon the character of the effusion, and a cytological examination helped to determine the cause. Aspiration should not be performed except (1) to relieve pressure symptoms such as dyspnoea; (2) to convert the condition into one of artificial pneumothorax if the underlying lung was actively tuberculous; (3) to encourage the underlying lung to re-expand owing to active disease in the other lung; (4) to treat chronic or recurrent pleural effusion. Removal of a quantity of liquid from the chest was not free from danger, and had led even to sudden death. More common complications, however, were violent paroxysms of cough, severe dyspnoea, oedema of lung, or activation of a tuberculous lesion which was previously quiescent in the lung. In order to prevent these dangers it was wise to introduce air or oxygen into the pleural cavity if more than 15 ounces of liquid were withdrawn. Air should be used if it was desired to maintain an artificial pneumothorax, and oxygen if one wished the lung to re-expand, because oxygen was more rapidly absorbed than air. In an acute case with high fever and toxæmia the effusion was the result and not the cause of the condition, so that although it was often necessary to aspirate in order to relieve pressure symptoms resulting from a rapidly accumulating effusion, in the absence of such pressure symptoms it was very rare that aspiration would relieve the toxæmic symptoms; indeed, it would often make them worse. Most of the acute cases settled down and did best without aspiration. In the very chronic or recurrent cases aspiration should be performed and the pleural cavity gently irrigated with Dakin's solution, oxygen introduced, and a negative pressure of about 6 cm. water left. The general condition of the patient should be treated by fresh air, good food, and, if necessary, aperients, tonics, etc., but no drug had any effect on the actual effusion. The after-treatment was of the greatest importance, for on it might depend whether the patient became consumptive; 40 per cent. of such cases subsequently developed pulmonary tuberculosis, and sanatorium treatment was the best for them. Professor A. RAMSBOTTOM (Manchester) said that all the evidence went to show that idiopathic pleurisy was in the vast majority of cases tuberculous; cold and injury, important medico-legally, were merely predisposing. It was noticeable how few soldiers, after chest wounds in the war, contracted pulmonary tuberculosis. Although by ordinary methods the bacilli could seldom be demonstrated in the fluid, by modified methods the bacilli had been found in seventeen consecutive cases. These observations were also borne out by inoculation experiments, and by following up the cases (43 per cent. within six years). Tuberculous pleurisy was rarely if ever primary; it was secondary to tuberculosis elsewhere, usually the lungs. The presence of pleural effusion was usually readily determined by clinical examination. It was frequently noticed that the affected side was smaller than the sound side, due no doubt to the fact that when effusion began the cohesive property of the pleural surfaces was lost; the lung collapsed and over-prepared for effusion, with the result that the chest wall fell in. Bulging of intercostal spaces, or increase in size of the affected side, usually indicated positive intrathoracic pressure. This collapsing of the lung on the affected side played a most important part in the displacement of the heart in the early stages, and in cases