ORGANIC GASTRIC SYPHILIS.

BY

DAVID SMITH, M.D., F.R.F.P.S.,

ASSISTANT PHYSICIAN, ROYAL INFIRMARY, EDINBURGH.

(With Special Plate.)

ORGANIC gastric syphilis has received very little study or attention in Great Britain. This is due in no small degree to the rarity of the condition and to the difficulty of obtaining confirmatory pathological evidence. The sceptic is apt to say that the lesion is a coincidence, and is simply some recognized condition occurring in a syphilitic patient. On this account it is all the more necessary for a definite diagnosis to be established between gastric symptoms occurring in a syphilitic subject and actual syphilis of the stomach. It does not necessarily follow that every patient with gastric symptoms and a positive Wassermann reaction is suffering from organic gastric syphilis, even if an organic lesion can be demonstrated. It must be remembered that both ulcer and neoplasm occur in syphilitic patients far more frequently than gumma does. Reference to the work of Chiari¹ and of Crohn² proves this.

History of the Disease.

In previous cases which have been recorded, the diagnosis was made from the fact that other specific lesions were present along with gastric symptoms, and because these cleared up under antisyphilitic treatment the diagnosis of "syphilis of the stomach" was made. I do not dispute the probability of the diagnosis being correct, but it is at least controversial. Into this category I put such cases as those described by Andral³ as early as 1834, and which Monod⁴ considers indisputable. The cases may have been syphilitic, but I cannot hold the opinion expressed by Monod that they are indisputable, because there are no data beyond the clinical history and the results of treatment to warrant the diagnosis. I admit that at that time no further data such as x-ray examination and Wassermann reaction could be given, but without such investigation no case can be considered indisputable.

Up till 1907 several cases of gastric syphilis were recorded by Continental authors, but proof is lacking that these cases actually showed syphilitic gastric lesions. At this date Pater,⁵ in a thesis, reviewed the then existing data, and also discussed 122 cases, rejecting those which did not present microscopical evidence of syphilis. He thus established the fact that cases with gastric symptoms sometimes have syphilitic lesions of the stomach. Since 1907 much has been written regarding syphilis of the stomach, but these contributions to medical literature have been made mostly by American and Continental writers. The majority of these records are based on radiography, the presence of syphilitic lesions elsewhere, a positive Wassermann reaction, or the results found at autopsy or operation. In 1912 Meyers⁶ reviewed fifty-eight cases, and the American literature has been added to by McNeil,⁷ Morgan,⁸ Downes and Le Wald,⁹ Smithies,¹⁰ Niles,¹¹ Eusterman,¹² and others. Eusterman's contribution is on twenty-three cases, and he reviews the literature. He establishes a sound case from the clinical standpoint. From the literature and the cases I have seen, I consider that syphilitic lesions of the stomach can, and do, exist. This is confirmed beyond all doubt, and with a certainty that should dispel all doubts, by McNee,¹³ who demonstrated the spirochaete in the lesion. Apart from this, British literature contains nothing but adverse criticism of the subject, with the exception of the cases referred to by Hurst,¹⁴ Fenwick,¹⁵ and Sherren.¹⁶ Muir¹⁷ states that "syphilitic lesions, though rare, occur both in the congenital and acquired forms of the disease." Bennett¹⁸ admits that gastric syphilis does occur, and urges that, "whenever one is confronted with a case of organic gastric disease the possibility of syphilis must be remembered." Wyard¹⁹ reluctantly accepts the possible diagnosis, and in spite of a degree of scepticism he states the case convincingly from the clinical standpoint. I cannot agree with his concluding remarks, however, that "the treatment of this disease is far from satisfactory, and that it may

properly be argued that its almost uniform refractoriness to recognized antisyphilitic methods is evidence of some weight that many or most of the cases so diagnosed are not, in fact, specific." The cases about to be described are in striking contrast to this statement.

Clinical Aspect.

I wish to focus attention on the clinical aspect, accepting the fact that pathologically and bacteriologically syphilis of the stomach is proved beyond all doubt, and therefore a clinical diagnosis should be possible and should be looked for oftener than it is. I do not hold the view that this diagnosis should be made because the Wassermann reaction is positive in a gastric case and because symptoms clear up under antisyphilitic treatment. It is a well-known clinical fact that diseases which cause a general constitutional disturbance frequently give rise to gastric symptoms, and these improve or get well with the increasing well-being of the patient. I do not propose to deal with the cases discovered at operation or at autopsy, because, strictly speaking, such are not in the province of the physician. I wish, however, by recording six cases which, in my opinion, are indisputable, to draw attention to the fact that syphilis of the stomach does occur, and that it can be cured by medical means, thus saving the life of the patient or averting operation.

Essentials of Diagnosis.

Before deciding that a patient is suffering from organic syphilis of the stomach the following points have to be satisfied:

1. Symptoms definitely pointing to disease of the stomach or duodenum must be given.
2. X-ray examination must show evidence of such disease.
3. The Wassermann reaction must be positive.
4. The symptoms must not yield at all to a "try-out" of recognized treatment for peptic ulcer.

Proof of the cases being genuine is furnished if

1. Symptoms clear up rapidly under light or ordinary diet in conjunction with intensive antisyphilitic treatment.
2. X-ray examination shows improvement or recovery.
3. The Wassermann reaction becomes negative or less positive.

It will be observed that confirmation of the diagnosis in the cases is obtained by assuming that the lesions are probably neoplastic, inflammatory, or ulcerative conditions in otherwise syphilitic subjects.

Review of the Cases.

The cases I describe were put on strict treatment in bed and the results were observed. No improvement took place after four weeks of treatment, and further x-ray examinations show the conditions unchanged. The patients were, with two exceptions, allowed up on full diet and treated with neokharsivan, potassium iodide, and mercurial inunction. In periods varying from seven to twenty-one days symptoms had disappeared and improvement in their general condition had begun. Treatment was continued, and the final x-ray photographs were taken as an indication of cure or very marked improvement. The patients are still well at the end of periods varying from one and a half to five years, and no further treatment has been given. Case 1 died of pneumonia three years after. During that time he remained well. Unfortunately no autopsy was allowed.

I am going to consider six cases only because in these no other lesion due to syphilis was found after thorough investigation. Five were males and one was a female. This corresponds roughly with the incidence in the sexes noted by American authors. The youngest patient was a man aged 36, and the oldest a woman aged 60. Marcus²⁰ records a case in a lad, aged 19. Two of the male patients admitted infection, one twenty years, and the other two years, previously. According to Meyers⁶ the alimentary lesion may appear as early as ten months after infection or as late as twenty-five years. None of the other patients admitted syphilitic infection, and no history could be obtained of a definite primary sore, except in the

two patients who admitted infection; no secondary or tertiary symptoms or signs were admitted. All six patients had a ++ or +++ Wassermann reaction. The shortest duration of symptoms was six months, and the longest eighteen months. The symptoms were progressive from the onset.

Symptoms.

The most frequent symptom is constant pain referred to the upper abdomen. The character of the pain varies from a mild discomfort to agonizing cramp-like seizures, which are superimposed on the general distress in the upper abdomen and come on some hours after food. These seizures are mostly unrelieved by taking food, but are somewhat relieved by vomiting. The underlying general distress is not at all alleviated by vomiting. In all cases the pain, whatever its character may be, is progressive, and the periodic remissions so frequently met with in gastric and duodenal ulceration are absent. In character, therefore, it more closely simulates the distress of gastric carcinoma. Nausea and vomiting are frequent symptoms. The vomitus does not contain blood so often as it does in carcinoma, and only one of the cases showed melaena. Rehfuß²¹ records four cases where blood was present more or less persistently in the gastric contents. Other observers have recorded fatal haemorrhages. Anorexia is common, and this, with the constant vomiting, leads to loss of weight, marked anaemia, and some degree of cachexia. The anaemia is often extreme, although there is no apparent loss of blood. Cachexia is not so marked as in cases of carcinoma. Jaundice occurs, but although I have seen several cases of this type I have not included them in this series.

Physical Signs and Gastric Analysis.

Physical examination of the abdomen may disclose a distinct mass caused by gummatous tumour formation in the antral or pyloric regions. This was well marked in two of the cases here recorded. There are, however, no physical signs peculiar to organic gastric syphilis.

Achlorhydria was demonstrated in five out of the six cases. The lesion in the case which showed free hydrochloric acid was a gummatous mass causing pyloric obstruction. This is in agreement with the findings of Downes and Le Wald.⁹ In this case there was almost complete retention of the gastric contents, and it was in marked contrast to the other cases, where the gastric contents left the stomach rapidly. There was no excess of mucus in any of the cases. Blood and lactic acid were absent; the analysis depends to a great extent on the situation and type of the lesion, and may be influenced by the general debility and anaemia due to the syphilitic infection, for syphilitic anaemia generally shows an achlorhydria. Some authors contend that the syphilitic virus in addition to the localized lesion may produce a general gastritis which is more responsible for the achlorhydria than is the lesion itself. This is rather negated in the cases here recorded by the fact that no excess mucus was demonstrated. Crohn²² thinks it is unlikely that any particular form of gastritis exists in these cases to a degree sufficient to cause an atrophy of the mucous membrane and a resultant anacidity or achylia gastrica. Section of the mucosa away from the actual gummatous area fails to show any unusual form of gastritis. The cells and membrana propria are not materially altered.

X-ray Appearances.

There is no definitely diagnostic x-ray picture of gastric syphilis. The situation and character of the lesion in the individual case resembles a stenosis, an ulcer, a filling defect, or an infiltration.

Diagnosis.

The diagnosis must be arrived at by careful consideration of all the available data. A purely clinical diagnosis is almost impossible, but in patients who give a history of definite gastric symptoms, especially when combined with achlorhydria, a careful examination of the blood, and if necessary the cerebro-spinal fluid, should be carried out

with a view to ascertaining the Wassermann reaction. As previously pointed out, we must not accept that organic syphilis exists even when x-ray deformities are found combined with achlorhydria and a positive Wassermann, for cancer and ulcer are far more likely to be present; the final word rests with the result of treatment. Anaemia and cachexia are rarely so marked as in carcinoma showing somewhat similar lesions. Sometimes syphilitic stigmata elsewhere will help in forming an opinion, although in the present series no such stigmata occurred; the absence of metastasis may help in differentiating syphilis from carcinoma.

Owing to lack of space the x-ray plates before and after treatment have been omitted in Cases II and III. They were easy of description, and this has been made in the text.

Summary of Cases.

CASE I.

Male, aged 57; ex-soldier. For one year complained of a constant gnawing pain in the epigastrium and left hypochondrium. The pain was not much affected by taking food, and was not aggravated by a change from a light to a heavy diet. He frequently vomited after meals, but without relief of pain. The pain was severe at night, but was not helped by taking food. His appetite was poor, and he lost 3 st. in weight during the year. He never noticed any blood in the vomitus, or darkening of the stools. No previous serious illnesses, and denied syphilis. Social condition good. He took no alcohol, and smoked in moderation.

On examination he was cachectic. Cheeks were sunken, and he looked emaciated; abdominal wall sunken; no tumour palpable; no tenderness; recti tense, especially on the left side; slight discomfort on palpating the left hypochondriac region; tongue clean; teeth bad; constipated; no enlargement of liver or spleen; all other systems healthy. No free hydrochloric acid was present; no lactic acid; no blood; no excess of mucus. Wassermann reaction +++; after treatment +-. X-ray examination showed filling defect and infiltration of the stomach wall, especially the greater curvature.

Figure 1 (see Plate) shows the condition before and Figure 2 after treatment.

CASE II.

Male, aged 38; miner. For eight months the patient complained of constant discomfort in the epigastric region. The discomfort became actual pain about one and a half hours after food. At irregular intervals he was seized with agonizing pain, which required morphine to relieve it, and the distribution of the pain suggested a gastric crisis. Vomiting frequently followed this agony, and gave relief, but the epigastric discomfort still remained. There was no marked difference between night and day symptoms. His appetite was poor, and he lost 2 st. in weight during eight months. No blood in vomitus and no melaena; no previous serious illnesses, and denied syphilis. Social condition good. He took no alcohol, and smoked in moderation.

On examination he was pale and emaciated. Abdominal wall tense; palpable tumour in the pyloric region; marked tenderness over upper right rectus in the area of the palpable mass; succession five hours after food; tongue clean; teeth good; no constipation; no enlargement of liver or spleen; all other systems healthy. Excessive resting juice; moderate mucus; high free hydrochloric acid; no blood; no lactic acid. Wassermann reaction +++; after treatment -. X-ray examination showed almost complete pyloric stenosis. Practically no barium had left the stomach in five hours. One month after treatment was begun x-ray examination showed that much of the barium had left the stomach in five hours, and there was only a small residue. Unfortunately this man had to go abroad, and further plates could not be obtained. The tumour could not be palpated when he left, and he had no symptoms. The stomach was emptying better, and showed greater motility. Treatment was continued, and he is still well at the present date, five years afterwards.

CASE III.

Male, aged 53; steel-worker. For one and a half years complained of flatulence three hours after food. The flatulence was relieved by taking food, but he was left with a feeling of weight in the epigastrium and right hypochondrium. At times he had acute pain, which was confined to the upper abdomen. The pain did not seem to have any relation to food, and sometimes came on during the night. He vomited now and again, and sometimes this gave partial relief. There was never any blood in the vomitus. On more than one occasion he had melaena. His appetite was poor, and he lost weight to the extent of 2½ st. No previous serious illnesses. Admitted having had syphilis twenty years previously. Social condition good. At one time took alcohol to excess, but for five years had been teetotal. Smoked in moderation.

On examination he looked pale and ill; he appeared to have lost weight, although not definitely emaciated; abdominal wall tense; no palpable tumour; marked tenderness over right rectus from the costal margin to the level of the umbilicus; no succussion; tongue clean; edentulous; very constipated; no enlargement of liver or spleen; all other systems healthy. No free hydrochloric acid; mucus normal; no blood; no lactic acid. Wassermann reaction ++; after treatment -. X-ray examination showed marked duodenal deformity, which entirely disappeared after six months' treatment.

CASE IV.

Male, aged 36; sailor. For six months complained of pain three to four hours after food. Relieved by taking food. Nocturnal pain frequently present, and relieved by taking food. Pain was agonizing, and sometimes required morphine to relieve it. There was in addition to this pain a constant aching in the epigastrium and right hypochondrium. He vomited frequently without relief. No melaena, and no blood in vomitus. Appetite poor, and he lost 2 st. in weight. No previous serious illnesses. He admitted having had syphilis one and a half years previously, for which he had some medical treatment, but no injections. Took no alcohol, but smoked heavily.

On examination he was cachectic. He was very weak and appeared ill. The condition of the skin pointed to marked loss of weight. Abdominal wall tense; no palpable tumour; marked tenderness over right rectus from the costal margin to the umbilicus; no succussion; tongue clean; very constipated; liver and spleen not enlarged; all other symptoms healthy. No free hydrochloric acid; mucus moderate; no blood; no lactic acid. Wassermann reaction +++; after treatment -. X-ray examination showed irregularity of filling of lower part of the body of the stomach and duodenal deformity.

Figure 3 shows condition before and Figure 4 after two and a half months' treatment.

CASE V.

Female, aged 60; housewife. For one year complained of increasing pain in the epigastrium. The pain was constantly present, and was not affected by taking food. She frequently vomited, but without relief of pain. At times she had agonizing cramp-like seizures in the right epigastrium and hypochondrium suggestive of gall-bladder colic. Appetite poor, and she became bed-ridden on account of weakness due to emaciation. She was literally "skin and bone." She did not know how much weight she had lost, but it must have been considerable. No blood present in the vomitus; no melaena; no previous serious illnesses; denied having had syphilis. Social condition fair; took no alcohol, and did not smoke.

On examination she was markedly cachectic-looking and very emaciated. Abdominal wall lax and sunken. There was a visible tumour extending from the middle line well into the right hypochondrium. It moved slightly with respiration. It was slightly movable on palpation, and not particularly tender. Tongue clean; edentulous; constipated; no enlargement of liver or spleen; all other systems healthy. No free hydrochloric acid; mucus scanty; no blood; no lactic acid. Wassermann reaction ++; after treatment +-. X-ray examination showed filling defect in the antrum. Surgeon's opinion: inoperable carcinoma.

Figure 5 shows condition before and Figure 6 after treatment.

CASE VI.

Male, aged 54; labourer. For one and a half years complained of constant dull pain in the left epigastric region and left hypochondrium. The pain frequently became acute almost immediately after taking food. Vomiting occasionally took place. The vomitus never contained blood, and there was no melaena. Sometimes the acute pain lasted for a day or two at a time. Appetite poor, and he generally felt weak. He lost 2½ st. in weight in a year. He had had no previous serious illnesses, and denied having had syphilis. Social condition poor. Moderate drinker and smoker.

On examination he was cachectic and emaciated. There was definite tenderness in the epigastrium one inch above the umbilicus and diffuse tenderness into the left hypochondrium. Tongue clean; teeth bad; no constipation; no enlargement of liver or spleen; otherwise healthy. No free hydrochloric acid; moderate amount of mucus; no blood; no lactic acid. Wassermann reaction +++; after treatment -. X-ray examination showed definite ulcer on the lesser curvature.

Figure 7 shows condition before and Figure 8 after treatment.

Prognosis.

The prospect of cure in cases of gastric syphilis is good, provided intensive treatment is employed. This is true of all the cases recorded here, even where a palpable mass has been present. Improvement takes place almost at once, and within six months the most obstinate case failed to show physical or x-ray evidence of disease. Most of the cases were negative on examination at the end of two to

two and a half months. In this I agree with Einhorn,²³ who records a case with a palpable tumour which yielded to treatment in six weeks. Le Wald,²⁴ on the other hand, thinks that beneficial end-results are slow, and records a case which took nine years to bring about return to excellent health. The general opinion is that the condition is capable of cure or marked improvement. Eusterman¹² states that the total cured and improved is 69 per cent., and that treatment instituted during the exudative stages gives brilliant results.

Treatment.

Treatment can be ambulatory, unless the physical condition of the patient is too low. Food should be given in a manner that will not irritate the stomach, but almost any kind of food can be taken. Meat should be minced, fish and fowl creamed, fruit and vegetables given as purées, and bread, etc., should be toasted so as to be crisp and require mastication. If these points are attended to I do not think the nature of the diet requires any further attention. Neokharsivan in doses varying from 0.3 gram to 0.9 gram is given in one, two, or more courses, as the individual case requires. Potassium iodide has a marked effect, and doses up to 30 grams three times a day should be administered. Mercurial inunction should also be freely used. Patients appear to stand the intensive treatment well, and no trouble of any kind was met with. Iodide and mercury should be given, with short periodic rests, even after physical signs and symptoms have disappeared.

Summary and Conclusions.

1. Syphilis of the stomach, though rare, does exist, and in searching for a diagnosis should be considered oftener than it is.
2. Six cases are recorded, of which four denied having had syphilis, so that denial of a primary infection should not exclude the possibility of the diagnosis.
3. In most cases the disease is characterized by continuous pain, which is progressive, and has superimposed on it other pains of varying type.
4. Vomiting is frequent and haemorrhage rare.
5. A palpable tumour is sometimes felt, and there is marked loss of weight and moderate cachexia.
6. Definite x-ray deformity is seen, and gastric analysis shows an achlorhydria, except where there is pyloric stenosis.
7. Syphilis should be considered in all atypical cases of gastric disease, or where ordinary methods of treatment fail.
8. Prognosis is good.
9. Treatment should be intensive.
10. The diagnosis is confirmed by the results of treatment.

I should like to thank Dr. John Henderson, in whose wards some of the cases were treated, and Dr. A. Bruce MacLean, to whom I am indebted for the x-ray photographs.

REFERENCES.

- 1 Chiari, H.: *Internat. Beil. z. Wissensch. Med.*, lxxxvii, 421; *Festschrift Rud. Virchow*, 1831, ii, 295.
- 2 Crohn, B. B.: *Affections of the Stomach*, 1927, 789.
- 3 Andral: *Cliniques*, Paris, 1834, ii, 201.
- 4 Monod: *Proc. Roy. Soc. Med.*, 1922, xv, Part I.
- 5 Pater: *Original Thesis*, 1907.
- 6 Mevers, J.: *Syphilis of the Stomach*, *Albany Med. Ann.*, 1912, xxxiii, 553.
- 7 McNeil, H. L.: *Syphilitic Ulcer of the Stomach*, *Journ. Amer. Med. Assoc.*, 1915, lxi, 430.
- 8 Morgan, W. G.: *Syphilis of the Stomach*, *Amer. Journ. Med. Sci.*, 1915, cxlix, 392.
- 9 Downes, W. A., and Le Wald, L. T.: *Syphilis of the Stomach*, *Journ. Amer. Med. Assoc.*, 1915, lxi, 1824.
- 10 Smithies, F.: *Syphilis of the Stomach*, *ibid.*, 1915, lxi, 572.
- 11 Niles, C. M.: *Syphilis of the Stomach*, *ibid.*, 1916, lxi, 564.
- 12 Eusterman, G. B.: *Syphilis of the Stomach*, *Amer. Journ. Med. Sci.*, 1917, clii, 21.
- 13 McNeel, J. W.: *Quart. Journ. of Med.*, April, 1922, lix.
- 14 Hurst and Stewart: *Gastric and Duodenal Ulcer*, 1929, III B, 242.
- 15 Fenwick, S.: *Lancet*, 1901, ii, 835.
- 16 Sherren: *Ibid.*, 1921, ii, 683.
- 17 Muir, R.: *Textbook of Pathology*, 1929, 477.
- 18 Bennett, I.: *The Stomach and Alimentary Canal in Health and Disease*, 1925, 157.
- 19 Ward, S.: *A Handbook of Diseases of the Stomach*, 1927, 249.
- 20 Marcus, J. M.: *Med. Clin. North America*, 1925, ix, 717.
- 21 Rehfus: *Diseases of the Stomach*, 1927, xliii, 901.
- 22 Crohn, B. B.: *Affections of the Stomach*, 1927, 799.
- 23 Einhorn, M.: *Syphilitic Tumour of the Stomach*, *Journ. Amer. Med. Assoc.*, 1902, xxxix, 1051.
- 24 Le Wald, L. T.: *Amer. Journ. Roentgenol.*, 1917, iv, 76; *Radiology*, 1923, i, 193.

DAVID SMITH: ORGANIC GASTRIC SYPHILIS.

FIG. 1.—The stomach of Case I before treatment.

FIG. 2.—The stomach of Case I after 2 months' treatment.

FIG. 3.—The stomach and duodenum of Case IV before treatment showing filling defect of lower part of stomach and duodenum. Palpation failed to make any difference.

FIG. 4.—The stomach and duodenum of Case IV after 2½ months' treatment. The stomach and duodenum now fill well.

DAVID SMITH: ORGANIC GASTRIC SYPHILIS.

FIG. 5.—The stomach of Case V before treatment. Note the antral filling defect in the position where there was a palpable tumour.

FIG. 6.—The stomach of Case V after 2 months' treatment. Note the well filled antrum. No tumour could be palpated.

FIG. 7.—Stomach of Case VI before treatment. Note the lesser curvature ulcer.

FIG. 8.—Stomach of Case VI after 2 months' treatment. The ulcer on the lesser curvature has healed.