ment of dosage intensities by Lee and Pack, following the plan employed by Martin and Quimby in their dosage intensity calculations for intra-oral cancer, suggests the need for greater relative intensities in breast cancer than in mouth cancer. They have demonstrated at the same time that these required doses of intensities, well beyond ten erythemas, can be carried through practically. Their minimum standards of intensity have not yet been determined, and it may be found that a very considerable percentage of the cases respond to a much lower relative quantity of irradiation.

In the lymphatic, glandular, neoplastic, and pseudo-malignant diseases—lymphosarcoma, Hodgkin's disease, and leukaemia—radium may be useful, and is occasionally of real value for some specific reason. In general, however, the widespread character of the involvement lends itself best to treatment by α -radiation. The treatment is palliative only.

The bone tumour problem is as discouraging as it is fascinating. Radiation has made but two distinct contributions, attributable alike to either α rays or radium, or both. Giant-cell tumours of bone are curable by external irradiation. The peculiar response of endothelial myeloma of bone to radiation led originally to its recognition as a definite clinical entity, and irradiation has remained the treatment of choice. The tendency to early and widespread dissemination of this growth, however, renders cure extremely doubtful. The true osteogenic tumours of bone are extremely radio-resistant, and radium has done nothing more than afford a moderate degree of palliative relief and growth restraint.

"DECEREBRATE RIGIDITY" IN MAN.

BY

J. S. MANSON, M.D., D.P.H.,

AND

FERGUS R. FERGUSON, M.D., M.R.C.P., D.P.H.,

ASSISTANT PHYSICIAN, MANCHESTER ROYAL INFIRMARY.

(With Special Plate.)

At the outset we wish to point out that, although the title of "Decerebrate Rigidity" in Man has been given to this short paper, we have made the publication rather to encourage interest and observation in this subject and to relate a very interesting clinical case than to record a detailed clinico-pathological study; for, from this latter point of view, our records are incomplete.

As Walshe⁶ points out, the variety of clinical features already included under this designation indicates that there has been no clear guiding conception of what a very definite condition true decerebrate rigidity is, and the term bids fair to become a vague label including all obscure forms of tonic spasm in a category which has a thoroughly false air of scientific precision.

Experimental Decerebrate Rigidity.

Since Sherrington⁷ detailed his experimental results of transection of the brain stem of animals and described the resulting picture of "decerebrate rigidity" or "reflex standing," there has been considerable difference of opinion as to what clinical picture constitutes its counterpart in man.

Experimental decerebrate rigidity occurs when the greater part of the anterior portion of the brain is removed by a coronal section through the hinder part of the mid-brain, in a plane bounded dorsally by the posterior limit of the superior corpora quadrigemina and ventrally by the region immediately caudad to the corpora mammillaria. This section cuts off the greater part of the red nucleus from the body; if the transection passes above the red nucleus, the muscle tone remains normal. Rigidity persists as the plane of the transection passes caudally through the pons and medulla to the level of entry of the acoustic nerves and Deiters's nucleus. Thus the brain stem centres responsible for decerebrate rigidity clearly lie in the pons and medulla.

Summary.

1. Radium has made a notable contribution and has added stimulus to the general problem of cancer investigation.
2. Since it is an agent for local application only, it must not be regarded as an ideal cancer cure.
3. It has, however, increased the curability of cancer.
4. It has reduced the limits and the severity of many operative surgical procedures.
5. It has supplanted some surgical operations entirely.
6. It has relieved operative surgery of the responsibility for attempting the unreasonable or impossible, as a frantic effort, in a large and varied group of cases.
7. It has made possible a larger measure of palliative relief in advanced cancer.
8. Advancement in radium therapy depends upon the intensive co-operation of pathologist, clinician, and physicist.
9. The technical problems of application and dosage intensity must be calculated and made with the greatest possible care and precision.
10. The combination of radium emanation, α -radiation, and operative surgery offers far more in the average case of cancer than adherence to any one agent alone: they are complementary, and not competitive.

REFERENCES.

- ¹ Failla, G.: *Amer. Journ. Roentgenol. and Radium Therapy*, xx, No. 2, August, 1928, p. 128.
- ² Ewing, James: *Amer. Journ. Path.*, March, 1929, v, No. 2, p. 99; *Amer. Journ. Roentgenol. and Radium Therapy*, February, 1926, xv, No. 2, p. 95.
- ³ Schmincke, A.: *Beitr. z. Path. Anat. u. z. allg. Pathol.*, 1921, lviii, 161.
- ⁴ Regaud, C.: *Compt. Rend. Soc. de Biol.*, 1912, lxxii, 523.
- ⁵ Quimby, E. H., and Pack, George T.: *Radiology*, 1929, xiii, p. 306.
- ⁶ Martin, H. E., and Quimby, E. H.: *Amer. Journ. Roentgenol. and Radium Therapy*, 1930, xxiii, No. 2, p. 173.

In this experimental decerebrate rigidity there is a great increase of tonus in the extensor group of muscles, with reflex inhibition of tonus in the antagonistic flexors in accordance with the laws of reciprocal innervation. On being suspended, the animal hangs with its fore limbs thrust backward, with retraction at the shoulder-joint, straightened elbow, and some flexion at the wrist. The hand of the monkey is turned with its palmar aspect somewhat inward. The hind limbs are straightened and thrust backward; the hip is extended, the knee very stiffly extended, and the ankle somewhat extended. The tail is straight, horizontal, or stiffly curved upward. There is a little opisthotonos of the lumbo-sacral region, and the head is kept lifted up against gravity. Sherrington regards this rigidity as reflex standing—the result of release or removal of inhibition of the postural reflex; it can be abolished by section of the appropriate afferent posterior nerve roots.

Decerebrate Rigidity in Disease.

The problem is to ascertain what is the counterpart in man, in a physiological sense, of this experimentally produced rigidity.

Kinnier Wilson¹³ has carefully analysed and recorded conditions—the results of cerebral haemorrhage, meningeal inflammation, or intracranial tumours—in which there has been, coupled with the impairment of consciousness, tonic rigidity of the trunk and limbs. He considers that these clinical pictures minutely resemble experimental decerebrate rigidity. In many of these cases frequent exacerbations are superadded in the form of tonic or postural fits, accentuating and exaggerating the background of tonic posture. The cases fall into three groups: (1) decerebrate rigidity (or attitude) and tonic fits combined; (2) cases of decerebrate attitude without tonic fits; (3) cases of tonic fits without persisting decerebrate attitude. Hughlings Jackson⁵ described attacks of rigidity in cases of cerebellar tumour. Many of these cases showed also tetanus-like seizures, in which the patient exhibited opisthotonos with extension of the lower extremities and flexion of the upper extremities, but he considered that the causation of the two conditions was different. In Wilson's opinion the involuntary postures of a limb or segment of a limb in many forms of nervous disease, as well as the transient positions assumed in involuntary movements such as chorea and athetosis, are in reality simply parts of the complete decerebrate attitude.

Walshe,¹¹ on the other hand, carefully studied the problem of Sherrington's decerebrate rigidity from the point of view of its occurrence and recognition in the human subject, and concluded that, while in the spasticity of hemiplegia and the paraplegia in extension, there is a clear example of hypertonus physiologically identical with that observed in the decerebrate animal; for the most part, the cases recorded as decerebrate rigidity in man are examples of different varieties of extrapyramidal syndromes or tonic fits, none of which can be identified with Sherrington's decerebrate rigidity. He recorded a case,¹⁰ which he considered a closer approximation to the experimentally produced condition than any previously encountered. The patient lay motionless and unconscious on her back, with the head in the median position. There was no trace of head retraction. The arms lay across the chest, semiflexed at the elbows, with the forearms slightly pronated and the wrists and digits flexed. The legs lay extended and adducted, with the feet plantar flexed. There was spasticity of moderate degree in all four limbs, definitely more pronounced in the arms than in the legs. The tonus was plastic and "clasp-knife" in character. The tendon-jerks were all very brisk, the abdominal reflexes absent, and the plantar responses extensor. The tonic neck reflexes of Magnus and de Kleijn were also obtained.

There are two points of defect from the complete picture of decerebrate rigidity: the first, the flexion posture of the upper limbs, and the second, the retention of temperature control. The flexed position of the upper limbs may be traced to the fact that in man the fore limbs are no longer simple locomotor props, but have taken on new and highly complex functions under the influence of the cerebral cortex. In consequence, they would not be expected to take up the extensor attitude of standing. In this case, it was at the stage of failing reflex activity, when the phenomena of decerebrate rigidity were finally disappearing, that tonic fits and the accompanying respiratory and cardiac irregularities appeared. This was regarded as demonstrating the fundamental difference between tonic fits and true reflex decerebrate rigidity, and as showing that there was no correlation between experimental decerebrate rigidity and the types of muscular spasm seen in extrapyramidal syndromes. Post-mortem examination revealed a suprapituitary cystic tumour and a well-marked secondary hydrocephalus; pressure from these on the structures in the anterior part of the brain stem produced the effect of decerebration by cutting off the cerebral hemispheres from functional connexion with lower levels.

The anatomical site of the lesions in Wilson's cases, with extended neck and spine, adduction at the shoulder, extended elbow, internal rotation, hyperpronation and flexion of the wrist and fingers, was similar to that of experimental decerebrate rigidity. Wilson regards the two states as physiologically equivalent, whereas other observers regard the hemiplegic posture in man—that is, flexion of the upper and extension of the lower limb—as the equivalent of decerebration in animals. Brain¹ considers that these views are not mutually irreconcilable. He has observed, although it is not always easy to demonstrate, that an extensor attitude of the upper limb, which exactly reproduces the posture of this limb in a "tonic fit," can be reflexly evoked in a hemiplegia by making the patient assume a quadrupedal attitude. This posture is the normal quadrupedal standing attitude of the anthropoid ape when on all fours. On the other hand, the "hemiplegic" attitude of flexion of the upper with extension of the lower limb is the primitive erect standing attitude of our subhuman ancestors. Man has passed through two stages in evolution: the more primitive characterized by quadrupedal standing, the later by erect standing. Thus, if by decerebrate rigidity is understood the posture of reflex standing, man may exhibit two decerebrate attitudes—quadrupedal and erect—the former of which (the attitude of "tonic fits") is the pathological equivalent of decerebrate rigidity in animals.

It will thus be seen that the cases appearing under the heading of decerebrate rigidity in the literature present considerable variation in the type of tone, in the posture assumed, and in the reflexes. Cobb,³ Weisenburg and

Alpers,¹² Zand,¹⁴ and recently Epstein and Yakovlev,⁴ have contributed, among others, to the literature on this difficult subject. In clinical practice, especially in children, cases are not rare in which the child is in such a fixed rigid attitude that it may be lifted into the vertical from the horizontal posture by placing one hand under the head. The actual position of the arms and legs in these cases varies considerably, but the type of tone increase appears to be more frequently extrapyramidal than pyramidal.

Notes on a Case.

The following case was under the care of one of us (J. S. M.) for a period of ten years, and during the greater part of the time he lay motionless and speechless, like a log, with his arms adducted and flexed at the elbows and wrists, and his legs rigidly extended—in the position regarded by Walshe and others as the physiological counterpart of experimental decerebrate rigidity.

A. B., male, was observed in November, 1918, at the age of 67, to incline to the right when out walking. Early in 1919 he complained of stiffness in his left shoulder and pain in the left leg, so that he had great difficulty in dressing. He was treated at Buxton for rheumatism, but returned unrelieved. About this time he lost all interest in his business and in his surroundings; he was frequently disorientated as to time and place, and would hold the newspaper upside down, although apparently reading it. He slept readily, and did not now complain of pain. Early in 1920 he exhibited great emotional distress at the death of his son, aged 24. A neurologist saw him in April, 1920, and noted a peculiar stiffness and slight weakness of the left side of the body, with ankle-clonus and an extensor plantar response. He considered the condition to be due to cerebral arterio-sclerosis, with softening in the right internal capsule. A. B. became gradually worse; the right arm became stiff, and he was only able to walk about the room with great difficulty.

In 1921 he became bed-ridden, and his apathy to his surroundings was more pronounced. The lower limbs were rigidly extended, with marked exaggeration of the tendon reflexes, bilateral extensor plantar responses, and ankle-clonus; incontinence of urine and faeces supervened. He could still use his arms for feeding himself.

During 1922 his condition altered but little; he recognized his attendants, but seldom spoke. In 1923 he was unconscious for three days, and it was thought that he was going to die. Towards the end of the year his wife had a cholecystectomy performed; when she returned home early in 1924 and explained where she had been, he remarked, "You look as if you had had your wings clipped." He made no further remark, and lapsed into his usual silence and apathy. In June, 1924, the family moved to a house in a brighter and sunnier situation, but he made no comment; he was utterly indifferent and apathetic; the increase in tone in the upper limbs now became so marked that he had to be fed.

The years 1925 and 1926 passed without much change, except the increasing rigidity of all four limbs. The upper limbs were strongly adducted at the shoulders, semiflexed at the elbows, with pronation of the forearms and semiflexion of the fingers. The rigidity in the lower limbs was so marked that it was impossible to flex the limbs, and when an attempt was made to abduct one leg, the other followed in the same direction. There was some retraction of the head; swallowing was fairly good with slow feeding, and the ocular movements were apparently full. Trophic sores appeared on the left leg and foot in 1927; the pulse, at times, thin and thready, at times full and strong, varied in rate from 68 to 110. In 1928 his general condition deteriorated and his muscles wasted. He lay like a log, not speaking, and apparently not recognizing any of those in attendance. His wife used to hear him mumbling in his sleep, but she could never understand what he said. It is interesting to note that in June, 1928, when one of his sons brought his 3-year-old twin daughters to see their grandfather, seeing one and not the other, A. B. said, quite clearly, "Where is the other?" He had not spoken since 1924, when he made the cryptic remark about the clipping of his wife's wings. He died on September 17th, 1928, at the age of 77, ten years after the onset of the first symptoms, and after having been bed-ridden and practically speechless for seven years.

In the post-mortem, permission was granted for the examination of the brain only. On removing the calvarium it was seen that the dura mater was thickened and excessively vascular in the right pre-Rolandic area; it covered, and was adherent to, a firm, rounded tumour, about two inches in diameter. The tumour had all the appearances of an endothelioma (growing from the falk cerebri). As is usual with these tumours, the surrounding brain tissue gave the appearance of having been hollowed out by the tumour. Beneath the tumour there was a softening extending into the basal ganglia; the medial surface of the opposito

hemisphere showed the imprint of pressure from the tumour, but the anterior cerebral artery was not occluded. The ventricle in the left hemisphere was somewhat dilated; there was no obvious softening in the left hemisphere, and the pons and medulla appeared normal. The cerebral vessels showed only a slight degree of atheroma. The specimen was sent to Sir Arthur Keith, and is now in the Royal College of Surgeons Museum (5018.2).

Microscopical examination confirmed the endotheliomatous nature of the tumour, and showed some degeneration in the pyramidal tracts in the mid-brain and pons, but not in the medulla oblongata.

We regret that we have no notes of careful observations on the type of tone increase or of the tonic reflexes of Magnus and de Kleijn and other observations, which, correlated with a complete pathological examination, would have made a very valuable record. In view of the incompleteness of the clinical details, a complete serial sectioning of the mid-brain and pontine regions was not considered warranted.

The early rapid progressive dementia and other unusual features encountered in this case, in contrast with those usually found in a frontal endothelioma, make it probable that a considerable portion of the symptomatology was due to cerebral vascular disturbances. Jefferson,⁶ in recording states of bilateral rigidity—which were accompanied by “tonic seizures” consisting of sudden accesses of rigidity—in middle meningeal haemorrhage, regarded the generalized rigidity as the result of disturbance of a very fine circulatory balance or a relative anoxaemia sufficient to suppress cortical control, but not great enough to render inactive the causal mechanism in the mid- and hind-brain.

We would call attention again to the two apposite utterances in 1924 and 1928, which contrasted very strongly with his otherwise complete silence and apathy for about seven years. It seemed that when an external stimulus was powerful enough to revive a thought or idea to a certain degree of intensity, it could be expressed in spoken words. It is interesting to speculate whether these gleams of comprehension could be explained by Spearman's theory⁸ of the two factors “g” and “s”—“g” representing the general mental energy, and “s” the special group of neurones activated by that energy. In A. B. the “g” factor was too feeble as a general rule to activate any special group of neurones “s” to such an intensity that utterance was possible.

Attention has been called again to the divergent views which are held as to what constitutes clinical decerebrate rigidity. Can these views be reconciled? Can we explain why these cases, which broadly might be described as being characterized by generalized rigidity, present such differences in the posture assumed and the type of increase of tone? In contrast to the case of A. B., recorded above, one of us (F. R. F.) has under his care at the present time a boy, aged 12, with the “decerebrate attitude” picture of Kinnier Wilson, probably the result of encephalitis. He came under medical observation two years ago for attacks of falling backwards, which were regarded as petit mal. There was no loss of consciousness, and his general and mental states were quite good. At the present time (as he lies in bed) his head is lifted forward, his lower back is slightly arched forward, his arms are extended, adducted, and pronated with slight rapid rhythmical tremor of the left hand. The legs are rigidly extended and the feet are markedly plantar flexed. He moves very little voluntarily, but can be lifted and turned like a “plank of wood.” When placed erect, he stands on tip-toes, and would fall backwards if not supported. His smile is slow, stiff, and spastic; intelligence is practically normal. There is generalized muscular rigidity. The tone increase is of the typical cogwheel type, and the limb assumes the posture into which it is placed for a few minutes before returning to the original position. There is no reflex evidence of involvement of the pyramidal tract—the abdominal reflexes are brisk, knee-jerks normal, and plantar responses flexor. He shows the tonic neck reflexes of Magnus and de Kleijn. From time to time he has “tonic fits” in which the abnormal posture is exaggerated.

Wilson says: “In the condition which has been regarded as decerebrate rigidity in man—that is, one of complete interruption of function, both cortico-spinal and strio-spinal, produced by lesions of various kinds, roughly speaking at

the level of the mesencephalon—what do we find? Since the striatal system is out of action, we might expect the universal rigidity of the post-encephalitic Parkinsonian, but it does not develop. Instead, we get an attitude in extension, which may fairly be regarded as indicative of an attempt at ‘reflex standing.’ In other words, the posture produced by release of function, when both old and new systems cease to act on the spinal neuraxis simultaneously, is that of release neither from the one nor from the other.” From clinical data, he has argued that the results of complete single release (hemiplegia and paraplegia) are not identical with those of complete double release (decerebrate rigidity) where both cortical and striatal efferent systems are affected. It would appear that the resulting picture depends on the degree to which these two systems are affected and their relative importance in a particular subject—animal or human, young or old.

It is hoped that the recording of this undoubtedly interesting clinical case will stimulate interest in this problem of “generalized rigidity,” and will lead to more complete examinations, which in turn will help to solve the problems of the motor pathways and the lesions which result from disease of them. In the current number of the *Journal of Neurology and Psychopathology*, Carmichael² has given an excellent short account of a human mid-brain preparation—the type of case to which generally little or no attention is paid from the point of view of scientific interest.

Practitioners frequently have not the opportunity of reading the works of experimental physiologists and of studying posture in detail, but it is hoped that by calling attention to clinical decerebrate rigidity, we will give neuropathologists a better opportunity for studying the problems of the motor system in the light of accurate, full clinical histories, and neurological examination.

It is a pleasure to acknowledge our thanks to Dr. E. E. Bowden for his assistance in making the post-mortem examination, and to Dr. Cecil Beadles for his interest in the specimen and the accompanying illustration.

REFERENCES.

- 1 Brain, W. R.: On the Significance of the Flexor Posture of the Upper Limb in Hemiplegia, with an Account of a Quadrupedal Extensor Reflex, *Brain*, 1927, i, p. 113.
- 2 Carmichael, E. A.: A Study of a Human Midbrain Preparation, *Journ. of Neurol. and Psychopath.*, 1930, ix, p. 324.
- 3 Cobb, S.: Spastic Paralysis in Children, *Med. Clin. N. Amer.*, 1920, iv, p. 417.
- 4 Epstein, S. H., and Yakovlev, P. I.: A Case of Decerebrate Rigidity with Autopsy, *Journ. of Neurol. and Psychopath.*, 1930, ix, p. 295.
- 5 Jackson, J. Hughlings: Case of Tumour of the Middle Lobe of the Cerebellum; Cerebellar Paralysis with Rigidity (Cerebellar Attitude); Occasional Tetanus-like Seizures, *Brain*, 1906, xxix, p. 425.
- 6 Jefferson, Geoffrey: Bilateral Rigidity in Middle Meningeal Haemorrhage, *British Medical Journal*, 1921, ii, p. 683.
- 7 Sherrington, C. S.: Decerebrate Rigidity and Reflex Co-ordination of Movements, *Journ. of Physiol.*, 1897, xxii, p. 319.
- 8 Spearman, C.: *The Nature of Intelligence and the Principles of Cognition*, p. 4.
- 9 Walshe, F. M. R.: *Med. Science Abstracts and Reviews*, 1921-22, v, p. 426.
- 10 Idem: A Case of Complete Decerebrate Rigidity in Man, with Observations on Various Reflex Reactions, *Lancet*, 1923, ii, p. 644.
- 11 Idem: The Decerebrate Rigidity of Sherrington in Man, *Arch. Neurol. and Psychiat.*, 1925, x, p. 1.
- 12 Weisenburg, T. H., and Alpers, B. J.: Decerebrate Rigidity following Encephalitis; Case with Necropsy, *Ibid.*, 1927, xviii, p. 1.
- 13 Wilson, S. A. K.: On Decerebrate Rigidity in Man and the Occurrence of Tonic Fits, *Brain*, 1920, xliii, p. 220.
- 14 Zand, N.: Decerebrate Rigidity from a Clinical and Physiological Standpoint, *Journ. Nerv. and Mental Dis.*, 1928, lxxvii, p. 105.

CYST OF THE URACHUS.

BY

ARCHIBALD RONALD, M.D., F.R.C.S. ED.

(With Special Plate.)

COMPLEXITY of development in the lower urinary tract affords ample opportunity for the occurrence of congenital abnormalities. Persistence of the intra-abdominal part of the allantois from the bladder to the umbilicus as a patent urachus, or the imperfect obliteration thereof with cyst formation, are among the less common varieties. Urachal cysts attract attention either from their increasing size or from the advent of infection. Some cases are discovered accidentally either at operation or at autopsy. The case reported below was unsuspected prior to operation. The patient, a young adult male, aged 26, had not suffered from any previous illness.

J. S. MANSON & FERGUS R. FERGUSON: "DECEREBRATE RIGIDITY" IN MAN.

Longitudinal section of Brain of A. B. showing tumour and softening.

A. RONALD: CYST OF THE URACHUS.

FIG. 1.

FIG. 2.