Observations

ON

THE MECHANISM OF ABDOMINAL PAIN.*

BY

CHARLES H. MAYO, M.D., D.Sc., LL.D.,

HON. F.R.C.S.,

SURGEON, ST. MARY'S HOSPITAL, ROCHESTER, MINNESOTA, U.S.A.

THE subject of abdominal pain is a most interesting and extensive one, and one which may be discussed from many points of view. The work on the nervous system is one of the glories of British physiology. Of late we have come to hear more and more about the involuntary nervous system, so that now the prize-fighter is not the only one who appreciates the importance of the solar plexus. In America our interest in the sympathetic nervous system was greatly stimulated a few years ago by the visit of Royle and Hunter of Australia. Encouraged by their work with spastic leg muscles, a number of surgeons throughout the world have revived the attempts to treat angina pectoris by section of a number of branches of the cervical sympathetic nerve, and are attempting to control hyperacidity and pylorospasm by section of one or more branches of the vagi. These involuntary nerves with their ganglionic hook-ups make me think often of radio apparatus with each tube amplifying the message that is passed. If, however, we do make use of this rather fantastic simile we must think of each ganglion as a group of stations all broadcasting at one time.

There has been much controversy over the nature of the stimulus which produces visceral pain, and Mackenzie and Lennander have actually claimed that pain is felt not in the viscera, but only in the overlying structures. Although some recent studies are tending to revive this view, it is to-day pretty well discarded, and when care is taken to avoid stimulation of nerve-endings in the parietal and visceral peritoneum it can be shown that the structures in the abdomen are sensitive to pulling, clamping, and tying. This is particularly true of the gall-bladder, the urinary bladder, the connective tissue around the pancreas, and the large and small omenta. The fact that the stomach, the intestine, the liver, gall-bladder, pancreas, and spleen appear insensitive to cutting and pinching does not prove that they are incapable of giving rise to pain, because it may be that there is no physiological necessity for these organs to respond painfully to the type of stimulus used.

In general it may be stated that most abdominal pain that is not due to peritoneal involvement is dependent either on undue contraction of a viscus, which brings pressure to bear on nerves in its wall, or on extreme distension, which leads to stretching of the nerves. This would explain the fact that pain is felt during the passage of abnormally intense peristaltic waves, such as those seen sometimes in the colon. That pain may be felt also during relaxation of a peristaltic contraction has been shown by the recent work of Payne and Poulton on the oesophagus. They believe that in a distended muscular tube peristaltic contraction will actually relieve pain if it happens to shorten the length of the muscle and thus relieve tension on stretched nerve-endings.

Kappis has emphasized Meyer's observation that distension of the intestine with gas will produce pain only when the mesentery is stretched and pulled on. Still another explanation for visceral pain may be found in the anaemia of the wall of the intestine when it contracts down so as to resemble a hard white cord. It may be that this can produce the type of pain that is felt in thawing fingers after they have been nearly frozen. Possibly in this case the stimulus is a chemical one—that is, some substance formed in the tissues when they are deprived of oxygen. It should also be noted that the blood vessels are sensitive to pain. Thus puncture of an artery, especially one of the smaller ones, will cause dull pain of a peculiarly unbearable type, and this may be accompanied by reflex sweating and nausea.

Even the solid abdominal viscera appear at times to be

sensitive to pain. Thus Foerster noted during the war that while a piece of shrapnel was being removed from the liver of a soldier complaint was made of a painful sensation in the organ. It is well known that the ovary is sensitive to pressure, and the stretching of the tunica albuginea by the growing follicle often produces pain which reaches its maximum just before the structure bursts.

Turning to the anatomical side of the problem, it should be remembered that the parietal peritoneum is innervated mainly by the intercostal nerve, and that the phrenic nerves innervate not only the diaphragm, but also the peritoneum on the under surface of the diaphragm. The sensory fibres in the visceral peritoneum and in the abdominal organs pass back to the spinal cord in association with the sympathetic nerves. The fibres from the stomach, duodenum, caecum, ascending colon, pancreas, spleen, liver, and gall-bladder return with the greater splanchnic nerves, while those from the distal half of the colon go back through the hypogastric and the pelvic nerves.

That the splanchnic nerve contains pain fibres can readily be demonstrated in either experimental animals or man. Even when general anaesthesia is used any stimulus applied to the nerve is likely to produce reflex movements or actual waking. It is easy to show also that splanchnic anaesthesia, or anaesthesia of the abdominal plexus, will render the whole visceral peritoneum, stomach, duodenum, liver, and gall-bladder quite insensitive to pain.

Langley has stated that the sensory fibres in the splanchnic nerves are white and medullated, and that they rise in posterior root ganglia in the thoracic region of the cord. They pass without interruption to the organs. He and Gaskell believed that the involuntary nervous system proper consists only of connector and effector neurones. The pain-carrying fibres, therefore, do not belong to this system, but are true sensory fibres similar to those found in the rest of the body.

It is very doubtful if the vagus nerves ever have anything to do with the transmission of pain sensations from the abdomen. It was frequently noted during the war that when the upper part of the thoracic portion of the spinal cord was severed by a gunshot wound, peritonitis and other abdominal diseases ran their course without pain. Stimulation of the central end of the cut vagus was shown by Foerster to cause nausea, but not pain.

It has long been known that disease in the gall-bladder or the liver may be accompanied by severe pain in the shoulder and in the side of the neck. This so-called viscerosensory reflex suggests strongly that the phrenic nerve must have something to do with pain in the upper part of the abdomen, and Ramström has described a branch which unites with the sympathetic fibres of the coeliac plexus to form the phrenic plexus. On the right side these nerves run to a special structure, the phrenic ganglion.

As is well known, the pains of peptic ulcer and of cholecystitis are not always felt in the same place in the epigastrium and back. The site of pain may vary from patient to patient, or from time to time in the same patient. This is not surprising when one remembers how many complicating factors there may be. Cholecystitis in many cases is associated with hepatitis or pancreatitis, and an ulcer near the incisura, or a gall-stone, may produce painful spasm of the pyloric muscle. The perforation of an ulcer into the pancreas or into the lesser peritoneal sac should also change the type and situation of the pain, because another group of sensory nerves then become involved. It is conceivable also that differences in the anatomical distribution and course of nerves in different persons might cause differences in the distribution of pain produced by ulcer or cholecystitis.

That our knowledge in regard to the mechanism of abdominal pain is still inadequate has been realized many times by surgeons as they have attempted to relieve the gastric crises of tabes. Those crises may perhaps be divided into three groups: (1) those in which pain, vomiting, and hyperalgesia of the abdominal wall are the most prominent symptoms; (2) those in which nausea and vomiting predominate, and in which there may also be pain in the larynx, ear, and heart, and perhaps bradycardia; and (3) those in which there is, in addition to the abdominal pain, hiccup, and pain over one or both

* Made in opening a discussion at a joint meeting of the Sections of Surgery and Physiology and Biochemistry at the Annual Meeting of the British Medical Association, Manchester, 1929.

shoulders. In the first group we suspect that the splanchnic nerves are mainly at fault; in the second we suspect the vagi; and in the third we feel sure that the phrenic nerves are involved.

In all of these types nothing abnormal can be seen or felt during abdominal exploration. On roentgenographic examination it has been found by Groedel, Lüden, and Carnot and Bruyère that the stomach is unusually active during crises, and that the height of the paroxysm of pain corresponds to the greatest intensity of a contraction. Similar observations were made by Daniélopou with the help of recording balloons introduced into the stomach.

It is remarkable that denervation of the entire stomach rarely abolishes the crises. Foerster has cut all the posterior roots involved in the nerve supply of the stomach, and has sometimes failed to give relief. This must mean either that sensation is reaching the spinal cord through anterior roots, or else that it is arising in some part of the cord or brain, whence it is projected into the abdomen. Actually, in a few cases, after section of the posterior roots had failed to give relief, section of the anterior roots resulted in cessation of pain. In one case the induction of paravertebral anaesthesia, in which the rami communicantes were blocked, was not followed by relief of pain. In another case, after section of the sixth, seventh, eighth, ninth, and tenth thoracic roots had failed to work relief, both vagi were cut below the diaphragm and posterior gastro-enterostomy was performed, but even this extensive denervation failed to influence the severity of the pain.

So far as I know, the phrenic nerves have not yet been cut for the relief of gastric crises. The spinothalamic tracts, which carry sensory impulses up the spinal cord, have been sectioned, and some of the patients so treated have been relieved. It may be possible that the surgeon did not always cut deeply enough, because the fibres from the more cranial segments lie deepest in the cord.

The lesion in *tabes dorsalis* is in the posterior root gangions and in the dorsal columns of the spinal cord. Gastric crises, however, may be caused by a variety of conditions other than *tabes dorsalis*. They may be associated with inflammation of the membranes of the spinal cord, injury of the spinal cord from gunshot, or fracture of the spinal column, or they may even result from angioneurotic oedema. Eusterman has described cases of typical gastric crises which were seen in association with duodenal ulcer and angina pectoris. In tumours of the cauda equina the clinical picture may be that of crises involving the bladder, the colon, and the rectum.

I have gone so into detail with regard to tabetic crises because they illustrate how diverse may be the paths through which painful sensations may leave the abdomen and how little we really know about the subject. In some cases one can find zones of cutaneous hyperaesthesia or hyperalgesia. Ever since Head's classical studies it has been known that visceral disease may be accompanied by these zones, and sometimes they are so strictly localized as to indicate the presence of disease in some one of the internal organs. Surgeons know well the tenderness of skin and muscle without rigidity associated with disease of the diaphragm and upper part of the abdomen. They have long recognized also that early pneumonia in the lower lobe of the right lung may simulate appendicitis. I might state, parenthetically, that severe, long-continued pain in a localized abdominal area may be the result of bacterial involvement of the spinal nerve supplying the region and the muscle under it. The source of this infection will usually be found in the mouth, the cervix, or the prostate.

There are certain chronic lesions in the abdomen which produce pain of a type that does not fit well into any of the categories just mentioned. Much has been written about the cause of the pain in gastric and duodenal ulcer. Wilson found that spasm of the duodenal cap and pain occurred simultaneously in twelve out of fourteen cases of duodenal ulcer. Störek found in the bed of a gastric ulcer exposed nerves with neuromatous formations. The pain of gastric ulcer may be compared to the pain of a varicose ulcer of the leg. I believe the exacerbations of pain in these ulcers are the result of vascular spasms and later small extensions in the margin of the lesion. When these little extensions heal, as they do after a week or two,

the pain disappears. Certain peptic ulcers do not cause pain, and when pain has been present its disappearance does not necessarily mean that the ulcer is healed.

As compared with the skin, the abdominal viscera contain very few nerve-endings, so fairly large areas in these organs must be insensitive. Although, under local anaesthesia, one can incise the stomach without causing pain, I doubt if this experiment proves that the pain of gastric or duodenal ulcer is any different from that produced by local extension of a varicose ulcer of the leg. It may be, however, that the pain produced by a peptic ulcer is the result of a long-continued summation of stimuli which, if acting singly, would not affect the consciousness of the patient. Actually, Adrian has shown recently with the string galvanometer that a stimulus applied to a single receptor organ will produce a rhythmic disturbance in the nerve, a disturbance which is repeated from five to a hundred times a second. For simple end-organs, such as those which record pain, the duration of the volley of impulses was found to vary directly with the strength of the stimulus; the number of impulses in each second did not vary to any great extent. The significance of this observation lies in the fact that one method of overcoming synaptic resistance (and getting an impulse to reach the field of consciousness) is to keep repeating the message. In the spinal animal this is exemplified by the summation of inadequate stimuli, which finally results in, let us say, the scratching response. In cases of gastric ulcer, therefore, it may well be that it is the long-continued summation of inadequate stimuli that eventually produces pain.

Experience is invaluable, and yet, now that we are being helped so much by the roentgenologist and the laboratory worker, we must not forget that two or three decades ago pain was often a misleading as well as a leading symptom. What is needed is mass accumulation of data for careful analysis. I believe with Ryle that we need more knowledge concerning abdominal sensations, that we may more carefully elicit and interpret symptoms of disease in an earlier stage—in fact, anticipate and possibly prevent the advance of disease. With the abdomen open the surgeon, by splanchnic injection of anaesthetic solutions, frees the whole abdomen of pain, and yet local anaesthesia in the abdomen is secured by injecting, within peritoneal folds, the lesser or greater omentum or local mesenteric areas, and this is often an advantage over general anaesthesia.

I think I have said enough to convince you that when the character and site of an abdominal pain tells something to the surgeon, it does so not because he knows much about the underlying anatomy and physiology, but because he has noted that particular pain many times before, and he remembers what he found when he opened the abdomen.

BIBLIOGRAPHY.

1. Adrian, E. D.: The impulses produced by sensory nerve-endings. Part IV. Impulses from pain receptors. *Journ. Physiol.*, 1926-27, lxii, 33-51.
2. Carnot, M. P., and Bruyère, I.: Crise gastrique du *tabes* de type moteur avec hoquets et vomissements incoercibles hématomateux consécutives. *Bull. et Mém. Soc. Méd. des Hôp. de Paris*, 1917, xli, 1103-1110.
3. Daniélopou, D.: Le gastrogramme et le réflexe oculo-gastrique au cours des crises gastriques du *tabes*. *Ibid.*, 1923, ii, 1102-1107.
4. Daniélopou, D., Radovici, A., and Carniol, A.: Réflexe oculo-gastrique et oculo-colique. *Ann. de Méd.*, 1922, xi, 143-157.
5. Eusterman, G. B.: Duodenal ulcer simulating the gastric crises of *tabes*: report of five cases. *South. Med. Journ.*, 1925, xviii, 319-323.
6. Foerster, O.: *Die Leitungsbahnen des Schmerzgefühls und die chirurgische Behandlung der Schmerzzustände*. Berlin, 1927.
7. Gaskell, W. H.: *The Involuntary Nervous System*. New York: Longmans, Green and Co., 1916.
8. Groedel, F. M.: Quoted by Daniélopou.
9. Head, Henry: *Studies in Neurology*. London: Frowde, 1920, 2 vols.
10. Hunter, J. L.: The postural influence of the sympathetic innervation of voluntary muscle. *Med. Journ. Australia*, 1924, i, 86-89.
11. Kappis, Max: Beiträge zur Frage der Sensibilität der Bauchhöhle. *Mitt. a. d. Grenzgeb. d. Med. u. Chir.*, 1913, xxvi, 432-550. Die Sensibilität der Bauchhöhle. *Klin. Woch.*, 1925, iv, 2041-2047.
12. Langley, J. N.: The autonomic nervous system. *Brain*, 1903, xxvi, 1-26.
13. Lennander, K. G.: Abdominal pain: especially pain in connection with ileus. *Journ. Amer. Med. Assoc.*, 1907, xlix, 836-840.
14. Lüden, M.: Quoted by Daniélopou.
15. Mackenzie, Sir James: *Symptoms and Their Interpretations*. Third edition. London: Shaw and Sons, 1918.
16. Payne, W. W., and Poulton, E. P.: Experiments on visceral sensation. Part I. *Journ. Physiol.*, 1927, lxiii, 217-241.
17. Ramström: Quoted by Foerster.
18. Royle, N. D.: A new operative procedure in the treatment of spastic paralysis and its experimental basis. *Med. Journ. Australia*, 1924, i, 77-86.
19. Störek, O.: Ueber Nervenveränderungen im Narbenbereiche des *Ulcus pepticum*. *Wien. Klin. Woch.*, 1921, xxiv, 109-110.
20. Wilson, M. J.: Duodenal ulcer: observations on the behavior of stomach and duodenum in the presence of pain. *Arch. Int. Med.*, 1928, xli, 633-641.