Remarks

ON

SOME ASPECTS OF THE PROBLEM OF THE EPILEPSIES.*

BY

S. A. KINNIER WILSON, M.D., F.R.C.P.,

PHYSICIAN TO OUT-PATIENTS, NATIONAL HOSPITAL FOR PARALYSED AND EPILEPTIC, QUEEN SQUARE.

It is only five years since this Section of the British Medical Association, at the Bradford Meeting, was occupied with a discussion on the nature and treatment of epilepsy; and the fact that epileptic problems are again offered for consideration may properly be taken as proof of continuing interest in their examination and hope for their solution. Indeed, if we are to judge by recent output,¹ epilepsy and its allied conditions form for the moment one of the "growing points" of neurological research; and this is matter for congratulation, since nothing but good can emerge from a vigorous and concentrated attack on the difficulties presented by an age-old yet ever new and common morbid clinical state.

I am, however, personally under no delusion as to the complexities and intricacies of the subject, and I cannot conceive of any advantage in minimizing them; on the contrary, now is the time to remind ourselves of Curnow's dictum, often quoted by Hughlings Jackson: "The tendency to appear exact by disregarding the complexity of the factors is the old failing in our medical history." Were epilepsy not so involved a problem it would long ago have received at least a working interpretation, but the truth is the condition defeats us on many grounds; its etiology is heterogeneous, its semeiology indeterminate, its pathology dubious, its pathogenesis conjectural, and its therapeutics empirical. For the purposes of this discussion I can only single out one or two aspects to which attention may be usefully directed; they have been chosen intentionally as in my view best calculated to indicate the range of the problems connected with the epilepsies.

I. THE CLINICAL FEATURES OF EPILEPTIC STATES.

At the outset the student must face difficulties arising from the *clinical polymorphism* of epileptic phenomena. We all believe we can diagnose an epileptic fit with readiness when we see one, and it is true enough that a major attack is, fortunately, like nothing else in the whole field of symptomatology; but the epileptic state is not by any means confined to major seizures. Motor, sensory, psychical, and visceral variants exist; and in not a few instances it is hard to say whether the symptoms observed belong to the epileptic group. To separate recurring involuntary movements of a tic-like or spasm-like character from local epilepsy is often difficult, and sometimes impossible. Moreover, during a major attack the observer will frequently notice movements which will not bear interpretation as essentially convulsive; they are co-ordinated, and have the appearance of being purposive, involuntary though they must be. At the other end of the scale are fits distinguished by relative motionlessness—an akinetic or inhibitory variety that refuses to be separated from the main class by any physiological or clinical criterion, unless, indeed, our semeiological scheme is to be convicted of arbitrariness. But apart from the seizure itself we have to consider pre-seizure and post-seizure disorders, which are integral parts of the whole, however grossly they differ from the symptomatology of the attack of convulsive movement. It will therefore aid us in taking a large, a scientific, view of the question if we remember the following points.

1. The epileptic aura is a *sensation*, and thus a symptom belonging to the *psychical* series. It is something in

consciousness. The wonderful historical study of the aura conducted long ago by Herpin² emphasizes the hallucinatory nature of this familiar phenomenon, for a description of which we are dependent on the patient himself. But I stress the fact that it need not be either distorted or caricatured; any sense may be involved, simply or elaborately; I mention in passing, as worthy of special remark, hallucinations of the muscular sense. Further, it may be more intricate still, consisting of a series of panoramic pictures, of forgotten memories recalled, of incidents apparently *re-lived*; and in this respect its relation to some of the phenomena we call "hysterical" is immediately apparent.

2. Interruption of the stream of *consciousness* may, or may not, accompany epileptic manifestations. Absolute unconsciousness may *supervene* without aura, fulminantly; but many fits, especially perhaps those of a local or unilateral, and some of a *visceral* kind, run their course with conservation of consciousness, enabling the subject to observe and comment on his own experiences.

3. The *motor content* of the fit consists usually of violent convulsive movements, not to be likened to any of everyday life—powerful repetitive muscular contractions at their wildest and most intense. Yet every clinician knows it is not always thus. Mild flickers of finger and thumb, quivering facial contractions round the mouth, repeated local twitching of a solitary limb-segment, something little more than the striking of an attitude in momentary spasm, can constitute the whole of the motor performance. We must note, again, the not infrequent occurrence of co-ordinated, quasi-purposive movement; and also absence of movement.

4. The *sensory elements* in an epileptic seizure, aura apart, sometimes make up practically the whole of it; further, in numerous instances the aura develops as though to herald the onset of the usual attack, without the latter making its appearance.

5. The *visceral components*, neurosympathetic in origin, are just as significant as the somatic, and just as frequent. Here are included circulatory, respiratory, vasomotor, vesical, rectal, renal, vagal, salivary, sudatory, pupillary, and other phenomena.

6. As for *post-seizure manifestations*, they embrace the most varied sets of opposites, from sleep to fugues, from stupor to aggressiveness and combativeness, from speedy or immediate recovery to prolonged headache, physical fatigue, and general malaise. In some of these respects they are completely indistinguishable from what occurs in the condition we still, for want of a better name, call hysteria.

With all this the clinician is familiar. My sole purpose in this sketch has been to underline that *heterogeneousness of manifestation*, ranging from the psychical to the visceral, which has to be included in any survey of epilepsy and has to be accounted for by any theory of pathogenesis.

II. THE INHERITANCE OF EPILEPSY.

Epileptic phenomena being demonstrably of extreme diversity, it becomes a question whether inheritance can be of any particular import in connexion therewith. Twenty-one years of practice and hospital experience have long since convinced me that the hereditary factor is persistently overrated, and the personal or constitutional factor underrated. The number of cases in which no hereditary element can be traced far exceeds that in which such a factor seems to be responsible. It is time protest should be unitedly voiced by neurologists against the portentous ascription of a sinister prognosis to every case of epilepsy because of a supposed inheritance and consequent incurability. Gowers³ found a family history of epilepsy in about 27 per cent. of his own and other collected cases (2,400); and Russell Brain⁴ in 28 per cent. of 200. But the percentage noted by other observers has been less still (in Spratling's series of 1,070 cases parental epilepsy existed in 16 per cent.; Davenport and Weeks,⁵ 19; Binswanger,⁶ 11). Even if we assume heredity for, say, 20 per cent., this means that in no less than 80 per cent. that factor is altogether wanting.

The matter can be approached from another angle. Were inheritance really a determinant, then in a large out-patient experience I should find myself treating many cases

*Made in opening a discussion in the Section of Neurology and Psychological Medicine at the Annual Meeting of the British Medical Association, Manchester, 1928.

of the condition among siblings. As a fact, this is distinctly rare. At the present time the sole familial pair or group is that of two epileptic sisters, whose family history is negative, and who are identical or homologous twins⁷—a striking example of the significance of the personal or constitutional component. Myerson⁸ cites information obtained (1925) from the superintendent of the Massachusetts State Hospital for Epileptics to the effect that among 1,500 inmates only four families of 11 persons in all were to be found. At the same institution Thom⁹ ascertained that 138 marriages of epileptics resulted in 553 offspring, among whom a history of epilepsy was obtainable in only 10 cases (1.8 per cent.). The most recent figures are those of Pilcz,¹⁰ who traced 161 children of 144 epileptic mothers; 78 died under one year, and of the others five were imbecile, three criminal, four psychopathic, two psychotic, and only three epileptic. It is, I submit, impossible to draw any other conclusion from these statistics than that direct heredity is of little moment in comparison with other inducing factors.

III. THE EXPERIMENTAL PRODUCTION OF CONVULSIONS.

The first point of importance that emerges from a great number of researches, old and new, on the experimental production of epileptic fits is that normal animals can be made to develop convulsions with ease and certainty. No inheritance is requisite, nor does this consideration, I venture to say, ever cross the mind of the experimentalist. Whatever the provenance of his laboratory animals, they are all potentially capable of developing epilepsy.

Study of methods reveals two possibilities; or, rather, two main technical procedures for the production of the convulsive state, and these may be classed as (1) neural, and (2) humoral, respectively.

It is of some relevance, I consider, to note at once that these may act separately or interrelatedly. This has been well demonstrated by Dandy and Elman,¹¹ who, to begin with, provoked convulsions in a normal cat by simple oral administration of absinthe in sufficient quantity. They next made cerebral (pre-Rolandic) lesions in the animals employed by either cortical or subcortical extirpation or the introduction of a foreign body subcortically (all the lesions were unilateral). Within from one to five months such animals became much more sensitive to the effect of absinthe, one-third to one-seventh of the original dose now resulting in fits. The addition of a pathological cerebral factor increased the facility, and aggravated the degree, of action of the convulsant; in other words, cerebral injury lowered the convulsion threshold. Experimental provocation of fits in the human subject also has in recent years been given much attention, in regard to both epileptic patients and normal persons, and a considerable gain to knowledge has thereby accrued.

(1) Taking the humoral factor first, we may define it comprehensively as that component in the production of experimental (both laboratory and human) epilepsy which consists in alteration of body fluids and biochemical constituents either mechanically or in connexion with exogenous and endogenous noxae. The table below is based on those published by Brock¹² and by Lennox and Cobb¹, with some modification.

TABLE I.—*Humoral Factors.*

Mechanical.

1. Increased intracranial pressure.
2. Increased permeability of tissues.
3. Alteration of water balance; oedema.
4. Pressure on or ligation of carotids and vertebrals.

Exogenous.

1. Convulsants (alcohol, lead, acid fuchsin, camphor, absinthe, thujone, cocaine, picrotoxin, etc.).
2. Insulin hypoglycaemia.
3. Foreign proteins.
4. Anoxaemia (variously induced).
5. Alteration of acid-base equilibrium; alkalosis (variously induced).

Endogenous.

1. Disorder of endocrines (parathyroid, thyroid, pituitary, ovary, etc.).

Without examining in any detail the "convulsigenic" significance of these various factors, I shall for the purpose of this discussion content myself by setting forth certain

general conclusions recently arrived at by Lennox and Cobb.¹³

1. Acidosis tends to inhibit, and alkalosis to augment, seizures.
2. An increased tension of oxygen in the tissues tends to inhibit and a decreased tension to augment seizures.
3. Oedema of the brain tends to increase, and dehydration to diminish, seizures.

Although these generalities merit attention the modes of action of the various factors have not been satisfactorily determined, and the separateness of their functions is not established. In other words, we are not in a position to say that the mechanical and the toxic components are always in essence contrasted or in their effect mutually and invariably independent. And not a few anomalies and difficulties remain for investigation. In order to produce opacity of cerebral arteries for radiographic purposes Moniz and Lima¹⁴ have injected harmless solutions of sodium iodide into the internal carotid, transiently ligatured; and in "a large number of cases" this has been followed by severe and rapidly developed Jacksonian fits, involving the body and limbs *on the same side as the injection and ligation*. At present no satisfactory explanation of this apparent paradox is forthcoming. The experimentation of Temple Fay¹⁵ appears to show that the mechanical factor (pressure, increased fluid, defective fluid absorption) is at least as significant in numerous instances as any of a chemical or biochemical nature (see further below).

(2) As for the second (neural) mechanism, it has been known for years that direct excitation of neural centres can result in the development of unilateral or generalized epileptic attacks. When in these days so much attention is being directed to humoral factors it is essential not to underestimate the importance of the other.

IV. THE OCCURRENCE OF CONVULSIONS IN PATHOLOGICAL STATES.

If the deductions of the experimentalist in respect of epileptic determinants are to prove of value they must be capable of substantiation from the clinical side—that is to say, there must be evidence that these, or some or other of these, play a part in the production of epileptic syndromes under pathological circumstances. What, then, are the neural and humoral (or humori-vascular) conditions encountered in clinical medicine that are likely to exhibit epilepsy as one of their manifestations?

Taking neural (cerebral) conditions first, we recognize in what a large number of diverse organic cerebral states epilepsy in one or other form may make its appearance, and these can be tabulated as follows.

TABLE II.—*Organic Cerebral States associated with Epilepsy.*

1. Cerebral tumours, of diverse kinds and sites.
2. Cerebral infections and toxi-infections (varieties of encephalitis, syphilis, tuberculosis, abscess; toxins of many specific fevers and exanthems, etc.) (Some of these are classifiable as humoral states also.)
3. Cerebral toxi-degenerations or sclerosis (disseminated sclerosis, tuberous sclerosis, Pick's cortical atrophy, Alzheimer's disease, diffuse sclerosis, Westphal's pseudo-sclerosis, etc.).
4. Cerebral trauma.
5. Certain varieties of congenital and heredo-familial disease.

In some organic cerebral affections fits are few and far between, and in the case of fevers and exanthems they often occur solely at the onset. In other instances they are frequent and persistent. If we seek a common factor for the development of convulsive attacks in these uncommonly diverse states it will be difficult to find, no doubt; but this question we shall for the moment postpone.

Passing now to humoral and vascular cerebral pathological conditions, we can classify them in the following way.

TABLE III.—*Humoral and Vascular Cerebral States associated with Epilepsy.*

1. Disease of blood vessels (thrombosis, embolism, haemorrhage, arterio-sclerosis, endarteritis obliterans, aneurysm, etc.).
2. Disorder of cerebral circulation (heart-block, anaemia, asphyxia of various origins, etc.).
3. Disorder of cerebro-spinal fluid regulation (hydrocephalus, external ("ocult") hydrocephalus, alterations in fluid pressure, etc.).

Neither of these two latter tables makes any pretence at completeness; nor, for that matter, have I by any means exhausted the catalogue of clinical conditions, temporary or otherwise, in the course of which epilepsy may reveal itself, for I might mention its occurrence during the administration of anaesthetics, in association with narcolepsy, under the influence of emotion, in alternation with certain psychical conditions, such as attacks of bad temper, and so forth.

V. THE MECHANISM OF PRODUCTION OF THE EPILEPTIC SEIZURE.

If now we put together the information garnered from the two preceding sections we shall probably not be far wrong in concluding that it can be summarized as implying the existence of three or four determinants, at most, for the production of epileptic phenomena.

An epileptic disorder is a disorder of a neuronie mechanism, without question; not of any specific mechanism, but simply of the individual's neural network in one or other of its parts. True, the major portion being motor, the disorder exteriorizes itself via the pyramidal system, but it commences anywhere, so to speak; witness the polymorphic auras.

The crux of the pathogenic problem remains; is this neuronie disturbance (irritative or release, it matters not) initiated in the neuronie mechanism itself, or is the latter influenced by humoral or vascular change? Does one accompany the other, or precede? At the present time no direct answer can be given to the question. What evidence has accumulated is indirect and collateral, and may be set down as follows.

1. Mechanical Determinant.

It has for many years been suspected, and often argued, that some connexion exists between the epileptic state and increase in amount and pressure of cerebro-spinal fluid; and the view has often been expressed that a relation can be established between the occurrence of the latter and failure in fluid absorption owing to arachnitis or other pathological local condition. According to Dandy,¹⁶ Fay,¹⁵ Foerster,¹⁷ and many more, x-ray photographs of the head in a large number of idiopathic epileptics (fluid being replaced by air for radiographical purposes) have shown the existence of enlarged subarachnoid spaces, more especially over the vertex and fronto-parietal regions; while, pathologically, subarachnoid collections of fluid, local oedemas, and such-like conditions have often been seen. These and other data suggest the possibility of external hydrocephalus—the "occult hydrocephalus" of Reynolds,¹⁸ who has found the condition in epileptic subjects at operation—which arises from defective fluid absorption and leads to pressure, and so to fits. Fay¹⁵ has recently added to our knowledge of this particular mechanism by his examination of the Pacchionian bodies in chronic epileptic brains. They are the portals through which the fluid in large part escapes, and it is therefore instructive to note that his colleague Dr. Winkelman has found various stages of sclerosis, calcification, atrophy, failure of development, and other abnormalities in these organs in epileptic subjects. Such changes follow infections, haemorrhages, local injuries, toxicoses, etc. They suffice to account for delay in fluid absorption, fluid accumulation, back pressure, and consequent supracortical oedema. Yet the problem still remains—to explain the occurrence of fits, given such pathological circumstances.

2. Vascular Determinant.

Many direct observations have been made of the vascular state of a patient on the threshold of an epileptic attack and of the exposed brain surface throughout an attack, and the consensus of opinion holds that cerebral and general pallor or anaemia is more or less constantly visible to the eye just as the seizure commences. This circulatory derangement or disorder is confirmed by the frequent observation of failure of the peripheral pulse at the outset, and the rarer one of constriction of retinal arteries at the same moment. These are old and long-established facts, which we owe to clinicians of other generations (Moxon,¹⁹ Troussseau,²⁰ Hughlings Jackson, Hilton Fagge,²¹ Broad-

bent²²), whose work should not be overlooked by newcomers in the epileptic field. The analogy of these phenomena with what obtains in syncope or fainting attacks is unmistakable, and was minutely studied by Russell²³ a number of years ago. In 1903 the circulatory steps were thus epitomized by Francis Hare²⁴:

1. Vaso-constriction, causing rapid rise in general blood pressure.
2. Cardiac inhibition (vagal), to counteract the effects of the above, causing sudden fall in general blood pressure.
3. Sudden cerebral anaemia, causing unconsciousness and tonic spasm.
4. Recommencement of the heart beat, causing rise of pressure, returning cerebral circulation, relaxation of tonic spasm, and clonic convulsions.
5. Re-establishment of blood pressure and cerebral circulation; cessation of convulsions; sleep recuperative of exhaustion and damage.

It is undoubtedly possible to harmonize the facts of clinical observation and of experimental epilepsy with this sketch of the phases in pathogenesis; but there are many obscurities.

What is the proximate cause of the sudden initial vascular constriction? And where does it start? To the former question no satisfactory answer is as yet forthcoming. We do not know whether the vasomotor change begins in the vasomotor centre of the bulb, or locally in the cerebral cortex, or distally, by way of reflex influence on the medulla or cortex. And even when we assume for vasoconstriction a pathogenic significance, I may point out that by itself transient cerebral anaemia is insufficient to determine the epileptic attack. Elsewhere²⁵ I have commented on and criticized the evidence in its favour, and have endeavoured to show how indeterminate the relationship is. Changes in cerebral circulation accompany the march of events in an epileptic fit, but whether the interconnexion is causal remains to be proved. Were cerebral anaemia the *per se* cause, then every deathbed, more or less, should be the scene of convulsions, which is, fortunately for all concerned, not the case. Nor can the occurrence of an integral epileptic phenomenon—the aura—be thus easily explained, for it is in numerous instances a precise psychical activation, a hallucination; and there is no reason to imagine that neural activity is accompanied by anaemia in the cerebrum of the normal subject. If, however, we assume it to come into consciousness before, cortical vaso-constriction, we still are faced with the enigma of how it arises itself, how it is related to the latter, and what is the state of affairs when it is not followed by convulsions. Is the existing condition then neural or vascular? I submit, further, that many local cerebral anaemias are unaccompanied by discharging phenomena (witness the flaccid palsies of embolism and thrombosis); and, on the other hand, that varieties of epileptic discharge occur in a way at once so limited and so continuous (some Jacksonian attacks, *epilepsia partialis continua*) that it seems impossible to conceive for them a vasomotor determinant such as is suggested for major fits.

In arguing thus I do not understate the importance of the vascular factor, which can properly be made responsible for some of the obvious manifestations of the seizure, and I do not hold entirely with my former teacher, the late Sir William Gowers,²⁶ when he stated dogmatically that "The vasomotor theory of epilepsy is alike unneeded, unproved, and inadequate. The phenomena indicate that there is a discharge of grey matter, and there is nothing to warrant us in going beyond the grey matter concerned in our search for the origin of the discharge." But it is my present considered opinion that proof of the initiation of the whole sequence of events by a causal vasomotor constriction is lacking, and that it is equally conceivable that a neural factor originates the vascular change.

3. Humoral Determinant.

The humoral or physicochemical determinant has, as we have seen, received an immense amount of attention in recent times, and some conclusions have been outlined above. Briefly, it must be considered as established that a number of conditions (hyperventilation, injections of various chemical substances, anoxaemia variously produced)

cause experimentally a marked fall of O_2 tension in the tissues and are calculated to give rise to convulsions (Argyll Campbell²⁷). And it is frequently stated that the "purpose" of convulsions is to counteract this O_2 deficiency. The relation of such deficiency to alkalosis and other non-neural components is sketched by Lennox and Cobb¹³ as follows:

"Contraction of cerebral vessels might lead to a decrease in the oxygen supply to the brain. Because of the anoxaemia there would be dilatation of capillaries with passage of fluid outward through their walls. One or more of these factors—decreased oxygen tension, oedema, increase in intracranial pressure—might lead to abnormalities in the activity of cortical neurones and a fit. Consequent on the convulsion there would be an accumulation of carbon dioxide and lactic acid in the tissues, which would initiate a reversible reaction. Accompanying the acidosis there would be an increase in the utilization of oxygen by the tissues, a passage of fluid into the capillaries, and a restoration of normal intracranial circulation."

The clinician is willing to believe that such a series of physicochemical interchanges accompanies a major epileptic seizure; but he notes that the description commences with a postulated "contraction of cerebral vessels," which, as I have just emphasized, itself remains unexplained, of uncertain origin, and of dubious application to various clinical symptoms which cannot be expelled from the epileptic category. Lennox and Cobb, it is satisfactory to observe, are aware of the hiatus, for they admit that "these processes are only contributory; they may be effective only in persons with a tendency to seizures, and with those having some pathology either in the brain, the body, or the emotional life."

4. Neuronic Determinant.

We are therefore, after this excursus into newer fields of investigation, brought back, not unexpectedly, to the position which for some is still unassailed (I do not say unassailable)—namely, that the essence of all epileptic semeiology, major or minor, pre-seizure or post-seizure, psychical or visceral, is neuronic derangement, and that this derangement is of neuronic derivation. Nervous tissue can be, and is often enough, influenced by mechanical, vascular, and humoral agencies; but the secret of epilepsy—the core of the problem—resides in the qualities of the neural mechanisms exhibiting discharge.

It is wise not in the least to undervalue the importance of recent acquisitions to knowledge, apropos vascular and humoral constituents and accompaniments of epileptic fits, but at the same time the following considerations must be given due weight.

1. The action of laboratory convulsants such as absinthe, which will produce fits in a normal animal, is much facilitated if the animal undergoes any cerebral injury.
2. Single humoral or vascular factors, by themselves, are often insufficient to produce fits; and even when combinations occur there may be no element of essential repetitiveness in the attacks thus generated.
3. Only a small percentage, as is well recognized, of cases with head injuries and with histories of infective and other conditions, leading presumably to the pathological states mentioned above, react with fits.
4. The development of epilepsy in some organic brain states (see especially No. 3 in Table II above) pleads in favour of a neural factor, seeing that no evidence of vascular or humoral anomalies is forthcoming in them.
5. A "susceptibility" or "tendency" can scarcely be ruled out in cases of direct heredity and of epilepsy in homologous twins, and it is at least as easy to express this in terms of neural instability as of extraneural abnormality.
6. The clinical occurrence of so-called reflex epilepsy strongly suggests a pure neural reaction.
7. Certain features of the epileptic state are clinically indistinguishable from hysterical manifestations; but for the latter, so far as I am aware, no attribution to mechanical, vascular, or humoral precipitants is thought of.
8. Certain phenomena of the epileptic fit (for example, inhibition and radiation) are much more readily explicable by neural than by any extraneural mechanism.

It is therefore impossible to find one single common factor for the totality of epileptic manifestations, unless the significance of some inherent nervous factor, some functional propensity, is allowed. However unsatisfactory the word "tendency" or "instability" may be, however useful in cloaking ignorance, I do not consider this a feeble conclusion; on the contrary, it is consonant with the facts of observation; the only trouble is that we cannot at present gauge "susceptibility" by reference to objective criteria. Nor, for that matter, can anyone estimate what is signified by susceptibility to infection or illness of any kind, unless by rather vague allusions to "immunity." But the question of immunity in epilepsy has never yet been seriously approached. For this reason I shall conclude by a brief note on the influence of intercurrent conditions on the epileptic state.

VI. THE INFLUENCE ON EPILEPTIC SEMEIOLOGY OF OTHER INTERCURRENT AFFECTIONS.

This perhaps deserves more attention than has hitherto been accorded it. The observation is as old as the days of Hippocrates, for that Father of Medicine stated it himself, that intermittent fever replaced or mitigated epilepsy, whence the adage "*Quartana epilepsiae vindex appellatur*." Lenoir²⁸ (1901) remarked that Esquirol had commented on the diminution or disappearance of epileptic attacks during "accidents fébriles," while Hammond²⁹ in the same year said that some authors, struck with the favourable influence exerted by infectious diseases over a pre-existing epilepsy, had advocated "the establishment of some of these different processes, such as malaria and erysipelas, to cure it." As long ago as 1882 it was noticed by Bourneville and Bonnaire³⁰ during an epidemic of measles in epileptics and idiots at Bicêtre that while the intercurrent affection lasted fits were much decreased in force and frequency, and similar observations were recorded by Séglas³¹ (at the Salpêtrière). Gowers³ quotes the case of a child ("with strong inheritance") who suffered from fits from 2 years of age till 7, when they ceased during an attack of typhoid fever and did not recur till the age of 27. He also states that scarlet fever provides an exception to this perhaps general rule. Aldren Turner,³² on the other hand, has seen a case in which a mild attack of scarlatina induced a period of amelioration lasting for several months; and another in which respite was obtained for a similar period after acute pneumonia. Bourneville,³³ once more, found during a typhoid epidemic among epileptic children (six cases out of seven) that the attacks were suspended while the pyrexia lasted, and that the improvement persisted for some time after recovery.

In my own experience measles and chicken-pox are the febrile illnesses which have sometimes effected a striking cessation of fits—nor has the improvement been limited solely by the duration of the intercurrent state.

So far as I have discovered no attempt has been made by Gowers or any of the other authors mentioned to explain the way in which this influence may be, theoretically, exerted. According to Francis Hare,²⁴ however, it is the pyrexia which is of significance, for it involves general vascular relaxation, and he argues that vaso-dilatation is similarly of value in cutting short attacks of both asthma and migraine. He quotes in this connexion Garrod, who noticed the effect of acute pyrexial gout on recurrent epilepsy.

"A gentleman had suffered from epilepsy from the age of 20 to 52; the fits were frequent, sometimes occurring as often as once a week. He then had distinct articular gout in one great toe, and afterwards experienced attacks of the same kind, from time to time, up to his death at the age of 72. From the first manifestation of decided gout there was an entire cessation of the epileptic convulsions."

It would be simple and proper enough to account for this particular ameliorating action on humoral lines; but in the case of the exanthems I scarcely feel that pyrexial vaso-dilatation covers the whole pathogenesis, just as I do not feel confident that the old observation that in children various acute diseases commence with convulsions, but in adults with rigors is to be elucidated by a theory of vaso-constriction, acting more precipitately in the former than in the latter.

Whatever the explanation, it may yet turn out to be of importance from a pathogenic and a therapeutic standpoint that epileptic manifestations are sometimes controlled by Nature's own methods.

REFERENCES.

- ¹ Pagniez, P.: *L'Epilepsie: conceptions actuelles sur sa pathogénie et son traitement*, Paris, 1929.
- ² Crouzon, O.: *Le syndrome épileptique*, Paris, 1929.
- ³ Lennox, W. G., and Cobb, S.: *Epilepsy*, London, 1928.
- ⁴ Muskens, L. J. J.: *Epilepsy*, London, 1928.
- ⁵ Herpin, Th.: *Du pronostic et du traitement curatif de l'épilepsie*, Paris, 1852.
- ⁶ Gowers, W. R.: *Epilepsy*, London, 1901.
- ⁷ Brain, W. R.: The inheritance of epilepsy, *Quart. Journ. of Med.*, 1926, xix, 299.
- ⁸ Davenport, C. B., and Weeks, D. F.: A first study of inheritance of epilepsy, *Journ. Nerv. and Ment. Dis.*, 1911, xxxviii, 641.
- ⁹ Binswanger, O.: Die Epilepsie, *Nothnagel's Spec. Path. u. Therap.*, Vienna, xii, 1899.
- ¹⁰ Wilson, Kinnier, and Wolfsohn, J. M.: Organic nervous disease in identical twins, *Arch. of Neurol. and Psychiat.*, 1929, xxi, 477.
- ¹¹ Myerson, A.: *The Inheritance of Mental Diseases*, Baltimore, 1925, p. 71.
- ¹² Thom, D. A.: A second note on the frequency of epilepsy in the offspring of epileptics, *Boston Med. and Surg. Journ.*, 1916, clxxv, 599.
- ¹³ Pilcz, A.: Die weiteren Lebensschicksale von Kindern welche während der Bestehens einer mütterlichen Geistes- oder Nervenkrankheit geboren worden sind, *Jahrb. f. Psychiat. u. Neurol.*, 1928, xlv, 153.
- ¹⁴ Dandy, W. E., and Elman, R.: Studies in experimental epilepsy, *Bull. Johns Hopkins Hosp.*, 1925, xxxvi, 40.
- ¹⁵ Brock, S.: The problem of the epilepsies, *Internat. Clinics*, 1928, iv, 179.
- ¹⁶ Lennox, W. G., and Cobb, S.: The relation of certain physicochemical processes to epileptiform seizures, *Amer. Journ. Psychiat.*, 1929, viii, 837.
- ¹⁷ Moniz, E., and Lima, A.: Accès épileptiques à aspect Jacksonien homolatéral, déterminés par l'injection d'iodure de sodium dans la carotide interne, *C. R. Soc. de Biologie*, 1928, xxviii, 1011.
- ¹⁸ Fay, Temple: Some factors in the mechanical theory of epilepsy, *Amer. Journ. Psychiat.*, 1929, viii, 783.
- ¹⁹ Dandy, W. E.: Ventricleography following the injection of air into the cerebral ventricles, *Annals of Surgery*, 1918, lxviii, 5.
- ²⁰ Foerster, O.: Encephalographische Erfahrungen, *Zeit. f. d. g. Neurol. u. Psychiat.*, 1925, xciv, 512.
- ²¹ Reynolds, C. E.: Meningitic epilepsy, *California State Journ. Med.*, August, 1921.
- ²² Moxon, W.: Croonian Lectures, *Lancet*, 1881, i, 649.
- ²³ Troussseau, A.: *Clinical Medicine*, New Syd. Soc., i, 1867.
- ²⁴ Fagge, Hilton: *Textbook of Medicine*, London, 1886, i, 750.
- ²⁵ Broadbent, W.: Croonian Lectures on the Pulse, *Lancet*, 1887, i, 710.
- ²⁶ Russell, A. E.: The pathology of epilepsy, *Proc. Roy. Soc. Med.*, Med. Sect., December, 1907, p. 71.
- ²⁷ Hare, Francis: Mechanism of the paroxysmal neuroses, *Australasian Med. Gaz.*, July, 1903.
- ²⁸ Wilson, Kinnier: *Modern Problems in Neurology*, London, 1928, Chap. I.
- ²⁹ Gowers, W. R.: *Diseases of the Nervous System*, London, 1893, ii, 755.
- ³⁰ Campbell, J. A.: Tissue oxygen-tension with special reference to tetany and convulsions, *Journ. of Physiol.*, 1925, ix, 347.
- ³¹ Lenoir, L.: *Thèse de Paris*, 1901.
- ³² Hammond, C. H.: *Medical Annual*, 1901, p. 241.
- ³³ Bourneville and Bonnaire: Relation d'une épidémie de rougeole observée à l'hospice de Bicêtre dans la section des enfants épileptiques et idiots, *Prog. Méd.*, 1892, x, 663.
- ³⁴ Siglas, J.: De l'influence des maladies intercurrentes sur la marche de l'épilepsie, *Thèse de Paris*, 1880.
- ³⁵ Turner, W. A.: *Epilepsy*, London, 1907, p. 50.
- ³⁶ Bourneville: Epidémie de fièvre typhoïde à Bicêtre, *Prog. Méd.*, 1899, xxvii, 147.

TUBERCULOSIS OF THE LARYNX.*

BY

WALTER G. HOWARTH, M.B., B.Ch., F.R.C.S.,

SURGEON TO THE THROAT DEPARTMENT, ST. THOMAS'S HOSPITAL.

WITH so large a subject as this it is not easy to know from what side to approach it, and I think that it will probably best serve the purposes of the discussion which is to follow if I make a few remarks from a very general point of view, and let those that follow amplify or criticize some aspect that may particularly appeal to them.

I am afraid that, in spite of the fact that for the past twenty years I have tried to study this disease at St. Thomas's Hospital, the Victoria Park Chest Hospital, and elsewhere, I have really nothing to say that is new, and the more cases that one sees the more is one inclined to wonder at the protean manifestations of this disease as far as the larynx is concerned. If one examines carefully the larynx in patients who are suffering from pulmonary tuberculosis, one may say that somewhere about 10 per cent. show abnormality in the larynx. I do not wish to imply that a definitely recognizable tuberculous lesion is present in all these cases, but that the larynx is not quite normal; it may be that the mucosa is more anaemic than is usual, or there may be a generalized erythema, or possibly some fullness in the interarytenoid region. Do such departures from the normal occurring in a phthisical subject possess more than usual significance, and is there such a thing as

a pretuberculous stage of the disease? I believe that many people consider that there is such a stage. Personally I am disinclined to admit this designation, and I would prefer to regard the local anaemia as a part of a more general one, and the catarrhal laryngitis as a not unusual accompaniment of the chronic cough, while the passage of sputum over the interarytenoid fold is surely sufficient to cause a little local hypertrophy without this being of necessity due to a definite tuberculous infection. At the same time, it is naturally advisable to keep a careful watch on such local conditions as they arise, and to make certain that they yield to the appropriate treatment.

What proportion of cases of proved phthisis exhibit laryngeal symptoms that are definitely tuberculous? The answer to this question is not very easy, as statistics are not by any means unanimous on the subject, and, like all statistics, they may be essentially fallacious, as often the information is not given as to what particular class of case is being dealt with. Thus at a large sanatorium (Midhurst) 8 per cent. show definite laryngeal disease, whilst at an institution such as the Victoria Park Chest Hospital, where the cases are year by year more carefully selected, the proportion may fall as low as 4 per cent. On the other hand, some recent American statistics from a State sanatorium mention 25 per cent. of laryngeal cases. However much statistics vary, it is, I believe, true to say that laryngeal complications of phthisis are certainly less common than they were even twenty-five years ago. Then, the average statistics of many skilled observers in Germany and elsewhere showed considerably more than 30 per cent. This improvement is, I think, undoubtedly due to the fact that the chest disease is recognized earlier and is usually treated most efficiently from the beginning of the disease.

It may be noted that I am assuming that tuberculosis of the larynx is secondary to pulmonary disease. This is, I think, almost universally agreed for the vast majority of cases, even though no definite lesion can be demonstrated in the lungs. There is to my mind, however, one lesion that is difficult to explain on this assumption, and that is the tiny nodule which may occur on the edge of a vocal cord in an elderly person, and which may give rise to the suspicion of cancer, but which the microscope proves to be tuberculous in spite of the fact that there was no trace of tuberculosis to be found elsewhere. I can recall at least two cases in which (with the concurrence of distinguished colleagues) I removed the supposedly cancerous cord with completely successful results, and no further manifestation of tuberculosis over a period of twelve years and seven years respectively.

When the larynx becomes affected in a case of pulmonary tuberculosis, how does the infection occur, and what parts are most commonly involved? In the large majority of cases secretion coughed up from the lungs produces a surface infection of the laryngeal mucosa, and the infecting bacilli can undoubtedly enter apparently intact epithelium, and still more so epithelium damaged by previous disease. When one considers the amount of sticky sputum that so often hangs about the interarytenoid region and the posterior ends of the vocal cords, it is perhaps surprising that infection does not more often take place. Certainly this stagnation of secretion in the region of the posterior commissure can be regarded as a very potent source of infection. In this connexion it is interesting to note that singers and some other voice users who do not tolerate any accumulation of mucus in the larynx are said to be less susceptible to laryngeal complications. This may be so, but there are, of course, other explanations.

When we allow the possibility of entry of bacilli through intact or abraded epithelium, or through the ducts of racemose glands, it seems perhaps unnecessary to consider the blood stream as the primary vehicle for the submucous deposit, and it may, indeed, be doubted whether infection ever occurs this way. At the same time, when once tuberculous disease has been established, spread to neighbouring parts of the larynx may undoubtedly occur by extension through the lymphatics.

It is, I think, agreed that in the majority of cases the posterior part of the larynx is most commonly affected, and it is here that we are accustomed to look for the earliest signs of disease. We are all familiar with the firm nodular

* Read in opening a discussion at a joint meeting of the Sections of Oto-Rhino-Laryngology and Tuberculosis at the Annual Meeting of the British Medical Association, Manchester, 1928.