13. So far as the specific prophylaxis of puerperal fever is concerned, it has been shown that while the exotoxins of the streptococci causing puerperal sepsis can be neutralized by any monovalent streptococcal antitoxin, the destruction of the organisms is only likely to be secured by the production of a type (serological) specific antibacterial serum.

PREVENTION.

The thesis is submitted, therefore, that in so far as the prevention of maternal mortality is concerned, and within the limits of our present knowledge, a significant reduction in maternal mortality would be immediately attained by (a) an extension of adequate ante-natal services, and (b) the institution of a new midwife-doctor combination as above defined (paragraph 11).

DIAGNOSIS.

With reference to the bacteriological diagnosis of puerperal fever, confusion has resulted from the wealth of bacteriological data relating to the complexity of the flora of the genital tract (1928). It is admitted that there is a great variety of organisms in the genital tract at the time of the puerperium, and that puerperal sepsis is not necessarily associated with streptococcal infection. The point that matters, however, is that the serious forms of puerperal sepsis, and certainly some 90 per cent. of the deaths from sepsis, are due to infection with haemolytic streptococci. If serious morbidity and death from puerperal sepsis is to be countered, therefore, the essential is to concentrate on the haemolytic streptococcus.

TREATMENT.

Prevention and treatment cannot properly be considered separately, since adequate ante-natal treatment prevents maternal deaths in general, and since the early treatment of puerperal sepsis involves isolation and the prevention of further cases of sepsis by contagion.

In so far as treatment may lead to the prevention of serious morbidity and mortality from sepsis, the first essential is early recognition of the infection, and to this end puerperal pyrexia has been made notifiable in England, and is in process of being made notifiable in Scotland. Up to the present time the vast mass of cases of puerperal fever receive comprehensive treatment only when the disease is so advanced that the prospect of recovery is remote.

In this connexion the advantages of the notification of puerperal pyrexia can be seen from the following chart, wherein the figures of pyrexia notifications have been analysed and those attributable to sepsis added to the puerperal fever notifications.

The chart shows that during the four years 1925-28, during which pyrexia has been notifiable, the elicited incidence of puerperal fever has increased from an average of 3 per 1,000 births prior to 1925 to 12 per 1,000 births in 1928. It will generally be conceded that such increasingly early diagnosis of puerperal sepsis is bound to have a profound effect on the efficacy of treatment. From the clinical point of view the valuable results obtained by Remington Hobbs (1927) with reference to glycerine drainage of the uterus are noted.

LITERATURE.

- Burt-White, H.: *British Medical Journal*, 1928, i, 974.
 Burt-White, H., and Armstrong, R.: *Ibid.*, 1929, i, 532.
 Hobbs, A. Remington: *Ibid.*, 1927, 1223; and *Medical World*, 1927.
Maternal Mortality in Aberdeen, 1928, H.M. Stationery Office.

A MODIFICATION OF LAGRANGE'S OPERATION.*

BY

W. H. KIEP, M.B., CH.B. GLAS.,

HONORARY OPHTHALMIC SURGEON, ROYAL EYE AND EAR HOSPITAL,
BRADFORD.

FIFTEEN years ago, at the historical meeting of the British Medical Association in Aberdeen, it was my privilege to see a very interesting specimen, shown by Mr. Hill Griffith, of an eye that had been excised on account of late infection following trephining by Elliot's method. The details of that case are to be found in the *Journal* of the Association (1914, vol. ii, p. 389). They left a rather profound impression on my mind. Here was a perfectly good operation—and, after all, no fistulization operation has found greater favour with the ophthalmological world than has Elliot's—performed by a perfectly good operator, and yet it was followed by late infection that necessitated removal of the eyeball. In those days I was not in a position to do any operations for glaucoma on the living human subject, but when, a few years later, my opportunity occurred, it struck me that, after all, it was not so surprising that late infection should occur in cases of corneo-scleral trephining, because the membranous covering of the fistula must essentially be rather tenuous and flimsy if that fistula be situated at the corneo-scleral margin. Obviously, therefore, a fistulization operation, to be reasonably safeguarded against late infection, must provide for a membranous covering in the region of the fistula sufficiently thick and impermeable by micro-organisms or their toxins to prevent the occurrence of an iritis, or at any rate of a septic irido-cyclitis. Now I should like to state at the outset that it is not my intention to criticize Elliot's operation for glaucoma, nor yet to describe a new operation for that disease. Most ophthalmologists are conversant with all the operative procedures for glaucoma (and their number is legion), and have had successful results with any or all of them. The methods of Young, Holth, Herbert, Heine, Bentzen, and others have been described elsewhere, and have found, no doubt, their staunch adherents. I merely desire to describe a modification of an operation advocated by one of the greatest protagonists of sclerectomy—namely, Lagrange—which seems to me easier to perform than the operation he describes, and to yield results quite as successful as those which he claims.

Every ophthalmologist knows well the procedure of Lagrange's operation for glaucoma. Those who were present at the Oxford Ophthalmological Congress of 1927 listened with rapt attention to his address and never doubted his statement that "we have never seen late infection." The only drawback to his operation is the technical difficulty of its performance. Whitehead of Leeds alludes to this in his contribution to the discussion on the results of operative treatment of glaucoma. Besides, it will surely be conceded that the size of the scleral wound which Lagrange employed is out of all proportion to the needs of the case. As many other operators have shown, a sufficient piece of sclera can be excised from the lip of a wound made with a small keratome and the trauma to the scleral envelope correspondingly diminished.

The modification of the operation which I beg to submit is as follows:

Conjunctiva and subconjunctival tissue are grasped with forceps at a point 6 or 7 mm. above the limbus, and divided with straight-pointed scissors. The capsule of Tenon, if prolonged as far forward as this, should be divided likewise in order to secure the thickest possible covering for the fistulous track. This composite sheet of tissue, composed of the three layers named, is then cut with the scissors by means of a curved incision running concentrically with the limbus (more or less as in Elliot's operation), the total length of this incision being about 1.5 or 2 cm. It is then reflected downwards in the direction of the limbus by easy dissection until we get to within 1.5 or 2 mm. distance of the corneo-scleral margin, when the scissors are laid down. The flap should not be grasped by forceps, but simply kept out of the way by

* A paper read in the Section of Ophthalmology at the Annual Meeting of the British Medical Association, Manchester, 1929.

means of moist cotton-wool swabs. A Took's corneal-splitting knife should then be taken, and with it a groove cut in between the layers of the scleral substance itself at a distance of 2 to 2.5 mm. from the limbus. It is neither necessary nor advisable to cut deeply with this splitter, but merely enough to make a small furrow running parallel with the limbus (corresponding to that between finger and a closely pared nail). Into this furrow a keratome is inserted and pushed downwards into the anterior chamber in much the same way as Heine does in his operation of cyclo-dialysis, except that the incision should be about 3 or even 4 mm. from end to end. The keratome should then be withdrawn as rapidly as possible. If this is done with sufficient speed very little aqueous is lost. The anterior lip of the scleral wound can now easily be gripped with Elliot's disc forceps and a small piece excised (like the paring of a finger-nail) by means of de Wecker's scissors. By this time very probably a small portion of the iris, close to its root, has prolapsed into the wound, and this is excised, a strictly peripheral iridectomy being aimed at. If the operator has been successful in his efforts thus far he will have lost little aqueous and still have an uncollapsed anterior chamber. He should then gently stroke the conjunctival flap upwards, and by gently pressing on the corneal surface of the limbus in the region of the iridectomy with a spatula he can cause the whole iris to resume its normal relationships, excepting only for the peripheral iridectomy. The conjunctival flap is then secured with one stitch. Should the iris not have prolapsed of its own accord, then its periphery should be gently withdrawn with iris forceps and a pinhole iridectomy performed. In this case more aqueous will undoubtedly be lost, but it should still be possible to complete the operation without causing complete collapse of the anterior chamber. If the iridectomy is a purely peripheral one there should be no necessity to poke an iris retractor into the wound, with consequent increase in the loss of aqueous, as the iris can practically always be caused to resume its proper position in the way indicated above—namely, by stroking the *surface* of the limbus and adjoining cornea with a spatula in the region of the iridectomy.

I am not so foolish as to claim that this operation is unattended by drawbacks. Chief amongst these is that, seeing the episcleral tissues are cut right down to the substance of the sclera itself and then stripped off this membrane, episcleral vessels, sometimes of relatively considerable size, are almost certain to be severed, and may cause troublesome bleeding. They can, however, be picked up with fine (toothed) conjunctival forceps and the haemorrhage stopped by torsion of the proximal end of the vessel.

In other respects, I have found this operation no more difficult and not more frequently attended by complications in the performance than is Elliot's, and it is less "tricky" and demands less skill in its execution than does Lagrange's. The advantages I claim for it are, roughly, as follows.

1. It is possible to do this operation with the minimum disturbance of the intraocular structures and the minimum risk of intraocular haemorrhage. Cases of acute or congestive glaucoma with shallow anterior chamber should be treated beforehand with myotics and Duke-Elder's method of administering hypertonic saline intravenously. The operation can then, say a week later, be performed as on an uninfamed eye.

2. In a large proportion of cases it is possible to do the operation without causing the anterior chamber to collapse completely. The valve-like nature of the fistula (as opposed to the rigid circular foramen obtained by trephining) is of great assistance in this respect. The advantages of maintaining a sufficient depth of aqueous fluid between the cornea on the one side and the iris and lens on the other need not be stressed here, because they are sufficiently obvious to any operator of experience. But I may say that restoration of the anterior chamber to its normal depth follows, on the average, in from forty-eight to seventy-two hours.

3. The thickness of the membranous covering of the fistula is a more efficient safeguard against late infection than it is possible to secure in cases of corneo-scleral trephining, because in the latter operation little more than conjunctiva *per se* covers the fistula, whereas, in the operation I have described, the fistulous track is covered by conjunctiva, subconjunctival tissue, and Tenon's capsule. The fistulous orifice, moreover, is situated not at the corneo-scleral margin, but at a distance of 2 mm. or even more from it. That the protection from late infection is efficient, I have, in my own mind, little doubt. One of my cases, with very free filtration, had the misfortune

to develop an attack of acute diplo-bacillary conjunctivitis about six months after operation. This attack passed off without being followed by any sign of iris irritation whatsoever (as observed with the slit-lamp, etc.).

4. Should, for any reason, a fibrosis ensue on this operation (as indeed may follow on any form of sclerectomy), a repetition of the operation may be performed without detriment. I have performed the operation successfully, and without difficulty, in a case in which a wide iridectomy had already been done for glaucoma, with insufficient results in the long run. A larger piece of sclera will, of course, have to be excised, but the previous history of the case will guide one in this respect.

In conclusion, I may perhaps be permitted to refer to a case in which this operation, in conjunction with other methods, was followed by truly gratifying results.

A married woman, aged 56, was seen by me in consultation with her medical attendant on the afternoon of March 14th, 1929. The presence of acute glaucoma in both eyes had already been diagnosed by her doctor, and the vision in the better eye (the left one) was no better than perception of hand movements, whilst that in the other eye was bare perception of light. She was admitted to the Eye and Ear Hospital, Bradford, a few hours later, and drops containing eserine 1 per cent., pilocarpine 2 per cent., and dionine 5 per cent. were instilled half-hourly, and, some five hours after I first saw her, 30 c.cm. of 30 per cent. aqueous solution of sodium chloride were administered intravenously according to the method of Duke-Elder. Next day the pupils were pin-point, and the patient was able to distinguish people at her bedside with her better eye, although the vision of the worse one (the right) was barely improved. Three days later the fields were, with a 5 mm. test object, as charted, and the vision of the right eye had improved to 6/60, and that of the left eye to 6/9. On March 21st the operation I have described was performed on the left eye, and on March 29th the patient was discharged from hospital, as she wanted to get home before her right eye was operated on. On April 17th she was readmitted to hospital, and the next day the right eye was operated on in the manner I have described. On May 11th the fields were, with a 5 mm. test object, as charted, and

V.R. with -4c -0.5 ↓90° equalled 6/18;

V.L. with -0.5 ↓90° equalled 6/6.

ACUTE INTESTINAL OBSTRUCTION.

BY

H. W. L. MOLESWORTH, F.R.C.S.,

ASSISTANT SURGEON, ROYAL VICTORIA HOSPITAL, FOLKESTONE.

In presenting for publication a review of 130 cases of intestinal obstruction which have been operated on by me, it is fitting that I should acknowledge the courteously given permission to make use of cases which I operated on whilst acting as surgical registrar to Sir Hugh Rigby, Mr. James Sherren, Mr. Russell Howard, Mr. Hugh Lett, Mr. A. J. Walton, Mr. Robert Milne, Mr. H. S. Souttar, Mr. E. C. Lindsay, Mr. W. S. Perrin, and Mr. G. E. Neligan, all of the surgical staff of the London Hospital.

Surgical lectures and papers on the subject frequently end on a wailing note, as though the summit of knowledge and of technique had been achieved. It is suggested that the existing high mortality is all due to late diagnosis, and that the technique of treatment had reached finality. With this generalized conclusion we venture to disagree. As surgeons it is absurd for us to ask that all our work should be easy child's play. When we are still ignorant of so important a subject as the nature of toxæmia in obstruction it is presumption to assume that the technique of treatment is final. No one who has seen much of intestinal obstruction can fail to be impressed with the fact that some of the most desperate cases survive whilst some apparently trivial cases die. In the former it would seem that by a happy chance appropriate measures were taken, and in the latter that over-confidence led to the omission of some such measure as intestinal drainage, which may be called for as an afterthought in less timely circumstances than at the primary intervention. The cases reviewed are consecutive, and it has not been the practice to refuse operation to any patient who was not actually moribund. No patient who was refused operation lived more than two hours.

It is apparent that the comparatively low mortality in external hernia may be justly attributed to the ease with