The method of formation would appear to be as follows. The fine spicules of asbestos cause, by mechanical action on the bronchioles and alveoli, either minute extravasations of whole blood, or serous exudates, which envelop them. Solution of any soluble fraction of asbestos takes place. The total amount of asbestos that is soluble must be extremely small, as proved by the x-ray pattern of the curious bodies, but in the case of chrysotile workers some solution is suggested by the free iron reaction. We must remember, however, that the Prussian blue reaction may be due to the iron of haemoglobin. Any surface in contact with a colloidal solution may act as an adsorbent, and in the present case the fine spicules must be considered to do so. Interaction between the soluble fraction of chrysotile and plasma proteins takes place, syneuresis occurs, and, with the loss of water, the adsorption is rendered irreversible. The adsorbent is permanently ensheathed with stable colloidal aggregates which become moulded into the familiar shapes by alveolar and bronchial currents.

Some support of this is adduced by the fact that micro-organisms adsorb colloidal material in the presence of blood serum and colloidal asbestos. Staphylococci so treated appear as large round yellowish-brown discs, and in the process their property of staining with aniline dyes has been lost. The organisms coalesce and form masses, and I think it probable that some of the coccal and spore forms of the curious bodies are similar organisms.

Finally, the answer to our second question must be found. Are the curious bodies diagnostic of pulmonary asbestosis? Asbestos is unique among the minerals in being fibrous, and the dust generated during the manufacturing process is also unique. As can be imagined from the formation of the curious bodies, there is no reason why any fine spicule of mineral should not have colloidal matter deposited around it and become moulded into a curious body. But, as no other mineral dust is fibrous, this occurrence must be so rare as to be negligible from a diagnostic point of view. The conditions which, apparently, must obtain for the formation of the curious bodies are the presence of plasma proteins and fine spicules of difficultly soluble material. These conditions are ideally found in asbestos workers, and for this reason I believe the curious bodies, if found in any numbers, are pathognomonic of pulmonary asbestosis.

I must express my gratitude to Dr. S. A. Henry and Dr. E. R. A. Merewether of the Home Office for their stimulating interest in the work, and for providing me with many specimens of asbestos, and to Mr. T. H. Byrom, F.I.C., and Mr. J. A. Darbyshire, M.Sc., for their invaluable assistance in the chemical and physical side of the work.

LITERATURE.

- Cooke, W. E.: *British Medical Journal*, 1924, ii, 147.
Cooke, W. E., and Hill, C. F.: *Journ. Royal Microsc. Soc.*, 1927, 252.
Cooke, W. E., McDonald, S., and Oliver, T.: *British Medical Journal*, 1927, ii, 487.
Stewart, M. J.: *British Medical Journal*, 1928, ii, 585.

CLINICAL ASPECTS OF PULMONARY ASBESTOSIS.*

BY

A. C. HADDOW, M.B., CH.B.,
LEEDS.

History.

DURING 1926 Dr. Ian M. D. Grieve, who was then with me in general practice at Leeds, drew my attention to the fact that several patients under our care of late with chest troubles were asbestos workers, and we agreed that they exhibited features sufficiently in common to justify investigation of what seemed an uncommon type of pneumoconiosis. We accordingly examined every asbestos worker who came our way, some fifteen in all, and had most of these radiographed. Then, in March, 1927, a female patient with long-standing disease died in hospital, and Dr. Grieve attended the post-mortem examination. With Dr. A. L. Taylor he examined the sections of fibrosed lung, and found a remarkable condition which Professor Stewart recognized as similar to the case described by Dr. Cooke in 1924.

* Read in the Section of Occupational Diseases at the Annual Meeting of the British Medical Association, Manchester, 1929.

Sir Thomas Oliver came to see our cases, and he described the clinical findings before this Section of the British Medical Association at Edinburgh in 1927. The two patients he reported on (both women) have since died, one in March, 1928, and the other in February of this year. One male patient died in March, 1928, and it was the certification of this case as death from "broncho-pneumonia, secondary to chronic asbestos poisoning, causing pulmonary fibrosis," which brought about the publicity of a coroner's inquest and official recognition of this disease.

Fatal Cases.

Of these four fatal cases the average age at death was 41, and the average years actually spent in the factory were under twenty, all in asbestos mattress-making for most of the time. The male patient had worked until two days before death (on a light job), but the three others had been incapacitated for an average of two and a half years (three years and five months, two years and five months, and one year and ten months). Only in one of them had tubercle bacilli been found in the sputum. She was one of those reported by Sir Thomas Oliver, and at her own request no necropsy was held.

I cannot personally trace any other death of an asbestos worker in my practice except that of a woman who died of perforated gastric ulcer in Leeds Infirmary, and whose lungs were examined by Dr. A. L. Taylor, with positive findings, two years ago. No death had ever been traced to asbestos dust in Leeds until the present series, and the only asbestos workers recorded by the medical officer of health for Leeds as having died of phthisis were found to have worked at a gas-mantle factory where asbestos is not employed in any department. I know that many workers had drifted away from the asbestos factory because they believed it was unhealthy, whilst many women had ceased work to attend to their homes, and any fibrosis present would readily be overlooked in later years when death overtook them.

Advanced Cases.

In practice these workers do not call for much attendance until they reach the stage of total incapacity. Until then they are naturally more prone to chest troubles during the winter months, and are sometimes unfit for weeks at a time, but at the present moment I have only four cases under treatment. These are all permanently incapacitated. They are all women, and are fairly well at present, though they were very ill in February and March. Three of them have been away from work for eighteen months and the fourth for four and a half years. They were all in the mattress department. Their average age is 35, and their average of years spent in the factory is fourteen, dating back to their teens. They average three children apiece, which accounts for most of the time lost from work. The children are healthy enough.

Clinically, they all suffer from shortness of breath on exertion, have cough without expectoration, are lean, although not losing weight at present, and complain of anorexia with resulting weakness (for in Yorkshire one is only as strong as one's appetite). Pain is variable, and occurs from time to time with signs of localized pleurisy. Night sweats are absent, and there has been no haemoptysis. No tubercle bacilli have been found. Their hearts are normal, with a pulse rate approaching 90 at rest. The average full range of chest expansion is only one inch in the upper thorax and even less around the lower ribs. On percussion there is always some dullness at both bases, and usually over the lower lobes generally. The breath sounds are diminished over the same area, and the voice sounds are usually well conducted. The respiratory murmur in the upper lobes is broncho-vesicular. Friction sounds can be heard at the bases and sometimes in the flanks. Moist rales and crepitations are for the most part absent now, but were abundant at the bases during the winter. These exacerbations have a serious bearing on the cases, and it will be noted that all the deaths have occurred at the end of winter.

Onset of Symptoms.—It is very difficult to ascertain the date of onset of symptoms, but the patients generally believe it was after five years of work. Depending as it does on so many factors, it must be extremely variable.

Our earliest clinical diagnosis is also in the neighbourhood of five years of work.

X-ray Findings.—Excellent radiographs of these four advanced cases were taken for me by Dr. Rowden recently. Exposure was for two-fifths of a second. All the chests showed abnormal radiolucency and marked limitation of diaphragmatic movement. The radiographs show this translucency above the level of the ninth rib posteriorly and the fourth rib in front. Below that level, roughly, in all of them there is definite infiltration, particularly on the right side. The outline of the diaphragm tends to be irregular and flattened, and the root shadows to be enlarged in a downward direction. Without going into further detail, the picture is that of asbestosis. But the shadow of the fibrosed zone is in itself characteristic. It is devoid of radiations from the root shadow and presents an amorphous, flocculent appearance, in which small cavities are indicated in some of the cases. In estimating the respiratory handicap we must not forget that the bases extend below the shadow of the dome of the diaphragm.

Lung Puncture.—Another line of investigation, suggested by Dr. Henry, has been carried out—namely, lung puncture. Asbestosis bodies can be extracted from the patients' lungs during life by means of a hollow needle and syringe. I punctured all these four cases, and Professor Stewart has found "bodies" in two of them. We do not recommend it as a routine method of diagnosis. No tubercle bacilli were found in any of them.

Sputum.—Much preferable is the examination of sputum by the antiformin method, which Professor Stewart has applied with striking success, and the only difficulty is to get the sputum. This is often available in winter, and if the patient has a receptacle in readiness the sputum will turn up some time. Three of these cases yielded the "bodies" in sputum, and the fourth had no sputum but was positive on lung puncture.

Conclusions.

The following conclusions have been established:

1. That the inhalation of asbestos dust will "in the long run" produce a state of fibrosis, the distribution being peripheral and mainly basal, the right base becoming most advanced and the upper lobes tending to develop compensatory emphysema as first the diaphragmatic and gradually the entire pleural surfaces become obliterated.
2. That the disease is not as a rule complicated by tuberculosis.
3. That the disease is usually first recognized after more than five years' exposure, although systematic examinations might reveal it earlier.
4. That the condition is usually found first during an attack of influenza or winter cold, when exacerbations of the disease occur.
5. That even in the advanced cases there is marked abatement during the summer months.
6. That the onset of anorexia signalizes to the worker that work is no longer possible.
7. That the patient may then live for several years and continue to bear children, but becomes progressively weaker and more emaciated, more hopeless, sleepless, and exhausted, until an attack of broncho-pneumonia or bronchitis brings death at last.
8. Further, that asbestosis bodies can be found in the sputum, if any, in all advanced cases, as well as in most early cases.
9. That they can be demonstrated by lung puncture in advanced cases; but since no patient is likely to permit more than one puncture to be made a negative finding is inconclusive.
10. That radiography is of great value in diagnosis and in observing progress.
11. That the treatment of these cases, being purely palliative so far, calls for great patience and vigilance over several years, and much relief can be afforded.
12. Finally, that the disclosure of the danger arising from asbestos dust has brought home to the workers the need for observing the measures provided for their protection, and is bound to be salutary, although the mortality statistics will probably be swelled by the wider recognition of this disease.

Memoranda:

MEDICAL, SURGICAL, OBSTETRICAL.

A METHOD OF EXAMINING THE SPUTUM FOR ASBESTOSIS BODIES.

HALF an ounce of so of sputum is added to an equal quantity of undiluted antiformin. This is gently agitated until the sputum is completely dissolved, after which it is diluted down with two or three ounces of water and allowed to stand in a large test tube for three or four hours. The bulk of the supernatant fluid having then been decanted, the remainder is centrifuged at a moderate speed for ten to fifteen minutes. The whole of the supernatant is now poured off, and the deposit transferred to an albuminized slide by means of a pipette. After thorough drying on a hot plate and final fixation over a bunsen, the film is very gently washed in water, dried, and mounted in Canada balsam. After a little experience the asbestosis bodies are readily picked up with the low power, and their true nature is then confirmed with the 1/6 inch or oil immersion lens. As a rule they are present in very small numbers, perhaps only one or two in a whole film. In the case of some of the older workers, however, numerous bodies, up to one or two per field in certain portions of the film, have been found. It is obviously necessary to cleanse thoroughly all glassware, etc., used in making these examinations, otherwise there is a risk of contamination of subsequent specimens.

The films can be treated with hydrochloric acid and potassium ferrocyanide to demonstrate the Prussian blue reaction given by the bodies.

Pathology Department, School of Medicine,
University of Leeds.

M. J. STEWART.

FOREIGN BODY IN THE LARYNX.

THE following case illustrates some of the difficulties of diagnosing the presence of a foreign body in the larynx, and the need, whatever the history may be, for making the most complete examination when this condition is suspected.

The patient, a waitress at a country hotel, consulted Dr. Frend of Hurstpierpoint on January 24th, stating that on the previous evening, as she got up from her chair, where she had been engaged in sewing, she felt a sudden pricking in the throat. After repeated questioning she admitted that she was in the habit of holding pins in her mouth, but stoutly denied that she had done so on this occasion. The meal which she had taken two hours previously contained nothing that could be impacted as a foreign body in the throat. Dr. Frend could see no abnormality in the larynx or pharynx, but with great difficulty he persuaded the patient to go to the Sussex Throat Hospital for further examination.

My examination by indirect laryngoscopy yielded negative findings, but as the patient still complained of the pricking sensation I asked Dr. Prowse to make an x-ray examination. I confess that I was surprised to see in the radiogram that a pin was lying, horizontally and in an antero-posterior plane, opposite the sixth cervical vertebra—that is, on a level with the entrance to the larynx. The laryngeal mirror had failed to reveal the presence of this pin, although the patient's larynx had presented no difficulty in examination. I decided to examine with the endoscope, and did so the same evening with the patient under general anaesthesia. I used Lynah's oesophageal spatula, and at once discovered a longitudinal tear about one inch in length on the posterior oesophageal wall. This discovery was soon followed by the sight of a pin-point abrasion on the posterior aspect of the right arytenoid. By making the necessary adjustment in the patient's position I brought the larynx into view, and was able to see the pin transfixing through the right arytenoid, with most of it lying free in the middle of the larynx. I grasped the pin with forceps and inserted the tube a little deeper; by this manoeuvre I was able to draw the pin forward wholly into the tube and thus eliminate any possibility of damage to the tissues. The patient wrote on February 17th that she had suffered no soreness or hoarseness, and that she had not felt the least discomfort.

The following features of this case are of special interest:

- (1) The absence of symptoms, even of the wheezing respiration described by Chevalier Jackson.
- (2) The unusual position of the pin—foreign bodies in the larynx usually lie with the long axis in the sagittal plane.
- (3) The failure to see the pin by indirect laryngoscopy. One concludes that it must have been tucked under the aryteno-epiglottidean fold and brought into view as a result of