in occasional cases nausea and vomiting persist for a time, even up to twenty-four hours. There is no unpleasant taste or smell lingering.

On questioning a medical man to whom I recently administered ethylene for an appendix operation and colopexy his criticism was that "the experience was not at all unpleasant either going under or coming out."

The Surgeon's Point of View.

From the surgeon's viewpoint several factors come into play, depending largely on the type of operation. The principal criticisms are directed towards insufficient relaxation and haemorrhage. In operations where neither of these is of serious moment the anaesthesia meets most requirements.

The Anaesthetist's Point of View.

Lastly, the anaesthetist, though dealing with an anaesthetic which requires an elaborate apparatus, and an anaesthesia demanding experience and constant watchfulness to maintain within its rather narrow margins, is working within fairly wide limits of safety. In the asphyxial stage which develops when too little oxygen is allowed respiratory failure definitely precedes cardiac failure. Ample warning is therefore given by the dark colour of the blood. If this be observed early, and dealt with immediately by an adequate increase in the oxygen content, it is rare that any further steps are necessary.

In an experience extending to over a thousand administrations I have only seen six instances of serious respiratory collapse with cessation of breathing. In four of these by administering pure oxygen under positive pressure, very easily carried out with McKesson's apparatus, a return of breathing was quickly obtained. In the other two cases this had to be supplemented for brief periods by rhythmic chest compression.

There is one other physical property of ethylene besides its odour to which reference must be made. It is very inflammable, and in mixtures with air or oxygen may readily be ignited and even exploded. In the United States cases of explosion have been recorded. So far as I am aware none has happened in this country. The nature of the hospital buildings and the drier atmosphere of the States probably combine to promote conditions more favourable to this occurrence. I think that, in this country, it is sufficient to remember that ethylene must not be administered in any room where there is an open fire or naked light, nor in any case where cautery or diathermy is to be used.

I have attempted briefly to outline the scope and limitations of ethylene as an anaesthetic. To those who are familiar with nitrous oxide and oxygen it presents no difficulties, and is well worth a trial. I personally should now regret not having it available for many cases in which I think it is the best anaesthetic.

ERYTHRAEMIA WITH A HIGH DEGREE OF ACHOLURIC JAUNDICE:

PROBABLY A MANIFESTATION OF COMPENSATORY "HYPERSPLENISM."

BY

F. PARKES WEBER, M.D., F.R.C.P.,

SENIOR PHYSICIAN TO THE GERMAN HOSPITAL, LONDON.

SOME excess of urobilin in the urine has not rarely been observed in patients suffering from erythraemia (splenomegalic polycythaemia). Lommel, and Löw and Popper, found the total excretion of urobilin and urobilinogen above the average, and the blood bilirubin may be in excess, but it is very rarely sufficiently so to give rise to actual (acholuric) jaundice. I have never myself seen or heard of the occurrence in England of another case of jaundiced erythraemia* like that of the present patient, whom I first saw in April, 1929, by the kindness of Dr. S. S. Chesser. She then showed, as she still (October 17th, 1929) does, considerable jaundice, without bilirubin in the urine (acholuric jaundice), but with a very highly positive indirect Hijmans van den Bergh reaction in the blood plasma—that is to say, for "blood bilirubin" (which, by the by, I believe to be the same as Virchow's "haematoidin").

The patient,† a Welsh woman, married, aged 40 years, living near London, has a typical "erythraemic" appearance and is markedly jaundiced (examined in April and May and October, 1929). Height 152.4 cm.; weight 71.6 kilograms in May, and about 1½ kilograms more in October, 1929. There is decided excess of subcutaneous fat over the thighs, buttocks, and abdomen. Examination of the thorax shows nothing special, excepting (by x rays) rather excessive bronchial markings and very slight enlargement of the aortic arch.

The spleen is enlarged and feels rather firm, the lower edge reaching three fingerbreadths below the costal margin. The liver (not abnormally hard) can be palpated three and a half fingerbreadths below the ribs in the right nipple line. No ascites. No subcutaneous oedema. Menstruation regular. Bowels not confined. Brachial blood pressure 165/95 mm. Hg. Urine (October 17th): specific gravity 1011; clear; rather high yellow; acid; a little albumin (moderate "cloud" on boiling); no sugar; excess of urobilin and urobilinogen; no bilirubin; excess of indican; one or two hyaline tube casts seen in the centrifuge sediment. Ophthalmoscopic examination shows a typically "erythraemic fundus." Basal metabolism (two estimations), 127 per cent.

* For references to the literature on the association of erythraemia with jaundice and also with hepatic cirrhosis see Weber and Bode, *Poly-cythaemia, etc.*, London, 1929, p. 13.

† The case was demonstrated, but not exactly described, at the meeting of the Clinical Section of the Royal Society of Medicine on November 8th, 1929.

Blood count (April 29th): haemoglobin 120 per cent.; erythrocytes, 7,880,000 per c.mm. of blood; colour index = 0.76; white cells 16,350 (polymorphonuclears 70 per cent., basophils 3 per cent., eosinophils 3 per cent., lymphocytes 21 per cent., monocytes 3 per cent.); no nucleated or abnormal red cells seen. Thrombocytes: 1,512,960 per c.mm. of blood. The blood serum gives an extremely strong positive indirect Hijmans van den Bergh reaction for "blood bilirubin." The resistance of the washed erythrocytes towards graduated hypotonic sodium chloride solutions seems to be within normal limits, haemolysis commencing with the 0.5 per cent. solution and becoming complete with the 0.32 per cent. solution. The blood serum gives negative Wassermann and Meinicke reactions. Dr. T. C. Hunt kindly estimated the cholesterol in the blood serum by the Autenrieth colorimetric method (as described by L. Pincussen, *Mikromethodik*, fourth edition, Leipzig, 1928, p. 143), and found it to be 190 mg. per 100 c.cm. A recent blood count (October 17th) showed decided increase in the polycythaemia: haemoglobin 125 per cent.; erythrocytes 8,900,000; colour index = 0.7; white cells 18,650. The erythrocyte count still more recently (November 7th) was 10,100,000.

The patient was married at 21 years of age and has had three children (all living and healthy); no miscarriages. There is nothing special in her family history. She is said to have had typhoid fever in childhood. Her first attack of jaundice was eight or nine years ago, and since then she has had many icteric exacerbations, but of late seems never to have become quite free from jaundice. She began to suffer from vertigo twelve years ago, and the giddiness is worse during the exacerbations of jaundice. The attacks of giddiness come on especially when she is walking in the open air, and during a recent attack she fell and cut her forehead (October 15th). For a long time she has been subject to occasional attacks of pain in the left lumbar region, passing down the outer part of the left thigh, but she has had no attack suggesting biliary colic. She suffers from occasional epistaxis and often has severe headaches. It was five years ago that she first noticed that her face had become redder and that she flushed very easily. She has been relieved by blood-letting, and this is to be carried out again. It is certain that her jaundice varies much in degree. In fact, at times there is no obvious jaundice.

Remarks.

It seems probable that the chronic acholuric jaundice and the icteric exacerbations in the present case are due to active destruction of the erythrocytes in the spleen (and possibly in other parts of the reticulo-endothelial system), but there is no evidence of abnormal "fragility" of the red cells. The jaundice might therefore be interpreted as the manifestation of a spontaneous attempt to compensate for the excessive production of erythrocytes in the bone marrow by increased haemolysis in the spleen—a compensatory form of "hypersplenism."* Periodic blood-letting would seem to be the safest treatment in such a

* If this is so an increase in the degree of jaundice should be accompanied by a decrease in the erythrocyte count. On one occasion we have noted a decrease in her jaundice accompanied by an increase in her erythrocyte count.

case; phenyl-hydrazine is perhaps contraindicated, since, by further destruction of erythrocytes, it might not only increase the jaundice but also favour the tendency to formation of bilirubin calculi, and possibly even produce hepatic cirrhosis.

My explanation of the whole morbid syndrome in the present case is therefore as follows. The primary "agent" is the erythraemia, which is an "idiopathic" increase of the total erythrocytes and erythroblastic bone marrow in the body—that is to say, an idiopathic increase of the "erythron," the term proposed by Professor A. E. Boycott in his recent presidential address before the Section of Pathology of the Royal Society of Medicine. If we regard the "erythron" as a tissue-organ or tissue-system (as we regard the reticulo-endothelial system and the total lymphadenoid tissue of the body), the increase in question must be of the nature of a primary diffuse but non-malignant neoplasm (a "neoplastic diffuse overgrowth"), and not of the nature of a secondary compensatory hypertrophy, such as we get in the polycythaemia of high altitudes, the polycythaemia of congenital heart diseases, and the polycythaemia connected with circulatory obstruction in the lungs—for example, in "Ayerza's disease." A true *neoplastic* overgrowth of the "erythron" is useless and harmful to the rest of the body, and in the present case a spontaneous compensatory reaction on the part of the spleen has taken place—namely, an increase of one of the splenic functions (a compensatory "hypersplenism"), with the object of destroying the superfluous erythrocytes. This in its turn gives rise to excess of the blood bilirubin and to a condition of acholuric jaundice, but is nevertheless insufficient to prevent the gradual further increase of the "erythron," which will have to be met by blood-letting. It is just possible that a diet qualitatively the reverse of that employed for pernicious anaemia may likewise be of some use.

SECONDARY COMPLICATIONS IN PATIENTS SUFFERING FROM EMPYEMA OF THE PLEURA.

BY

L. HOFBAUER, M.D.,

CHIEF OF THE DEPARTMENT FOR RESPIRATORY DISEASES, FIRST MEDICAL CLINIC, VIENNA.

By an operation for empyema dangers which threaten life are avoided, but the *restitutio ad integrum* is seldom attained. On the contrary, prolonged suppuration often follows (with, sometimes, death due to exhaustion), and repeated operations may be required. At the least, disability, disfigurement, and marked pathological conditions ensue. The surgeon's life-saving intervention may be criticized by the patient's relatives, and therefore fall into discredit. Taking the view that the untoward consequences are due to the wide opening of the chest, Lister adopted a method of treatment which avoided this, while Wenckebach prefers the thoracic puncture, followed by substitution of the exudate by air. Many excellent results have been obtained by this method, especially in cases of tuberculous empyema; but not all cases of empyema can be treated in this manner. Hathaway has shown that coagulated fibrin in the form of blood clot is an excellent culture medium for pyogenic bacteria. The clot, therefore, must first be removed, and this is only possible with a widely opened pleural cavity.

Post-operative complications are to be found, however, whether the chest is widely opened or not. They may be classified in the following groups: (1) deformities, especially of the thorax; (2) functional troubles, including dyspnoea and disturbances of the circulation; (3) dangers to life, such as arise from the "pyothorax *ex vacuo*."

Massive adhesions, the result of insufficient resorption of the exudate, are generally believed to be the cause of the first group. Attempts were made, therefore, to empty the

effusion as thoroughly as possible; often, however, there are insurmountable difficulties to doing this completely, and even when all the effusion has been removed the patients suffer from the complaints already mentioned, sometimes in the highest degree. Those cases in which complete elimination of the exudate is attained by the Monzingo form of drainage show the greatest changes. Severe symptoms of breathlessness, oppression, and a peculiar type of dyspnoea are present, although the x-ray examination shows that the pleural space is entirely free from exudate.

FIG. 1.—Dislocation of the diaphragm by bending. ——— Position of the organ in a sitting man. - - - - - Position of the organ in a bending man.

As soon as these patients try to bend sideways they experience a feeling of constriction in the chest, with pain and the sensation that they cannot get enough air. In many patients this symptom appears only when bending over to the diseased side; in others, when they bend to the sound side.

These complaints are comprehensible when it is remembered that they correlate with the presence of adhesions of the diaphragm to the bony wall and disappearance of the phrenico-costal angle. The condition becomes obvious when comparison is made with the normal displacement of the diaphragm following the bending of the body sideways. The lower part of the diaphragm is displaced upwards into the thoracic cavity, while the upper half descends into the abdominal cavity. The reason for this dislocation is that the intestines fall on the lower half of the diaphragm (Fig. 1); the abdominal pressure on this area becomes increased, therefore, and the upper part of the diaphragm is pulled downwards. The phrenico-costal angle opens on this latter side to its fullest extent.

The presence of scar tissue in the phrenico-costal angle interferes with its opening. The patient feels pain when

FIG. 2.—Pressure on a fixed midriff on bending. - - - - - Position of the diaphragm in the sitting man. ——— Slope caused by the altered pressure in bending.

he inclines to the healthy side, because the scar tissue is stretched and pulled. These symptoms are more marked the more the patient inclines the upper part of his body to the horizontal position, because the dislocation of the intestines increases the pressure under the lower part of the diaphragm the more the body is bent. On the other hand, when fixed to the thoracic wall the diaphragm cannot follow the increased abdominal pressure when the patient bends his body to the diseased side; it becomes expanded (Fig. 2), and the patient feels pain owing to stretching of the adhesions and involvement of the nerves.

* I use the term (morbid) "agent" as something which provokes a compensatory or conservative "reaction" (compare my short paper, "A Note on Action and Reaction in Pathology and Therapeutics," *St. Bartholomew's Hospital Reports*, London, 1903, xxxix, p. 139).