the lungs nothing abnormal beyond signs of some congestion at the bases could be discovered. The cervical glands on the left side were enlarged.

A lumbar puncture was performed. The cerebro-spinal fluid showed a slight increase in globulin and total protein; the sugar reduction was diminished, the chlorides were normal, and the cell count 18 per c.mm., the lymphocytes and polymorphonuclears being in equal proportions. No micro-organisms were seen and the culture was sterile. Blood was also taken for culture, but only staphylococci, which were considered to be due to a contamination, were found.

The gradual "march symptoms" of a hemiplegia formed a definite indication of a cerebral abscess in this case. The boy's condition was extremely grave on admission, and in addition he had valvular disease of the heart. However, the localization of the abscess being adequate, it was decided that the chance of operation should be given, but unfortunately he did not survive.

There are two theories of the production of cerebral abscess after tonsillectomy. One is that infected pulmonary emboli result from the sepsis in the tonsillar bed; the other, that septic material is inhaled while the patient is under the anaesthetic, and that the cerebral abscess is secondary to one or other of these conditions. The present case—in which early broncho-pneumonia was demonstrated at the necropsy—would appear to fall into the group of the latter theory. The possibility of cerebral abscess, though rarely associated with tonsillar infection, is not to be ignored.

Vertical section of brain, showing location of abscess in sub-cortical region of right frontal lobe.

A post-mortem examination was performed by Dr. R. T. Chadwick, who reported as follows.

The brain dura mater was dark in colour and congested. The arachnoid and pia mater were intensely congested and opaque; the blood in the vessels was fluid. No signs of emboli and no pus were discovered. The convolutions on the right side were flattened

and the ventricles were filled with clear cerebro-spinal fluid, the left being normal in size, the right flattened by the abscess.

The abscess, which was filled with greenish-red, thick pneumococcal pus, was $2\frac{1}{2}$ inches in diameter and circular in shape; it was situated posteriorly to the fissure of Rolando, extending downwards from the vertex about $1\frac{1}{2}$ inches on the external surface, and the same distance on the internal surface of the right hemisphere, lying about $\frac{1}{8}$ inch deep to the brain surface. On the vertex the meninges were intensely congested, and the abscess showed signs of pointing. The medulla was congested, and there were no pressure marks. No sign of tuberculosis or post-basis meningitis was detected, but the brain substance was intensely congested generally.

The tonsillar fossae were found to be healing after the operation, and no sloughs were present. The cervical glands were slightly enlarged, but there were no abscesses therein. The lungs showed an early and moderately intense broncho-pneumonia; the air passages were intensely inflamed. There was great right-sided dilatation of the heart, the mitral orifice admitting two fingers. Old endocarditis was present on the mitral cusps, but the tricuspid and aortic valves were healthy. There was general congestion of the alimentary system, and the mesenteric glands showed evidence of past healed tuberculosis. The kidneys were greatly congested.

ORBITAL CELLULITIS IN CHILDREN.*

BY

E. TEMPLE SMITH, F.R.C.S.ED., D.O.Oxon.,
F.C.S.A.,

HONORARY OPHTHALMIC SURGEON TO THE ROYAL ALEXANDRA HOSPITAL
FOR CHILDREN AND THE ROYAL NORTH SHORE HOSPITAL,
SYDNEY, AUSTRALIA.

SINCE orbital cellulitis may occur in children of a few weeks old there is no age limit. The essential feature is proptosis of the eye associated with signs of inflammation. Proptosis alone is not sufficient for its diagnosis. Proptosis unassociated with inflammation may be caused by a cyst, a mucocele, a new growth, or what not. Until one meets the combination of a congested eye and pink swollen lids with proptosis the diagnosis cannot be established. Once made, a grave prognosis should be given and the help of the specialist sought.

The diagnosis is not, however, always easy, as it is at times very difficult to be sure whether or not proptosis is present when chemosis of the conjunctiva and a swollen lid are the most prominent features. But it is on the existence of this sign that the diagnosis will depend. Proptosis is mechanically produced by congestion of, or exudation into, the orbital tissues behind the orbital fascia. This is a dense and well-fitting septum separating the orbit proper from the conjunctival sac and the anterior segment and interior of the eyeball. It matters not how severely infected an eye may be, orbital cellulitis, with its cardinal symptom of proptosis, will not arise unless the orbital fascia is penetrated.

It is clear, then, that orbital cellulitis must not be diagnosed because oedema of the eyelid is present, or even cellulitis of the eyelid. The latter is often due to the spread of infection from a hordeolum or impetiginous sore in the neighbourhood; and oedema can often be traced to a cellulitis of the scalp or face, or dacryocystitis, and is then present merely by extension, owing to the loose cellular structure of the lids.

Again, an intense purulent conjunctivitis, gonorrhoeal or diphtheritic, may occasionally give rise to a pink and

swollen lid; ocular diphtheria is frequently at first monocular. The diagnosis will be established by the presence of a profuse discharge or of a membrane on the everted lids. For the determination of the presence or absence of proptosis lid retractors are essential. These may be improvised by the use of two hairpins bent over near the blunt end. If a local cause has been discovered for the swollen lid, whether abscess, conjunctivitis, dacryocystitis, or foreign body, it will be treated on appropriate lines. If none such is found and proptosis is present or strongly suspected, a grave view must be taken of the case, as it is one of orbital cellulitis.

Orbital cellulitis in children has, so far as my experience goes, three chief causes:

1. Skin infections, such as impetigo, hordeolum, fly-bite, boils, or any facial wound, with accidentally superadded virulent infection. This is more likely to be met with in older children. The temperature is generally very high; septic cavernous sinus thrombosis and general septicæmia with pneumonia are likely to supervene, and they often end fatally and rapidly. The infection passes by the angular vein to the cavernous sinus. Fortunately such cases are rare.

2. Nasal sinus infection, the commonest cause, is met with in young infants as well as in older children. The temperature is not usually very high and the child is not very ill. The eye is pushed outwards as well as forwards. This is a strong diagnostic point. I will go so far as to say that if the eye is not displaced outwards the case is not nasal in origin. In these cases the services of the rhinologist should be sought; he will, if he thinks necessary, open up the antrum or ethmoidal cells. I believe the most usual condition is an ethmoiditis. If the proptosis is only due to cedema it will gradually subside. If, however, pus has formed in the orbit a counter-opening between the nose and eye may be needed. Convalescence is slow but sure, and the eye is not usually in jeopardy. I have seen a considerable number of cases of this type of cellulitis in the last fifteen years. They are sometimes associated with the exanthemata.

3. Osteomyelitis or periostitis of the orbit is a third principal cause. Until about three years ago I had not seen such a case. I was then asked to attend a young

*An address given to the Paediatric Section of the New South Wales Branch of the British Medical Association on April 11th, 1930.

infant with a high temperature and pink oedema of the eyelids of both eyes, who was very ill. There was no injection of the ocular conjunctiva. As both eyes were affected it was difficult to say whether or not proptosis was present. I believed, however, that it would be by the morrow, and that it was an early case of septic cavernous sinus thrombosis, that nothing could be done, but that my rhinologist colleague should be asked if he concurred. He did concur. On the next day I still found no ocular injection to speak of and no definite proptosis, although the lids were more pink and swollen. I realized then that I had made an erroneous diagnosis, being deceived by the binocular affection, which will generally point to a common origin. I asked that my colleague should see the child again. He found an osteomyelitis of the lower rims of both orbits. Incisions were made over the malar bones, and the child recovered. This was not a typical case, however, as it was binocular—surely a rare occurrence. Furthermore, although there was osteomyelitis of the malar bones, the lesion was anterior to the attachment of the orbital fascia; therefore, though oedema of the lids was present, there was no orbital cellulitis.

I have since seen three cases of orbital cellulitis (monocular) caused by osteomyelitis of the orbit, and not, apparently, associated with or due to sinus infection. I am sure that these cases are metastatic. They have not been coincidental with a similar lesion in the long bones and epiphyses, although they are without doubt analogous. The child becomes very ill indeed, with a high and septic temperature. Pink oedema of the lids and definite proptosis are early signs. The eye is pushed forward without being pushed outwards. Presumably the eye might be pushed to some extent away from the side of origin of the infection, wherever it might be; but, in the early stages at any rate, when the proptosis is merely due to oedema and not to formation of pus, there is no lateral displacement. This is in contradistinction to those of nasal origin. In the last three cases I have seen, the point of maximal swelling and tenseness of tissues has been at the lower outer side of the orbit. Possibly a trauma may have been the exciting cause.

In the first case I was not called in for several days, as a tentative diagnosis of a nasal origin had been made, and the rhinologist had explored the sinuses with an indecisive result and with no improvement in the patient. I found the outer lower orbital margin tense and swollen, and incised it here, getting pus freely. The boy had a high temperature for many weeks, and was very ill. Although there was free evacuation of pus, the orbital oedema and proptosis persisted, and in spite of all efforts the cornea sloughed, and the eye had to be eviscerated. Subsequently he developed a mild osteomyelitis of the facial surface of the malar bone.

The next case was that of an infant, seen in private, with proptosis and similar symptoms. I temporized for a day, and then opened the orbit at the outer angle, evacuating pus. Again the orbital oedema was so great and persistent that the cornea sloughed, and the eye had to be eviscerated, though the child recovered. Sequestra formed for a long time.

The third case was that of a boy of 5, with the same symptoms of high temperature, severe illness, inflammatory central proptosis, and the chief swelling at the lower outer orbital margin. He had been ill two or three days. This time, profiting by my experience, I opened the orbit within half an hour of seeing him, and passed a sinus forceps through the orbital fascia down to the bony floor and rim. No pus was seen, but on the next day it appeared alongside the tube which had been left in. The wound was kept open, the proptosis went steadily down, and the eye was never in danger. He had a septic temperature for several weeks, and had some meningeal symptoms, which, however, subsided on lumbar puncture. *Staphylococcus aureus* was found in the blood and urine, so that there was evidently a very generalized infection. Later he had an attack of osteomyelitis of the tibia.

To recapitulate: apart from cases of orbital cellulitis due to facial infection, in which the orbital condition is only an incident in a fatal septicaemia, there are two main causes. The first, and the most common, is nasal sinus infection, with outward displacement of the proptosed eye and not very severe general illness. Here the rhinologist can give most help; and if he is called in early incision into the orbit will probably be avoided. The second is metastatic osteomyelitis or periostitis of the orbital walls, with

severe illness and temperature and central proptosis. The lesson I have learned from these latter cases is that early incision (as in the analogous disease of the long bones and epiphyses) is essential if the eye is to be saved. When once the orbital tissues are thoroughly infiltrated and oedematous it seems to be difficult or impossible to bring about resolution before the cornea has sloughed, owing to interference with its circulation as well as to exposure. Our aim should be to prevent this phlegmonous condition by prompt incision down to the bones involved. Hence the importance of early and accurate diagnosis. In every case of a pink and swollen eyelid the general practitioner should realize the significance of proptosis, if this exists. In its absence he may temporize; in its presence he should act with decision. No case of inflammatory exophthalmos should be regarded lightly.

Although I have dwelt in more detail on the signs and symptoms of the type of orbital cellulitis due to bone infection by the blood stream, that is not because it is the commonest, but because it is the gravest, and demands the most promptitude in its treatment. According to Roemer, more than 60 per cent. of cases are due to nasal sinus infection. He is referring to adults, and I think that this applies much more so to children, owing to the thinness of the orbital walls. In my own practice orbital cellulitis in adults is exceedingly rare, whereas in children it is at least not uncommon.

My aim in this short paper is twofold: (1) to emphasize the importance of proptosis when associated with inflammation; (2) to attempt to make clear the difference in signs and symptoms of the two main types of orbital cellulitis, and the difference in outlook and treatment which each demands.

Memoranda:

MEDICAL, SURGICAL, OBSTETRICAL.

RENAL CYST CONTAINING CALCULI.

(With Special Plate.)

THE following particulars of a renal case upon which I recently operated will, I think, be of general interest.

A man, aged 62, had always had good health until seven years ago, when he noticed a swelling in his left lower abdomen. He consulted a surgeon, who, after examination, advised an operation. After the operation he was told that there had apparently been a cystic condition of his left kidney, the wall of which had been removed, and a drain put down into the cavity. He made a good recovery, and had no trouble until three years ago, when the swelling returned; on this occasion his doctor aspirated and drew off a large quantity of fluid.

A few months ago he came into my hands, and I found a large mass occupying a considerable proportion of the left hypochondriac and lumbar regions of the abdomen. It was dull on percussion, and its range of movement was practically nil. X-ray examination showed a "corn-cob"-like mass of calculi (see plate).

A cystoscopy having been performed, and a pyelogram taken, the efficiency of the right kidney was proved good. No solution could be made to travel more than an inch up the left ureter, but on the right side there was a normally placed organ. It was then decided to explore him.

Operation.—The old scar on the anterior abdominal wall was removed, and the abdomen was opened as carefully as possible. The sac of the cyst was, however, so adherent to the abdominal wall that it was at first punctured, and a dark grumous-looking fluid escaped. This being controlled, the cyst was slowly separated from adhesions with which it was completely surrounded. This was no easy matter, since it was necessary to avoid as much as possible damaging the numerous vessels of the mesentery which ran across it. Ultimately, however, the whole cyst was shelled out. With the exception of the descending colon, bowel was never seen throughout the whole operation; we had been apparently working in a retroperitoneal position from the first, the slowly enlarging kidney having pushed everything aside in its advance anteriorly. The cavity was drained and the aperture in the mesentery closed. The patient made an uninterrupted recovery.

Pathology.—On the sac being freely opened more than two pints of grumous fluid escaped. The sac was directly continuous with an enormously dilated kidney pelvis; this in turn expanded into wide, deep calyces. The kidney pelvis and calyces contained hundreds