ON BOXER'S THUMB.

BY

I. H. LLOYD-WILLIAMS, M.C., M.B., B.S.,

HONORARY MEDICAL OFFICER, HAYWARDS HEATH HOSPITAL, SUSSEX.

(With Special Plate.)

SUBLUXATION of the first metacarpo-phalangeal joint is a condition occurring chiefly among boxers, with whom it has attained almost the status of a stigma of the prize ring. It does, however, occur as the result of other accidents. The deformity may be either a primary condition, as with boxers, or a residual condition arising from a complete dislocation. In boxing the subluxation is produced by a blow on the distal portion of the thumb when it is not efficiently supported by the clenched fingers. The impact of a properly delivered blow should in no case directly affect the thumb, but it may be damaged by the opponent's guard.

Anatomy.

The head of the first metacarpal is somewhat flatter than the heads of the other metacarpals, and strong lateral ligaments bind it to the phalanx. Crossing the palmar aspect of the joint in question are the tendons of the adductor obliquus, opponens, and flexor brevis pollicis muscles. The first two are inserted by a common tendon into the ulnar side of the phalanx, and the short flexor into the radial side. In each tendon is a sesamoid bone provided with a separate synovial membrane for its articulation with the metacarpal. Below the sesamoids the tendons are fused together by fibro-cartilaginous tissue which is intimately attached to the anterior ligament and capsule of the joint. The capsule, which is lax in the flexed position, is drawn down during flexion and is thus prevented from being caught between the bones. Situated dorsally are the extensors and a weak ligament.

Movements.

The *active* movements of the joint are flexion and extension. The *passive* movements are: (1) a lateral hinge movement corresponding to adduction and abduction of the other digits; (2) a gliding movement in the horizontal plane, the base of the phalanx being displaced towards the palm of the hand. A similar gliding movement of phalanx on metacarpal can be produced in the fingers, but it is of less extent. With very long lax ligaments hyperextension and some degree of subluxation may take place normally as a result of active movements without apparent injury to the joint.

The accompanying skiagrams may be considered typical of (a) the normal, and (b) the subluxation in boxer's thumb. The subject of this particular joint had had a complete dorsal dislocation, and subluxation resulted. Note particularly a dislocation of the phalanx in the horizontal plane towards the palm, and the wide gap between the sesamoids and the head of the metacarpal.

Symptoms.

Immediately after the injury the patient complains of pain in the joint and of weakness, particularly as regards grip. There is not usually much swelling at the time. If left untreated, weakness continues and is accompanied by limitation of movement, both active and passive. In old-established cases the joint is almost immobile. It assumes a position of semiflexion, and becomes somewhat fusiform in shape from thickening of the capsular structures. The swelling tends to be more abrupt than in rheumatoid arthritis.

Treatment.

Digital reposition of the bones is easily accomplished by manipulation in the horizontal plane, with or without traction in the long axis of the thumb. A correct position is, however, not generally maintained. The problem, therefore, is not so much to reduce a dislocation as to make this reduction permanently effective.

Dislocation does not occur unless both the lateral ligaments and the capsule of the joint have been ruptured. The posterior ligament is relatively inefficient compared with

the lateral and anterior ligaments. I have confirmed this by numerous experiments on the cadaver. A complete dislocation, in which the joint capsule and ligaments are ruptured, involves also a tearing of the attachments of the sesamoids to the metacarpal head. But in no case was there intervention of fibrocartilaginous or other tissue between the articular surfaces or between the tendons and the metacarpal, although I had imagined at one time that such intervention was a possible cause of failure in maintaining reduction.

Recurring or persisting deformity is caused by the unopposed traction of the short muscles of the thumb in the presence of rupture of the lateral ligaments and capsule. Treatment must therefore be directed to neutralizing this muscular pull. The first essential is a dorsal reduction of the lower end of the phalanx whereby it is brought into correct anatomical alignment, while maintaining a certain degree of flexion of the joint. This semiflexed position is best maintained by a special splint which exerts backward

FIG. 3.—Illustration of the splint.

pressure on the lower end of the phalanx and the tendons, and counter-pressure both on the distal end of the phalanx and on the metacarpal. The moulded metal splint illustrated was made for me by Messrs. Lindsey and Sons of Brighton. It should be worn for about fourteen days to allow the healing of the capsule and ligaments; massage and passive movements should then be instituted, and the splint kept on for another week or fortnight.

Note.—I have tried both moulded poroplastic and strapping, but these methods have not proved satisfactory in my experience. Poroplastic does not lend itself to fine adjustments; and the essential counter-pressure producing a certain amount of flexion is not obtained by strapping.

OSTEITIS FIBROSA WITH INITIAL SYMPTOMS
RESEMBLING RHEUMATIC INFECTION.

REPORT OF CASE.

BY

J. SALISBURY CRAIG, M.B., CH.B.GLAS.,
DISPENSARY PHYSICIAN, WESTERN INFIRMARY, GLASGOW;

AND

JAMES H. SHEARER, M.B., CH.B.GLAS.,
STRATHAVEN.

(With Special Plate.)

THE cases of generalized osteitis fibrosa described by Wilder,¹ Hunter,² and others in 1929 presented signs of progressive weakness, loss of muscle tone, anaemia, pain in the bones, and decalcification of the skeleton, which were attributable to hyperfunction of a parathyroid tumour. The following case has been considered worthy of record since it conforms in many respects to those previously noted, and because it was one in which an initial clinical picture, suggestive of subacute rheumatism, was suddenly altered by the occurrence of a pathological fracture.

In May, 1929, the patient, an unmarried woman aged 50, complained of pain over the posterior aspect of the neck, the left deltoid region, and the left upper arm. Tenderness over those regions was then considered to be due to fibrositis. Anorexia, dyspnoea, mild pyrexia, and pain in the left infra-axillary region

ARTHUR JACOBS: HORSESHOE KIDNEY.

FIG. 1.—Calculous hydronephrosis in the right segment of a horseshoe kidney. The anomalous axis of the calyces and the relation of the ureters are shown.

FIG. 2.—Resected segment of horseshoe kidney. The large calculus can be seen within the dilated pelvis, the anterior wall of which has been opened.

FIG. 3.—Hydronephrosis in the left segment of a horseshoe kidney.

J. SALISBURY CRAIG AND
J. H. SHEARER:
OSTEITIS FIBROSA.

FIG. 1.—Pathological fracture of left humerus. Areas of rarefaction in radius and ulna.

I. H. LLOYD-WILLIAMS: BOXER'S THUMB.

FIG. 1.—Radiograph of normal thumb, showing the normal position of the bones and sesamoids.

FIG. 2.—Radiograph of boxer's thumb. Note the space between the sesamoids and metacarpal and the subluxation of the phalanx.

FIG. 2.—Areas of rarefaction in right humerus, radius, and ulna.