membrane, sloughing—the result of overdosage—and stricture of the external meatus.

It is a striking fact that in all except the advanced cases of penile cancer the removal of the primary lesion results in a decrease of the size of the inguinal glands, and in some cases complete disappearance. This, of course, means that the enlargement is due principally to sepsis, and not to malignant disease. The treatment of these enlarged glands depends upon the clinician's interpretation of the pathological processes at work. My view is that to dissect out the lymphatic tissue at the same time as the primary growth is eradicated is a most dangerous practice. First, lymphatic oedema may result, comparable to the condition known as "brawny arm," following the radical operation for carcinoma of the breast; secondly, there may be an extensive cellulitis, due to operative intervention in tissues heavily charged with infection.

As the malignant cells do not penetrate beyond the inguinal regions it is perfectly safe to allow three months to elapse after the removal of the growth of the penis, in order to observe any changes in the lymphatic tissues. As I have already stated, sepsis is the chief factor in causing the glands to enlarge, and the elimination of the primary focus will cause the disappearance of the adenitis within a few weeks. If at the end of three months the

glands are still larger than normal, and the skin over them is thickened, we can be quite certain that these changes are due to malignant disease. It is my practice, then, to excise the diseased tissues, rather than to speculate with radium.

Surgical Removal of Primary Growth.

Finally, we have to answer the all-important question: When is the primary growth to be irradiated, and when is it to be removed by operation?

Surgeons are unanimous in their opinion of the value of the Gould operation in eradicating the disease. The functional result, so far as micturition is concerned, is also satisfactory. There is complete control. Narrowing of the external meatus frequently occurs, and requires intermittent dilatation with bougies. This defect, and the loss of the whole organ, are the two great disadvantages of this method of treatment.

On the other hand, by irradiation the penis is preserved, and though stricture of the external meatus may sometimes result, its frequency is not comparable to that observed after amputation. If irradiation fails, we can still fall back upon amputation, with the knowledge that delay in surgical removal has not eliminated the hope of cure.

OPTIC ATROPHY, WITH RETINAL CHANGES, CAUSED BY HIGH-TENSION CURRENT.

BY

W. BAINBRIDGE, M.B., CH.B.ED., D.P.H.,
HONORARY OPHTHALMIC SURGEON, HULL ROYAL INFIRMARY;
OPHTHALMIC SURGEON, HULL EDUCATION COMMITTEE.

I PUBLISH the following case because similar cases must be rare, and it deserves to be recorded. The history was given to me as follows.

In June, 1927, the patient, aged 31, an acetylene burner by trade, was going into an electric crane to do some repairs when he happened to take the wrong lead to the crane and got a shock with a current of 460 volts A.C. He was taken to the local hospital. His left wrist was burned, and also his toes. His

The condition of his eyes, when first seen, was as follows: Pupils active to light and accommodation. Conjunctivae and corneae normal. Right eye, vision 6/36. The fundus examination showed the disc to be pale and atrophic, and the arteries slightly smaller than normal. There were a large number of rounded white areas in the macular region surrounded by pigment. To the nasal side of the disc there was an angioid streak. The retina appeared healthy in the periphery. Left eye, vision 6/24; upper part of the disc pale. There were similar rounded areas to those seen in the right eye, and angioid streaks on either side of the disc. The periphery showed retinal degeneration. A small degree of myopic astigmatism was present in the right eye, and a small degree of myopia in the left. Both fields of vision contracted all round.

Wassermann reaction negative; urine normal. He was examined by Dr. Muir, honorary physician to the Hull Royal Infirmary, who reported that he could find no disease of the central nervous

Right eye.

Left eye

eyes were bloodshot for four months. He received no treatment for them in the hospital because he did not complain about them; they did not hurt him, and his sight was not affected. He was in the hospital for a month or five weeks. Six or seven months afterwards his eyes became misty and used to hurt him. He went to his doctor about them, who advised him to see the nearest eye specialist, who tested his eyes and prescribed glasses. As his eyes still troubled him he again visited his doctor, who sent him to the Hull Royal Infirmary, where I first saw him on October 8th, 1929. Regarding the family history, he has three sisters and five brothers, all with good sight; his father and mother have also good sight.

system. I saw him again on October 15th, 1929, January 10th, 1930, and May 13th, but the condition of his eyes has remained the same.

As the case was one of peculiar interest, I sent him up to London to see Mr. A. H. Levy, surgeon to the Central London Ophthalmic Hospital, who reported as follows:

"The fundi show most peculiar changes. Besides the optic atrophy, which is well marked in the right, there are macular changes, and also vascular changes. The brown lines look somewhat like angioid streaks, except for their colour, and the whole gives me the impression of a very slow non-infective inflammatory process."

Mr. Levy considered that the condition had definitely been caused by the accident.

Ocular injuries through electricity are not common, as shock of sufficient severity to produce fundus changes is generally fatal. The modes of injury are as follows: lightning, exposure of the eyes to electric flash, and electric shock.

The following fundus changes have been reported:

(a) Hyperaemia, oedema, changes in calibre of arteries and veins.

(b) Localized lesions, including scar formation observed round disc (Terrieu), near macula (Haab). Hole at macula with pigmentation at periphery (Kramer).

(c) Fully established choroido-retinal changes, with and without gross interference in visual acuity (Lundsgaard, Junius, Schirmer, Robinson, and Terrieu).

(d) Choroido-retinitis leading to detachment (Lundsgaard).

(e) Detachment in one eye and optic neuritis in the other (Pfahl).

(f) Optic atrophy (Ulbrich, Haas, Junius, Dor).

I am greatly indebted to Mr. Levy for his assistance in looking up the references to reported cases; also to Messrs. Rayner (Hull) for making drawings of the fundi reproduced.

BIBLIOGRAPHY.

- Wagenmann, A.: *Die Verletzungen des Auges, Graefe-Salmisch Handbuch*, 1913, second edition, Part V, p. 1656.
Terrieu, M. F.: *Les troubles visuels provoqués par l'électricité, La Presse Méd.*, 1912, p. 193.

MAJOR DERMATOSES OF CHILDREN.

BY

W. J. O'DONOVAN, M.D.,

PHYSICIAN, SKIN DEPARTMENT, LONDON HOSPITAL.

A "dermatological bad bargain" must be our first consideration. Chronic seborrhoea capitis is not dramatic, it is unpleasant to look at, inveterate in its resistance to ordinary treatment; it is a bar to the education of its owner, a hindrance to its possessor's marriage or employment, and, throughout its course, a means of reproach to the profession. I shall first describe the clinical picture as we see it in young children, outline its clinical variations and complications, and suggest a line of treatment that I have most frequently found to be efficacious.

CHRONIC SEBORRHOEA CAPITIS.

The disease appears, when developed in a child of 5 or more years of age, as a scalp condition whose history is seldom definitely obtainable. The scalp is red or pink, scaling all over, and generally dry. Moist patches from exudation of clear serum are covered with heaped-up minor aggregations of damp, yellowish scales. Over the greater part of the head the pink surface is dry, and covered with copious, easily detached white scales. The hair has no lustre; its surface, if dark, is greyed. Generally the hair is plentiful, but at times in an exacerbation of the disease there may be a general loss of hair producing notable alopecia; regrowth is the rule. Were the condition confined to the scalp it would be bad enough, but extensions on to the body are frequent, particularly to the flexures. The folds of the groins, the umbilicus and the armpits, although hairless, are red, and covered with copious, easily detached white small scales. At times all these sites may be moist, exude freely, and show scabs and macerated epidermis, heaped over the reddened surface as a pale yellow malodorous and pultaceous accretion. During acute phases it is common for a moderate adenopathy to develop in the neck and groins. These children otherwise are well, but particular attention should be called to a hall-mark of skin chronicity which they invariably show. The eyelids are reddened at their margins, while yellowish crusts are formed at the roots of the eyelashes. The lids are pink and roughened from frequent rubbing, matted bunches of eyelashes are often seen, and a considerable disfiguring loss of eyelashes takes place also. In about one-third of the cases two further extensions can be seen. First, the eyebrows may be involved; their site is reddened and scaling, or moist and scabbed. Secondly, behind the ears

red fissures may develop; the surface of the ears is pink, the whole ear is thickened, while stiff and macerated scales and caked exudate often cover a retained yellow, sticky, purulent discharge, that has been formed at the crescent of attachment of the auricle to the scalp. Both sexes are afflicted with an equal incidence. The disease is far more common in hospital and institutional work than in private practice. For this, the factors of heredity, uncleanness, a disinclination on the part of the parents to perform unpleasant dressings, and improper feeding, must be taken into account. Although superficial pustular dermatitis speedily clears up when removed into the best hospital surroundings, nevertheless the scaling erythema of scalp and flexures remains, apparently responding to the varied treatment of the house-physician, but, quite unaccountably, exacerbations occur that bear no relation to changes in diet, climatic changes, or to variations in the general health. It is soon borne upon the observer that these patients have a fragile skin; they are photosensitive, and exposure to the sun on a bright day makes pink skin red, and red areas moist. These patients respond by aggravation if they are washed with any household soaps, or if mercurial ointments are applied in an attempt at antibacterial treatment. Such children are not of the nervous type so brilliantly described by Cameron. They do not suffer from dental sepsis, there are no signs of intestinal toxæmia, and their differential blood count is normal. They are not anaemic, or wasted, or mentally dull, although they have a strange tolerance of their persistent malady.

In treatment, personal cleanliness is of the first importance; underclothing and bed linen should be changed and boiled daily. We can assume that by the time these cases have reached the specialist's hands, bland treatments, such as bran baths, calamine liniment, various zinc ointments, soft paraffin dressings, a period of salt-free and also of low sugar dietaries, have already been tried without success. Thyroid extract and ergosterol may have been exhibited, and yet the disease remains. My practice is to order an x-ray epilation to the whole scalp at once. If the child is an inmate of an institution, or under the care of an authority through which such a treatment is not easily obtainable, I prescribe thallium acetate, 8.5 milligrams per kilo of body weight, dissolved in a draught of sweetened water. Whilst awaiting the epilation that should occur after a latent period of three weeks, I swab the scalp and any other lesions with a 2½ volume peroxide lotion. This is speedily mopped away with boiled water, and the scalp and lesions are then sprayed with Sir Lenthal Cheatle's malachite green, which is a solution composed of equal parts of 2 per cent. malachite green in 80 per cent. alcohol, and 2 per cent. mercuric chloride in 80 per cent. alcohol. This application is, to some patients, painful for a short time at first, but it is an invaluable aid for infected dermatoses. The scalp should be cropped closely before the application of x rays or giving of thallium, so that local medication may be facilitated. The backs of the ears, the axillae, and the groins should also be sprayed. Frequent application of white lint fomentations for twenty-four hours, or a starch poultice, is occasionally needed before a direct attack can be instituted in this manner. Application of unguentum hydrargyri oxidi flavi to the eyelids twice a day is also advisable. A watch must be made for bed-wetting, and naturally constipation will receive proper treatment. It is a common tradition that soap and water is bad for these patients. Hard water and common soaps undoubtedly are, but I have never regretted bathing these patients daily, using Midgley's or Allen and Hanburys' basic over-fatted soaps; a second water is always used, however, to remove any curds from the skin, and hot, soft towels are advisable to lessen the need of skin friction and to shorten the washing process, since, in winter, these children frequently develop attacks of bronchitis.

During treatment, patients are kept in bed, and at the last stage in the latent period after x-ray treatment, it is highly advisable to use the weapon of protein shock therapy. This is best given on the nineteenth evening after the x-ray treatment, so that the period of fever corresponds to the time of epilation. The customary dose