A CASE OF HUMAN INTESTINAL MYIASIS.

BY

J. F. D. SHREWSBURY, M.B., CH.B., D.P.H.,
LECTURER IN BACTERIOLOGY, UNIVERSITY OF BIRMINGHAM.

(With Special Plate.)

REPORTS of cases of myiasis—infestation with fly larvae—are infrequent in the medical literature of this country. Possibly the incidence of infestation is higher than the published reports suggest, especially in regard to intestinal myiasis, since in those cases in which larvae are only present in small numbers the condition may pass unrecognized.

The case here reported is that of a little girl, aged 3 years, a doctor's daughter. The child suddenly developed a mild enteritis, which began with an attack of vomiting and persisted for about a fortnight. During this period her father noticed, on two separate occasions, a single larva in her motions. Eventually, some three weeks after the onset of her illness, two more larvae were found in the chamber one morning, immediately after the child had urinated. No faeces were passed on this occasion, but, in view of the previous findings, the larvae were most probably passed per anum. There were never any symptoms of the urethral or vaginal irritation that would certainly have accompanied the presence of fly larvae in the urethra or the vagina. No more larvae were evacuated, and the child made a speedy and complete recovery from her enteritis.

The larvae were originally sent to Dr. P. C. Cloake, honorary assistant physician to the Queen's Hospital, Birmingham, who brought them to me. I, in turn, sent one larva to Major E. E. Austen, keeper of entomology, British Museum (Natural History), who identified it as the larva of the dipterous fly, *Anisopus fenestralis*, and generously gave me permission to incorporate his report, which follows:

Anisopus (Rhyphus) fenestralis, Scopoli.

"This insect breeds normally in fermenting organic matter, such as cow dung, the effluent from farmyard manure heaps, sap exuding from wounds in trees, water in tree holes, and the decaying roots of certain plants. In one instance the species was found breeding in a human liver, which had been exposed to the air after having been preserved in 10 per cent. formalin for seven years; the larvae have also occasionally been found in cider and home-made wine, and in one instance, they caused damage to the comb in a beehive. As its name implies, the adult insect, which, although quite incapable of biting, somewhat resembles a mosquito in appearance, is often to be found on the windows of living rooms."

The two larvae I received from Dr. Cloake were unequal in size, the larger one being about half as big again as its companion. The larger larva measured exactly 1 cm. in length by 0.5 mm. in breadth. The larvae were, however, identical in structure and colouring.

I am indebted to the courtesy of Mr. F. W. Edwards, Major Austen's colleague, for the following expert description, by which the larva of *Anisopus fenestralis* may be recognized.

"Complete absence of legs, but distinct brown head present. Body consisting of three thoracic segments and ten abdominal. The thoracic segments show conspicuous reddish-brown markings on the sides, and less conspicuous bands are present on the abdominal segments. The first eight abdominal segments each consist of a narrower portion at the base, divided by a suture from a larger posterior portion. The respiratory system has two pairs of openings, one on the first thoracic segment and the other at the extreme tip of the body; the latter is surrounded by five small pubescent lobes. The last two abdominal segments are small. The anus is surrounded by a plate on the eighth segment."

In human intestinal myiasis, infestation can occur apparently by one of two routes: either the ova of the offending fly are ingested and pass through the stomach unharmed, the larvae hatching out in the intestines, or the fly actually oviposits on or near the anus of an individual at stool. The ova hatch rapidly in this situation, and the larvae make their way through the anus and travel upwards into the large bowel. This path of infestation is probably not uncommon in districts where the old earth closets are still in use (Hewitt, 1912), since many of the

dipterous flies that provoke myiasis are attracted to sewage and the females oviposit on faecal matter. In the case reported, however, infestation most probably resulted from ingestion of the ova. Little children frequently suck their fingers, and are also fond of rubbing their fingers up and down window panes. As Major Austen has pointed out, *Anisopus* is commonly found on the windows of living rooms, and oviposition is certain to occur occasionally on window frames when the fertilized female is unable to escape from the room to find suitable faecal or vegetable material.

Human intestinal myiasis in this country, judging from the published cases, is not apparently of grave import as a general rule. Enteritis, of a more or less serious degree, invariably accompanies the passage of the larvae through the intestinal tract, but the symptoms clear up rapidly after the larvae have been evacuated. The larvae do not appear to attack the intestinal mucosa, subsisting on the food material available in the faeces. When the larvae are present in large numbers, however, the symptoms are proportionately more severe and the patient may be seriously ill. In the adult, the passage of fly larvae per anum is at least an unpleasant, and frequently a terrifying experience, and in the published cases the psychological effect upon the adult patients is usually evident. There is, moreover, always the risk that the larvae may be carrying intestinal pathogenic bacteria, or may pave the way for an infection with such organisms.

Treatment does not seem to exert much effect upon the larvae, which leave of their own accord, per anum, when approaching maturity. The usual vermifuges are of little value, but intestinal disinfectants, mild purgatives, and high enemata may hasten their departure in certain cases.

I wish to thank Drs. W. R. Blore and P. C. Cloake for their permission to publish the clinical material, and to express my grateful appreciation of the generosity of Major Austen and Mr. Edwards in allowing me to utilize their reports.

LITERATURE.

Hewitt: "An Account of the Bionomics and the Larvae of the Flies *Fannia (Homalomyia) canicularis* L. and *F. scalaris* Fab., and their Relation to Myiasis of the Intestinal and Urinary Tracts." Reports to the Local Government Board on Public Health and Medical Subjects (New Series, No. 66). Further Reports (No. 5) on Flies as Carriers of Infection. London: H.M. Stationery Office. 1912.

"FAT NECROSIS" OF THE BREAST.

BY

H. A. COOKSON, M.B., F.R.C.S. Ed.,
PATHOLOGIST, ROYAL INFIRMARY, SUNDERLAND.

(With Special Plate.)

DRS. LEE AND ADAIR¹ were, I believe, the first to describe this condition, in a report to the American Surgical Association in 1920, when they described two cases. The same authors reported three further cases in 1922,² making five in all. A sixth case was recorded by Mr. Geoffrey Keynes³ in the *British Journal of Surgery*, 1924-25, and a seventh case was the subject of a contribution by Professor Hadfield⁴ in the *British Journal of Surgery*, 1925-26. Professor Hadfield refers to Mr. Rowntree⁵ as having described a case of this nature in the *Lancet* in 1925. The condition therefore appears sufficiently uncommon to be an excuse for describing another case of the condition.

Case Report.

A woman, aged 56, was admitted to the Royal Infirmary, Sunderland, complaining of a tender lump in the left breast. Eighteen months before admission she had fallen and injured the breast, and six months after her accident she had noticed a lump. This lump had recently slightly increased in size, and there was slight pain. On examination it was seen that the breasts were large and pendulous, and that there was a small nodule at the periphery in the lower and medial quadrant of the left breast. The skin was of normal appearance over the tumour, but was adherent to it. There was free movement on underlying structures, and no glands were felt in the axilla. The lump was tender on pressure, but felt firm; it was softer in the middle than at its edges. Mr. Hamilton Ross, the surgeon in charge of the case, excised the nodule by an elliptical incision and using a diathermy knife, after which the specimen was sent to the pathological department for examination.

W. ARNOTT DICKSON: PLEURAL EFFUSIONS IN GENERAL PRACTICE.

Empyema cured by absorption in boy aged 6.

Fig. 1.—May, 1929.

Empyema cured by absorption in boy aged 6.

Fig. 2.—After recovery.

C. ALEX WELLS: APPENDIX CONCRETIONS
OPAQUE TO X-RAYS.

X-ray photograph, showing shadow of stone at site of the appendix,
(from behind).

J. F. D. SHREWSBURY: A CASE OF HUMAN
INTESTINAL MYIASIS.

Microphotograph of the head and anterior segments of the larva of
Anisopus (Rhyphus) fenestralis, Scopoli ($\times 40$).