FIG. 1.—Case I, tuberculosis. Before thoracoplasty.

FIG. 3.—Case II, bronchiectasis. Before thoracoplasty; lipiodol injections demarcating cavities in the right lung.

FIG. 2.—Case I. After thoracoplasty. (Sides reversed in photograph.)

FIG. 4.—Case II. After thoracoplasty; lipiodol injection showing collapse and flattening of peripheral and central cavities; those near the mediastinum are only partially collapsed.

W. E. COOKE: THE MACROPOLYCYTE.

FIG. 1.—Normal polymorph. ($\times 1000$.) FIG. 2.—Macropolycyte from a case of staphylococcal osteomyelitis. ($\times 1000$.) FIG. 3.—Macropolycyte from a case of streptococcal septicaemia. ($\times 1000$.) FIG. 4.—Macropolycyte from a case of pernicious anaemia, showing hypersegmentation of nucleus. ($\times 1000$.) FIG. 5.—Macropolycytes of the megakaryocyte type in pernicious anaemia. ($\times 1000$.) FIG. 6.—Macropolycyte of the megakaryocyte type showing large granules in the cytoplasm. ($\times 1000$.) FIG. 7.—Macropolycytes of the megakaryocyte type in pernicious anaemia. ($\times 1000$.)

7. Pneumectomy appears to be almost totally contra-indicated in phthisis. In bronchiectasis, although apparently the ideal procedure, it is followed at present by an almost prohibitive mortality (50 per cent.). Further improvements in the technique may eventually make it the operation of choice. The more recent cautery pneumectomy, as practised by Graham in America, has been followed by a much lower mortality than any other lobectomy operation, and offers reasonable hope of eradication of the septic lobe.

My thanks are due to my medical colleagues of the staff of the Brompton Hospital for the opportunities afforded me in this work, and to Dr. Stanley Melville for the very excellent skiagrams, a few only of which are reproduced here.

THE MACROPOLYCYTE.

(With Special Plate.)

BY

W. E. COOKE, M.D., F.R.C.P.ED., M.R.C.P.LOND.,
D.P.H.

DEFINITION.

THE diameter of a normal polymorph in stained films is from 10 to 12 microns. The term "giant" polymorph or macropolyocyte—a name suggested by Sir Humphry Rolleston—may be applied to those polymorphs having a diameter of more than 14 microns. Fig. 1 represents a normal polymorph with five nuclear segments. This has a diameter of 11.5 μ .

ETIOLOGY.

The macropolyocyte is found in two widely different conditions—such opposite conditions, in fact, as to make one doubt whether the causation can be the same in the two cases.

The first is in the acute pyogenic infections. Fig. 2 illustrates a macropolyocyte from a case of staphylococcal osteomyelitis showing the following blood picture:

Total leucocytes	...	...	18,000	per c.mm.
Polymorphs	...	...	84	per cent.
Lymphocytes	...	...	12	"
Large mononuclears	...	...	3.5	"
Myelocytes	...	...	0.5	"

Polynuclear count:

I.	II.	III.	IV.	V.
41	43	15	1	0 per cent.

Fig. 3 illustrates a "giant" in a case of streptococcal septicaemia.

Total leucocytes	...	...	26,000	per c.mm.
Polymorphs	...	...	79	per cent.
Lymphocytes	...	...	13	"
Large mononuclears...	...	...	7.5	"
Myelocytes	...	...	0.5	"

Polynuclear count:

I.	II.	III.	IV.	V.
34	43	21	2	0 per cent.

The staining reactions of the nucleus and cytoplasm are identical with those of other polymorphs in the films. They are rare—encountered perhaps twice or three times a year by anyone constantly examining blood films—and seem always to be associated with a polymorphic leucocytosis and the presence of myelocytes in the peripheral blood. They are, I think, "sports."

The other condition in which macropolyocytes occur is pernicious anaemia, and in this disease they are frequently seen. They measure 16 to 21 microns in diameter and are of two distinct types. The first type of macropolyocyte is similar in nuclear structure and cytoplasm to the other polymorphs, but the cell is larger, and the nucleus is frequently hypersegmented, as shown in Fig. 4, which illustrates this type. In the second pernicious anaemia type the total bulk of the nucleus is great, absolutely and relatively to the amount of cytoplasm. The nucleus may be of the band form, or have one, two, or as many as fourteen divisions. The basi-chromatin is not so densely massed as in the normal cell, and the cytoplasm contains larger granules, which are neutrophile in reaction. Figs. 5, 6, and 7 illustrate this type. These cells look very like megakaryocytes.

The etiology of the macropolyocyte in pernicious anaemia is probably that of the disease itself. The polymorph, even in the absence of the "giant," behaves in a curious manner.¹

The normal polynuclear count may be taken to lie between the following values:

	I.	II.	III.	IV.	V.
(1) ² :	11	25	47	15	2 per cent.
or (2) ³ :	5	35	41	17	2 per cent.

which give readings of 36 and 40 in the left-hand columns and 64 and 60 in the right.

In infections the count moves to the left, as in the cases of osteomyelitis and septicaemia quoted above:

I.	II.	III.	IV.	V.
41	43	15	1	0 per cent.
34	43	21	2	0 "

In pernicious anaemia frequently, possibly in all uncomplicated cases at some period of the disease, the polynuclear count shows an increase of the cells of Classes 4 and 5, as in the following examples:

	I.	II.	III.	IV.	V.
Case 1 ...	8	20	39	22	11 per cent.
Case 2 ...	6	27	40	22	5 "
Case 3 ...	7	20	44	24	5 "
Case 4 ...	7	16	29	39	9 "
Case 5 ...	12	34	26	20	8 "
Case 6 ...	10	15	38	24	13 "
Average...	8	22	36	25	9 "

The left-hand columns give an average of 30 and the right 70. The count is actually right-handed. As far as my experience goes, this does not occur in any other disease.

We have, then, at the outset in pernicious anaemia some factor producing a dysleucopoiesis, and accentuation of this may lead to the production of the macropolyocyte.

Some recent papers on the transformation of monocytes in frog's blood, and cells of lymph nodes, into giant cells are suggestive. M. R. Lewis⁴ states that giant cells are formed from monocytes in hanging drop preparations of frog's blood. E. Barta,⁵ working on tissue cultures of lymph nodes, finds that giant-cell formation takes place at some distance from the surface, and ascribes this to deficiency of oxygen. The addition of embryonic extracts greatly diminishes the effects of oxygen want, and cells in the deeper layers then behave like those with a sufficient oxygen supply. There are many more factors in experimental conditions than simple oxygen want. Alteration in pH, gaseous diffusion, surface tension, saline ratios, and lipid changes must take place and also influence cell metabolism.

Anoxaemia is not the only state responsible for the production of macropolyocytes in pernicious anaemia, but whatever the primary general state in this disease is, the marrow seems to take control in the later stages. In addition to the behaviour of the polynuclear count, and the presence of macropolyocytes and ichthyoid cells, there is direct proof of this local influence. Walterhöfer and Schramm⁶ have scraped and irrigated the medullary cavities of long bones in cases of pernicious anaemia that were apparently moribund. The alteration of the local chemical state effected produced surprisingly good results. That the results may be only temporary does not affect the argument.

These considerations lead to the conclusion that the macropolyocytes and pernicious anaemia may be due, either to a primary defect in the chemical composition of the blood, the marrow changes completing the vicious circle, or to a congenital or an acquired condition of the haemocyto-blast itself leading to defective haemopoiesis and local biochemical abnormalities, with, later, general effects.

PROGNOSTIC IMPORT.

The first type of macropolyocyte, that found in pyogenic infections in association with myelocytes, is, I think, an "accident" and has no prognostic value, except, perhaps, as signs of marrow "stress." Of the second type, the first subdivision, an exaggerated form of the polymorph frequently found in pernicious anaemia, is present at the crises of the disease, and is an index, I think, of local urgency in the haemopoietic tissues, which may or may

not end in temporary recovery. The presence of the second, or megakaryocyte type, is of grave significance, and no case in my experience has lived beyond a few days from its appearance.

REFERENCES.

- ¹ Cooke, W. E.: *The Arnetb Count*, 1914, 19. ² Cooke, W. E.: *Journ. Path. and Bact.*, xix, 496. ³ Arnetb, J.: *Deut. med. Woch.*, Leipzig, 1904, Bd. xxx, S. 54. ⁴ Lewis, M. R.: *Arch. exper. Zellforsch.*, 1926, 2, 228-243. ⁵ Barta, E.: *Ibid.*, 1926, 2, 6-30. ⁶ Walterhöfer, G., and Schramm, O.: *Arch. f. klin. Chir.*, 1921, cxviii, 794-811.

THE EXPERIMENTAL BASES OF PROPHYLAXIS IN GONORRHOEA.

BY

KENNETH M. WALKER, M.B., F.R.C.S.,

LECTURER ON VENEREAL DISEASES, ST. BARTHOLOMEW'S HOSPITAL.

THE study of the initial stages of gonococcal urethritis is of extreme importance from the point of view of prophylaxis. Knowledge of the pathology of acute gonococcal urethritis is almost entirely derived from the observations of Finger, who inoculated moribund patients with gonococci and made a detailed examination of their urethrae thirty-six to forty-eight hours afterwards. His description is as follows:

"The gonococci begin to proliferate on the surface of the epithelium which they are about to attack; they are not spread diffusely, but in small groups. At the end of thirty-eight hours they have scarcely as yet penetrated into the epithelium; at the most they have slipped in among the superficial coils at the points presenting the feeblest resistance. If the epithelium is denuded at any point, exposing the connective tissue, the organisms penetrate there more rapidly. They invade the lumina of the crypts of Morgagni with extreme rapidity, and are found in them, lying free on the surface of the epithelium. From this moment, and, indeed, well before this, diapedesis has commenced, the leucocytes are everywhere numerous in the superior layers; a certain number have gained the surface, and one already finds intracellular gonococci."

Finger's observations would lead us to believe that until at any rate thirty-eight hours after inoculation very little penetration of the epithelium by the gonococci has taken place, and that the adoption of appropriate prophylactic measures would stand every chance of success if carried out within this period.

Experimental work has, however, been carried out by other observers using organisms other than gonococcus, which suggests that the diffusion of organisms within the urethra might be more rapid than Finger's observations on the gonococcus would lead us to suppose. Notable among these experiments are those of Palladino Blandini. This observer placed a loopful of *Bacillus prodigiosus* within the meatus of a guinea-pig, and twelve hours later made a necropsy under aseptic conditions and investigated the fate of the organisms since their introduction into the urethra. His choice of *B. prodigiosus* was determined by the fact that on account of its brilliant colouring matter it could easily be recognized. Palladino Blandini found that a widespread diffusion of prodigiosus from the urethra had taken place within twelve hours of inoculation, but that if the animal were allowed to survive for more than twenty hours these diffused organisms were destroyed by the tissues. That the method of diffusion was not through the blood stream was proved by the fact that blood cultures taken from the heart were sterile. In 1909 I repeated and confirmed these findings, and four years later Teale and Embleton proved that what occurred in the urethra was equally true of organisms placed on other mucous surfaces or inoculated into serous cavities. The prodigiosus is capable, therefore, of passing through uninjured mucous membrane, and may be carried from thence along the lymphatic stream and eventually into the thoracic duct. Teale and Embleton's experiments showed, therefore, as mine had already done, that with regard to the generative tract the spread of organisms is into the submucous tissues rather than along the surface. At the necropsy of a guinea-pig killed twelve hours after a loopful of organisms had been placed within the meatus no infection of the surface of the posterior urethra, bladder, ureter, or pelvis was found, although extensive dissemination by the lymphatics had taken place. The organism employed for this experi-

ment was, however, *B. prodigiosus*, and it is possible that what is true of that organism is not true of the gonococcus. *B. prodigiosus* is a non-pathogenic organism, whereas the gonococcus, at any rate in the human being, provokes immediate and intense reaction of the tissues. Nevertheless, on repeating this experiment with a mixture of *Staphylococcus aureus* and diphtheroids I obtained similar results, and these organisms, unlike *B. prodigiosus*, are pathogenic to guinea-pigs.

In the light of these experiments Finger's contention that there occurs little or no penetration of the human urethral mucous membrane by the gonococcus for thirty-eight hours seemed open to doubt, and for many years I awaited an opportunity of throwing new light on the situation. Would organisms placed within the meatus of a human being behave in the same way as organisms inoculated into the urethra of a guinea-pig? Two years ago a chance of carrying out a somewhat similar observation on a human being presented itself.

A patient was admitted under my care into the Royal Northern Hospital suffering from an impassable stricture in the bulbar urethra and an early carcinoma of the glans penis. At 11 a.m. I placed two loopfuls of *B. prodigiosus* within his meatus. At 6 p.m. I opened the urethra on the bladder side of the stricture without having previously attempted to pass any instrument. After dividing the stricture through the wound in the perineum, a sterile cotton-wool swab on a holder was passed along the urethra from the opening in the perineum to a point 2 inches from the meatus and an agar plate inoculated. A second swab was similarly passed until it reached a spot 1 inch from the meatus, and another plate inoculated. My object in doing this was to ascertain whether in the seven hours' interval the organism had spread along the surface of the urethra towards the bladder. The fact that both plates remained sterile showed that no surface spread had occurred. I then carried out a partial amputation of the penis. Later multiple transverse sections were made through the amputated penis. These sections were stained with methylene blue and carefully examined under an oil immersion objective for organisms within the tissues. On the surface of the urethra no organisms could be found, but in the submucous tissues large clumps of small bacilli morphologically resembling *B. prodigiosus* were visible. These collections of bacilli were either in the form of more or less circular colonies or else were spread out along a line, and although no proof could be furnished of this, their distribution suggested that they were lying within the submucous lymphatics.

This observation, therefore, so far as it went, led me to believe that what was true of the behaviour of *B. prodigiosus* within the urethra of a guinea-pig was equally true of its behaviour when placed within the meatus of a human being.

By the courtesy of Dr. Mervyn Gordon I was able to repeat my observations in a monkey. Similar results were obtained.

A loopful of Hoffmann's bacillus and *Staphylococcus aureus* was placed within the meatus, care being taken to inflict no damage whilst the inoculation was being made. Eight hours later the monkey was killed by means of chloroform. Transverse sections were then prepared through the urethra at different levels and organisms searched for. In this case a few Gram-positive bacilli and cocci were found lying in the lumen of the urethra and also deep to the epithelial cells, but further up, an inch from the meatus, larger clumps were found in the submucous layers.

The above experiments suggest, although they do not prove, that in any infection of the urethra there is an immediate dissemination of organisms through the lymphatics as well as a slower spread which is associated with the signs of inflammatory reaction. In the majority of cases the tissues are able to deal with the immediate lymphatic invasion provided that it is not too liberally reinforced by fresh supplies from the surface. The chief point of practical interest is the bearing of these observations on prophylaxis in gonorrhoea.

Clinical observations, and particularly those carried out at the prophylactic stations established by the American Army in Paris during the war, showed that the chances of success of any prophylactics are directly proportional to the time that has elapsed since exposure. Since the gonococcus is easily destroyed by weak disinfectants while lying on the surface of the urethra, irrigation with weak permanganate solution is found to be effective in the great majority of cases, provided the patient presents himself for disinfection within a few hours of exposure. The longer the interval between exposure and disinfection, the more often does the disinfection process fail. No absolute time limit, however, can be laid down to guide us in arriving at a decision as to