triple rhythm, daily range of temperature 97° to 98.4°, is illustrated in Fig. 2. The diphasic ventricular complex is shown with characteristics already noted. The initial phase of ventricular activity is necessarily prolonged, for the stimulus has first to traverse the left ventricle before reaching the right.

A record closing five months' observation is shown in Fig. 3. The tracing shows reversion to the physiological type, but the steep negativity of r_2 and r_3 is ominous.³

Of three cases encountered two were males. All had reached the degenerate period of life, with thickened arteries and high systolic and diastolic readings, and with symptoms indicative of cardiac insufficiency. The associated cardiac lesions were respectively mitral, aortic (Wassermann reaction positive), and hypertrophy (with renal cirrhosis), the common factor being myocardial with degenerative arterial change.

Branch block adds to the gravity of prognosis, as the much quoted statistics of Willis⁵ show. In an analysis of twenty-four cases of branch block, Cowan and Bramwell⁴ found that in only one case did the block show a tendency to disappear. Lewis,⁶ Ritchie,⁷ and Hewlett⁸ have recorded cases in which block was present as a temporary manifestation in association with febrile or toxic disturbance, as in the case here quoted. In Hall's case⁹ the block was permanent but stationary, in contrast with the ordinary rule that the lesion, if chronic, is progressive.

I wish to acknowledge my indebtedness to Dr. Foggie, in whose ward the patient was.

REFERENCES.

- ¹ *Clinical Electrocardiography*, p. 28. ² Gross: *The Blood Supply of the Heart* (quoted in ⁴). ³ MacLwaine and Campbell: *BRITISH MEDICAL JOURNAL*, 1923, ii, 456. ⁴ *Quart. Journ. Med.*, No. 73, October, 1925. ⁵ Willis: *Clinical Electrocardiography*. ⁶ *BRITISH MEDICAL JOURNAL*, 1913, i, 484. ⁷ Cowan and Ritchie: *Diseases of Heart*. ⁸ *Heart*, 1921, 9, 1. ⁹ *BRITISH MEDICAL JOURNAL*, 1924, i, 779. ¹⁰ *Presse Méd.*, March 20th, 1926.

TUBERCULOSIS AND MILK.*

BY

NATHAN RAW, C.M.G., M.D.,

LORD CHANCELLOR'S VISITOR IN LUNACY.

IN the *BRITISH MEDICAL JOURNAL* of 1903 I ventured to suggest that bovine bacilli conveyed by milk played an important part in causing tuberculosis in children. This has since been confirmed by many other workers, and is now accepted as a serious menace to child life.

The International Congress on Tuberculosis which was held in Washington in 1908 will always be remembered on account of the discussion which took place on the part played by the bovine bacillus in the production of tuberculosis in man. We were honoured on that occasion by the presence of the great master, Robert Koch; and those of us who took part in the secret session will never forget it. Up to that time Koch had not publicly admitted that bovine infections in man were possible, but he later modified his views, and we are all now agreed that an important amount of tuberculosis, especially in children, is directly conveyed from the dairy cow.

At that conference I read a paper, entirely based on clinical and hospital observation, pointing out the possibility that surgical tuberculosis in children, such as enlarged lymph glands, mesenteric disease, and affections of bones and joints, were separate and distinct infections from those of pulmonary tuberculosis, and might be caused by milk. The brilliant pioneer work of our president, Thebold Smith, on the role of the bovine bacillus, stimulated many workers, myself included, to further investigation, and it is particularly fitting that, eighteen years afterwards in this city of Washington, Dr. Park should tell us of the immense reductions in infant mortality and suffering as the result of their researches.

The incidence of bovine tuberculosis varies in different countries, and depends upon the extent to which cattle are infected with tuberculosis and the amount of uncooked milk, butter, cheese, and other dairy products consumed. In England and Scotland dairy cows are very heavily

affected, with the result that surgical tuberculosis is very common in children.

In the light of our present knowledge we cannot say with certainty which particular strain of bacillus has caused the infection. We are accustomed to recognize three types of tubercle bacilli: human, bovine, and avian. I feel convinced that there are several intermediate strains of bacilli, which we cannot yet recognize with certainty, and it is not possible to say without bacteriological and animal tests to which group those from any lesion belong. For this reason, for the purpose of determining milk infections, clinical and hospital evidence is valuable.

Pulmonary tuberculosis is in a class by itself, and is practically always caused by the human type of bacillus, which also causes laryngitis, secondary intestinal ulceration, and fistula. It is rare for such cases to develop any other lesions. On the other hand, bovine infections in human beings rarely cause pulmonary lesions; in fact, they seem to be antagonistic to each other, and it is possible that a primary bovine infection, such as enlarged neck glands, may produce immunity against pulmonary tuberculosis for life.

Should this theory prove correct, the beneficial effects of sterilization of tuberculous milk may be followed by diminished resistance to human infection in the form of pulmonary tuberculosis. In any case it is imperative that the milk supply be pure and safe for children.

Avian bacilli, though considered to be non-pathogenic to man, may contaminate a milk supply, but with what effect we do not know. In Denmark the danger of infection of pigs by avian bacilli is so real that legislation has been introduced forbidding chickens on the farm.

To sum up. In my opinion—

Human bacilli produce:

1. Pulmonary tuberculosis.
2. Tuberculous laryngitis.
3. Secondary intestinal ulceration.
4. Fistula in ano.

Bovine bacilli produce:

1. Enlarged lymph glands.
2. Abdominal tuberculosis.
3. Lesions of bones and joints.
4. Meningitis.
5. Acute miliary tuberculosis.
6. Lupus.
7. Rarely, secondary extension to lungs.

These infections are antagonistic to each other, and human and bovine bacilli are rarely found in the body at the same time. The practical measures that should be taken against bovine infection are eradication of tuberculosis from dairy cows and pasteurization of all milk for human consumption.

Eradication will for a long time be impossible; pasteurization is a safe and certain remedy. The results in several large cities where compulsory pasteurization is law are truly wonderful, and have resulted in the saving of the lives of thousands of children every year, and endless pain and suffering with lifelong deformity. I sincerely trust the splendid pioneer work in New York and other American cities will soon be universal. At the same time it is imperative that every effort should be made to stamp out tuberculosis in dairy cattle.

It is estimated that in England about a million dairy cows are affected with tuberculosis. The cows with udder disease are, of course, the most dangerous. The compulsory seizure, isolation, and slaughter of these cows is a slow, laborious, and very costly procedure. The new Tuberculosis Order recently made in England will in time have a most beneficial effect in removing diseased cows from dairy herds, and so reducing the amount of bovine infection in children. In addition it is to be remembered that the financial loss to stock breeders and farmers from tuberculosis is enormous.

The question of the possible immunization or vaccination of cattle against tuberculous infection is of supreme importance, and as practically all cattle are born free of any tubercle-infection taking place after birth—the problem ought to admit of a solution.

I would like to lay before this conference the results of my own work, during the last eighteen years, in attempting to produce immunity in animals against tuberculosis. In 1906 Professor Koch gave me in Berlin a pure culture on

* Read before the International Conference on Tuberculosis, Washington, October 2nd, 1926.

glycerin agar of human bacilli from a virulent case of pulmonary tuberculosis. Professor Calmette, then of Lille, supplied me with a virulent bovine culture from the mesenteric glands of a cow, and Professor Bang of Copenhagen gave me a culture of avian bacilli from the liver of a chicken.

These cultures have been subcultured on glycerin potato and transferred to glycerin agar every month for twenty years in my laboratory, and the present growths represent the 241st generation. At the end of 1914 I was afraid the work would be interrupted, but I took the cultures with me to France, and as I had a good laboratory in the hospital I was able to continue the subculturing there for the next four years. The growths are still profuse and true to type, but they are now completely non-tuberculo-genic and non-pathogenic to animals.

Every year since 1906 I injected these bacilli into animals with a view of testing their pathogenicity; until 1913—that is, the 84th generation—I noticed no change in their virulence. After that time attenuation became marked, and in a series of inoculations of these bacilli into animals in 1914 they were observed to be avirulent. These experiments conclusively prove that virulent tubercle bacilli can be attenuated to such a degree as to be avirulent and non-pathogenic to highly susceptible animals. Some hundreds of animals have been inoculated to test the powers of immunization of healthy animals, and with excellent results. For the purpose of immunizing calves a vaccine was prepared from attenuated cultures of human bacilli. It is a bacillary emulsion of killed bacilli, and contains all the products of the bacillus. Living bacilli are never used; they are destroyed by heat before preparation.

Calves that are to be immunized should be separated from the mother at birth and fed on healthy milk for a week, when the first dose of vaccine is given; this is followed a week later by the second dose. They then associate with the other animals. The injection is given hypodermically in the usual way, under the skin of the abdomen, with strict antiseptic precautions. The method is only applicable, of course, to healthy animals.

During the last four years over 600 calves in valuable pedigree herds have been vaccinated by careful veterinary surgeons, and tuberculin tests at intervals of six months have so far been most satisfactory; the duration of immunity in 120 calves has been over three and a half years.

The problem of tuberculosis resolves itself into the question of immunity, and, in my opinion, we want to aim at producing an immunity in the animal against the amount of infection it is likely to encounter in ordinary life. It is almost impossible to immunize any animal against a lethal dose of virulent culture. The protection of the community by some safe scientific method of active immunization is the only effective way by which this preventable scourge can be eradicated.

Turning to the question of human immunity the problem is much more difficult, as children are susceptible to both human and bovine infection. At the special request of the parents in a large tuberculosis colony to which I was attached, and where each father has pulmonary tuberculosis, I gave all the children, over a hundred in number, two doses of protective vaccine prepared from dead attenuated bovine bacilli, with the object of protecting them from human infection. So far they appear to be healthy, but it will require at least ten years of observation before we can give any definite opinion as to its efficacy.

In conclusion, I am of the opinion that it is possible to immunize cattle against ordinary infection by using killed cultures of attenuated human tubercle bacilli, in precisely the same way that we can give immunity to man against ordinary infection of typhoid and paratyphoid by using dead cultures of typhoid bacilli. Animals cannot withstand large doses of virulent bacilli as a test of immunity, and I cannot see that it is necessary to give them. Our aim is to protect animals and children against the ordinary infection they are likely to meet with in life, and not against a lethal laboratory dose of virulent bacilli. In my opinion it is not necessary to use living organisms for the purpose of producing a relative immunity against tuberculosis.

TREATMENT OF TUBERCULOSIS.

A NOTE ON THE EFFECT OF COLLOIDAL ANTIMONY.

BY

PERCY MOXEY, M.B., CH.B., D.P.H.,

CLINICAL TUBERCULOSIS OFFICER AND DEPUTY M.O.H. FOR THE PORT AND BOROUGH OF SOUTHAMPTON.

PULMONARY tuberculosis is so widespread and accommodation in sanatoriums is so limited that any form of treatment to be of much value to the community at large should be one than can be carried out under ordinary conditions with the patient living at home, or even continuing his work. It should also be such as can be carried out by the patient's own medical attendant.

In looking for a drug which would at least alleviate some of the most distressing symptoms of phthisis it occurred to me to make trial of antimony, which has been used with some success in the treatment of leprosy, kala-azar, and bilharziasis. The difficulty at first was to find a preparation which caused no unpleasant symptoms. A 2 per cent. solution of tartar emetic in normal saline was tried for some time, half-grain doses of antimony tartrate being administered intravenously twice a week. Improvement followed a course of these injections, but the immediate effect was often a feeling of collapse.

I next decided to try colloidal antimony (Crookes's) in 1/2 c.cm. doses intramuscularly twice a week, and found that with this there was no shock, and it could be given in the out-patient clinic without fear of untoward symptoms. On the day after the injection the patient feels a little tired, but when this has passed off there is a distinct improvement; appetite increases, the breathing becomes easier, and the patient feels much brighter and more alert. The effects on the lesion seem to resemble those produced by sanocrysin, but without its dangers; for a week or two moist signs increase and the expectoration becomes more profuse, but these eventually diminish, the evening temperature drops slowly to normal, and thereafter pursues a much more level course.

The colloidal antimony is most conveniently injected into the flexor muscles of the forearm, and in some thousands of injections during the past year I have had no ill effects from it. There is often a little pain with the first few doses, but this is seldom complained of at subsequent injections. The dose at first is 0.5 c.cm., increased later to 1 c.cm. The temperature is as a rule improved after three or four injections, but if the pyrexia persists, both the quantity and the frequency of the injections may be increased. In severe cases I have given daily injections for a fortnight without ill effect. Occasionally when a patient has not reacted to intramuscular injections I have given them intravenously, with immediate result; on the other hand, intravenous injections not uncommonly cause some malaise.

How long the injections of antimony should be continued it is as yet too early to say, but in general one would suppose that they should be persevered with until the physical signs have cleared up.

To patients with troublesome cough during the day I give also a mixture thrice daily of vinum antimoniale 10 minims, oxymel scillae 1 drachm, in half an ounce of water. This reinforces the antimony administration, and also soothes the cough and assists expectoration. For the morning cough, often so distressing, I give asparagin 1 grain first thing, dissolved in a drachm of water; this loosens the tenacious sputum. At night I give the liquid extract of *Sanguinaria canadensis* in 2-minim doses in water to allay bronchial irritation and promote a good night's rest. The two drugs asparagin and sanguinarin have been used by me now for many months and in over 200 cases, and I am convinced of their value, although where there is haemoptysis or even blood-stained sputum I omit the sanguinarin. In very acute cases I administer these drugs alone for a week or two before beginning the antimony injections, lest the antimony, by increasing congestion in the lungs, might do fatal injury. If the sanguinarin irritates the stomach it should be discontinued. For pleuritic pain in phthisis I use a simple liniment of turpentine, alcohol, and compound camphor liniment in olive oil, to be rubbed on the chest and back daily.