the paretic or paralysed limbs splinted in the position of selection. We have found that this can be commenced in approximately four to six weeks from the onset, and the frame finally discarded in a further two to three months.

Notes of six early cases treated on these lines are appended.

Case 1.—A boy, aged 3½ years, was first seen three days after onset. He had complete paralysis of both legs, but was able to contract the flexors of the hips slightly with assistance. Back weak. High temperature. Cerebral cry. He was immobilized on a frame, with club-foot shoes holding the feet. Two days later removal of the frame was suggested as the child was thought to be dying, but it was retained, and the acute stage subsided two days later. The frame was discarded in two months. Three months later the boy began to walk, and two years later was leading a normal life.

Case 2.—A boy, aged 3 years, was first seen five days after onset. There was complete paralysis of both legs, thighs, and glutei, and very marked weakness of the back and the abdominal muscles. He was immobilized on a frame with two light plaster cases holding the legs in the desired position. He began to discard splints eight weeks later, and four months after the onset all muscles were acting voluntarily. Back and abdomen good.

Case 3.—A boy, aged 5½ years, was first seen five days after onset with the left leg and foot flail-like. The right leg quadriceps and glutei were normal, but the back and abdomen were weak. He was immobilized on a frame, with a plaster case for the left leg. He first began to discard splints five weeks later, and eighteen months from onset had regained full control of the limbs and was leading a normal life.

Case 4.—A boy, aged 5 years, was first seen four days after onset; the left leg was flail-like, the right quadriceps weak, the back and abdomen very weak, and the left arm weak. He had a high temperature, with periods of unconsciousness. He was immobilized on a frame, with plaster casing of the left leg and arm. He first began to discard splints three months later, and two years afterwards was walking with calliper splint, his arm being apparently normal.

Case 5.—A girl, aged 8 years, was first seen seven days after onset with complete paralysis of the right leg and shoulder. She began to discard splints six weeks later, and two and a half years afterwards had full control of the leg and shoulder, but the grip of the right hand was slightly weaker than that of the left.

Case 6.—A girl, aged 6½ years, was first seen eight days after onset. The left leg and right arm were flail-like, and the right leg weak. The temperature was 100° to 101°, and she was very irritable. She was immobilized on a frame with plaster cases on both legs, and her arm was tied to the top of the splint. One week after fixation the temperature and irritability were much diminished. She first discarded splints six weeks later, and a year afterwards was an apparently normal child.

Of the fourteen cases since July, 1926, in nine both lower limbs, the back, and abdominal muscles were affected. Of these, six are now gradually discarding their splints. In three cases one leg was affected; of these, two are now walking in callipers, and one is free of all splints. In the two remaining cases both upper arms and shoulder girdles were affected. These two children are wearing plaster cases, which are being discarded while in the prone position.

I am indebted to Dr. Mitchell Smith, under whose care these patients were, for permission to publish these notes.

THE ELECTRICAL TREATMENT OF THE PARALYSIS OF POLIOMYELITIS.

BY

R. J. MORRIS, C.B.E., M.D.DURH., M.R.C.P.LOND.,
HARROGATE.

In a paper published in the *Archives d'Electricité Médicale* in February, 1912, Professor Bergonié appealed against the discouraging prognosis given of infantile paralysis in spite of the efforts made by medical electricians to show that this need not be so, and that it is only so when, through lack of courage or through scepticism, treatment is omitted or given as a matter of form, or is badly applied by incompetent persons, or is insufficient as regards intensity, duration, or instrumentation. Electrotherapeutic treatment, more than other methods, requires clinical experience and practical knowledge which it is impossible to acquire through books; and this applies very particularly in cases of infantile paralysis. The part of the medical electrician will begin at the end of the acute stage by making a

precise and complete electrodiagnosis as a basis for everything which shall be done afterwards. To examine the leg, for example, a compress soaked in warm weak saline solution is wrapped around the foot and ankle, and a flexible metallic electrode bandaged to it; the patient may be seated or recline on a similar electrode of large size placed about the buttocks. These electrodes are then connected to the poles of a source of faradic current of low tension (coil with coarse winding); this current is passed through a metronome which rhythmically interrupts and reverses it about forty times a second. The intensity of the current is regulated by a progressive rheostat; if the intensity is very gradually increased, the coil being of the proper coarse winding, and the electrodes large, the process will be painless. It is possible to see and feel the muscles which respond, and those which have lost their faradic excitability. To ascertain which muscles show signs of degeneration it is only necessary to move the commutator of the apparatus and send the galvanic current by the same electrodes to be rhythmically interrupted and reversed by the same metronome at the rate of twice a second. By this means the muscles can be divided into two classes: (1) those which have preserved their faradic excitability; (2) those which have lost it and show the slow contraction of degeneration. Having completed the electrodiagnosis the treatment can be continued with the same apparatus.

It will be necessary to commence with the galvanic current rhythmically interrupted and reversed and to follow with the faradic current; the instrumentation is simple, and consists of a source of supply of galvanic current (24 to 36 volts), a metronome to interrupt and reverse the current rhythmically, and a finely graduated rheostat; the milli-ampere-meter may be dispensed with. This electrization of all the muscles of the limb has the advantage that by sight and touch we can be certain that the affected muscles are well acted on, and we can see the progress and modify the technique. There is no objection to this electrization *en masse* on the theory that the healthy antagonists should not be stimulated at the same time as the partially paralysed muscles. The results are more favourable than in cases subjected to local electrization. The duration and intensity of the electrization are very important. It has been proved by experience and numerous observations carried on over a long period that the condition of the patient improved more and more as the treatment was increased in duration and intensity; it can therefore be laid down as a principle that the intensive treatment of the paralysis of infantile paralysis can be repeated with a duration and intensity not formerly used. Bergonié states that he was driven to this conclusion little by little by ever better results. He quotes, from a large number of cases similarly treated, one of infantile paralysis of the hemiplegic kind. After two months the muscles of the leg responded to faradic stimulation, while in the arms, on the contrary, the biceps, triceps, and deltoid muscles did not respond. Treatment was commenced with an electrode of 300 sq. cm. at the level of the shoulder, and the hand was enveloped in the other electrode. Electrization commenced with half an hour a day at one sitting, then three-quarters of an hour, then one and a quarter hours, always at a sitting, finally one and a half hours at two sittings, then two hours—one hour in the morning and one in the evening; this was continued for two months. The child played and slept during the electrizations; his arm, roused by extensive rhythmic movements, did not make him suffer at all; he did not even wake when the current was intensified. The muscles made such progress that the biceps alone was a little backward, but it contracted at will.

The practice of long electrizations *en masse* by galvanic currents rhythmically interrupted and reversed automatically is the best treatment in these cases. Bergonié frequently stated that, from his experience of over twenty-five years, even in the most serious cases of infantile paralysis a good prognosis can be given if the electrical treatment is intensive, well regulated, and assisted by the adjuvant treatments—muscular relaxation, massage, passive and resistance movements, exercises, and, in particular, warmth to the limbs, as discussed by him at the Congress of Physiotherapeutics in 1910.