FIG. 1.—Showing two posterior grafts six months after implantation as thin osteoperiosteal shavings.

FIG. 2.—Showing consolidation of grafts one year after operation, and after using foot for six months. Note commencing equinus deformity due to faulty implantation of grafts and unequal growth at lower end of tibia.

FIG. 3.—Taken two months later, showing fracture of graft and further equinus deformity in tibia.

BERNARD HUDSON: A CASE OF ARTIFICIAL PNEUMOTHORAX.

FIG. 1.

FIG. 2.

FIG. 3.

FIG. 4.

epiphyseal line is seen to be of primary importance. They are to be used as osseous ligaments, and should retain sites of implantation corresponding with ligamentous attachments. In the neighbourhood of growing epiphyses due regard must be paid to the laws of growth. This case may therefore be looked upon as an instance of the extraneous fixation of the epiphysis to the diaphysis (by a graft), which the late Sir William Macewen showed would interfere with or prevent normal linear growth.

A CASE OF ARTIFICIAL PNEUMOTHORAX.

(With Special Plate.)

BY

BERNARD HUDSON, M.D.CANTAB., M.R.C.P.,
SWISS FEDERAL DIPLOMA; MEDICAL SUPERINTENDENT, VICTORIA
SANATORIUM, DAVOS-PLATZ, SWITZERLAND.

A MAN, aged 26, was under treatment for eighteen months (1922-23) for pulmonary tuberculosis affecting the upper lobe of the right lung. Recovery was excellent, and, by the sanction and advice of his doctors, he went to Kenya Colony in September, 1924, and invested in a farm there. The place suited him, and all went well until some months ago, when he became very ill, apparently (as far as can be made out from the history) with an acute exacerbation, which spread rapidly through the right lung, but chiefly affected the base. He saw a doctor, who advised him to get away as quickly as possible, and go straight to Davos-Platz. He arrived here under my care on February 22nd, 1926. He was exceedingly ill on arrival, the daily temperature varying from about 99° in the morning to 102° and over at 7 p.m. (taken by the mouth). He had a racking cough, and was bringing up a large quantity of sputum loaded with tubercle bacilli; he had lost about 2 st. in weight, and was looking a typical picture of what is known as "galloping consumption."

On examination the whole of the right lung was dull, and the lower half absolutely so; the whole lung was full of moist râles, and at the base there was loud bronchial breathing, increased vocal resonance, and increased vocal vibrations. Some moist crackling râles were heard in the upper lobe of the left lung, and the disease apparently had well started in this lung.

The radiographic appearance (Fig. 1) showed extensive trouble in the right lung, cavitation at the apex, and a dense black shadow in the lower part. It will be seen that the heart is drawn well over to the right side; the left lung shows commencing tubercle in the upper lobe. The appearance of this radiograph made me consider the possibility of an effusion at the right base, and I accordingly explored with a large-bore needle, but got nothing out. As the patient was obviously going downhill, and showed no signs of any improvement, I thought it worth while to see if it would be possible to establish an artificial pneumothorax on the right side, although there seemed very little hope of being able to do so, because of the probable presence of strong adhesions. However, on March 13th an attempt was made, and, to my great astonishment, a free space was found at once, the manometer swinging well, showing a strong negative pressure; 150 c.cm. of air only was introduced and a small initial pocket made. This amount of air sufficed to raise the pressure from a negative one to a strongly positive one, the pressures being as follows:

Induction	...	...	-2 -6	...	150 c.cm.	...	+2 -4
March 16th	...	...	-2 -6	...	200 c.cm.	...	0 +3
March 21st	...	...	-2 -7	...	300 c.cm.	...	+1 -4

A radiograph taken shortly afterwards showed the production of a small pocket (Fig. 2). Two more insufflations were made, the results being as follows:

March 31st	...	...	-2 -7	...	300 c.cm.	...	0 -3
April 10th	...	...	-3 -8	...	500 c.cm.	...	0 -1.5

The fillings were continued at intervals of a fortnight, about 600 c.cm. being introduced each time, and the radiograms (Figs. 3 and 4) show the development of a fair-

sized artificial pneumothorax in the lower part of the right lung, many adhesions having given way, but leaving two or three powerful bands, which are very plainly seen in the illustrations.

An interesting point is that after the third insufflation the patient remarked that he felt his heart once more beating on the left side instead of on the right. On inspection of Figs. 3 and 4 it will be seen that the heart, which was formerly drawn over to the right side by adhesions, has returned to its proper position on the left.

July, 1926.—The pneumothorax, although only a partial one, caused an undoubted and pronounced improvement in the patient's condition. The cough and expectoration have very much diminished, the pulse rate, instead of being 120 to 130, is now running at about 80, the breathing is easier and quieter, and the temperature has come down. The patient is now getting up for a short period each day without any ill effects. The signs in the left lung are diminishing in extent and intensity.

February 1st, 1927.—The further progress of this case is as follows: The patient kept very well all through the summer, and was able to take moderate exercise, the temperature being always quite normal, and the cough and the sputum diminished. The general condition improved considerably, and he gained weight. Fillings were continued every three weeks, the pneumothorax cavity gradually increasing in size, and the adhesions thinning and stretching. On January 17th the patient's temperature began to rise again, and he developed pain on the side of the pneumothorax; this was shortly followed by the appearance of a small serous effusion, which is following the ordinary course of such effusions; and the patient is already improving daily. Incidentally it is of interest to note that with the establishment of this effusion the adhesions have given way, the pneumothorax now being nearly "complete"; the other lung is holding up in a very satisfactory manner.

I am indebted to Mr. P. G. Sutton, M.S.R., of the X-Ray Institute, Davos-Platz, for the radiographs.

PARAPLEGIA (MEGALOCYTIC) CURED BY HYDROCHLORIC ACID.

BY

FRANK C. EVE, M.D.CANTAB., F.R.C.P.LOND.,
SENIOR PHYSICIAN, ROYAL INFIRMARY, HULL.

DIAGNOSIS in the following case was possible by measuring the average diameter of the red cells, and by that alone. Haematology, and all its complexities, gave no light. How forcibly this illustrates my favourite doctrine that simple recorded measurements in clinical medicine are richly rewarded, yet greatly neglected.

A florid, healthy, active young man of 37, a cargo superintendent, decent and clean living—noticed tingling in both legs up to the groins in January, 1925; later his hands tingled, he was drowsy, and had sacral pains, worse on walking. Next his ankles gave way, micturition became difficult, and the pavement felt like cotton-wool.

He was much worried sexually at this time by the exactions of his second wife, to whom he had been married a few months. In July he could no longer walk, and in August I saw him in consultation with Drs. Eddie and Robertson. I diagnosed posterior combined sclerosis, and he was admitted into the Royal Infirmary under my care. There he lay on the balcony for five months, and then went to the seaside under Dr. Sproule. The plantar reflexes were extensor, knee-jerks absent, and he could not move his feet. Hydrochloric acid was not present in the test-meal secretions, and x rays were suggestive at first of pyloric new growth, but became normal after belladonna. Haemoglobin was 75 per cent., and the pathologist reported more than once that there was no evidence of pernicious anaemia. As he had had sore tongue before admission I still felt incredulous.

In view of the recent illuminating work of Dr. A. F. Hurst on this subject I ventured to send him a blood film and an outline of the case. At his instance, Dr. Hampson kindly carried out the measurements of