She still complains of a feeling of breathlessness, but this was a feature of the "nervous breakdown" two years ago. Anaesthesia to pinprick is still present, but is patchy in distribution. Epicritic sensibility appears to show definite improvement. The feet are bluish in colour and the skin is dry and atrophic; numbness, tingling, and coldness of the feet are still complained of, though less so. Knee-jerks and ankle-jerks are present and equal; the plantar reflex is flexor. No retinal haemorrhages can be found.

She is now permitted to replace one portion of the liver with underdone steak or kidneys on three days a week, and, oddly enough, prefers to replace the cooked portion rather than the raw. The raw liver can also be conveniently given in sandwiches or in a jelly, which can be flavoured to taste with bovril, lemon, etc.

It is, of course, too early to make any claim to permanent effect, but the rapidity of improvement appears to be remarkable in a patient of this age.

My thanks are due to Dr. H. M. McCrea for his kindness in giving me full particulars of the patient's past history and treatment, and to Dr. J. Bamforth for the blood counts.

THE THYROID AND MANGANESE TREATMENT.

A SUGGESTION WITH REGARD TO ITS POSSIBLE MODE OF ACTION.

BY

HERBERT W. NOTT, M.R.C.S., L.R.C.P.,
GUILDFORD.

NEARLY three years have elapsed since this treatment was promulgated; it has been widely and successfully practised, and the time seems to have arrived for a tentative investigation as to its possible mode of action. Some good will have been gained if such an inquiry helps to dispel certain views which are quite misleading. For example, the reports of acute pneumonia being rapidly cured by rectal injections of a solution of potassium permanganate have led some to disagree with the title of the treatment and suggest that the thyroid substance may be unnecessary, the possibility being overlooked that, by using potassium permanganate in the way we do, the thyroid gland may be relieved of its load and be allowed to function normally. Yet this was clearly foreshadowed in a paper in this JOURNAL in 1925 (vol. i, p. 443), where, under the heading of "goitre," the following results of treatment were given.

Parenchymatous Goitre.—Four cases are reported to have quickly responded to the treatment, two of them without the use of thyroid extract. One is inclined to relapse, unless she keeps up the treatment occasionally.

Exophthalmic Goitre.—Three mild cases find themselves very much better, with increased weight, and diminution of symptoms, after a few weeks' to a few months' treatment. Two advanced cases describe themselves as "less nervy," and the swelling of the neck has diminished in size. One case of enlarged thyroid gland is reported as having been on the point of having the operation of thyroidectomy, but shortly after the treatment was used operation became unnecessary. In another very severe case no relief at all was obtained after five weeks' continuous treatment, with injections.

Since then nothing has been more obvious to numerous observers than the satisfactory reduction in the size of goitres, so that, whether the thyroid substance is produced by the individual directly, or whether it is necessary to administer it where greater inefficiency exists, the treatment need not be looked upon as incorrectly named.

Some have loosely ascribed its action to the liberated oxygen, without explaining how this occurs. That it cannot be due entirely to oxygen seems quite certain, or why should we not get similar good results from the use of sodium permanganate?

Basic Principles Underlying Treatment.

In this paper it is suggested that an increase of oxidative processes in the tissues and fluids of the body is produced by the interaction of manganese, potassium, thyroid gland substance, and the liberated oxygen derived from the permanganate on reduction. In support of this hypothesis, the more recently acquired knowledge of the elements comprising the combination will be considered in conjunction with the accepted laws of oxidation. An argu-

ment will be adduced tending to show that a reinforced catalytic action must be set up in the blood and cells of the organism by which firmly established metabolites are expelled, to the relief of the autonomic nervous system and the internal secreting glands.

As biology teaches that life itself depends upon oxidation, so also biochemistry has taught that, when oxidation is incomplete or hampered, disordered function results; and as this is the very core of our subject, oxidation will be considered first.

Oxidation.

Bayliss summarizes oxidation as follows:

"Oxidation in the tissues is due to the presence therein in the first place of an organic substance which is readily oxidizable by molecular oxygen. A part of the energy set free in this process is made use of to form from other molecules of the substance an unstable peroxide. Benzaldehyde undergoing oxidation in air is an instance of such reaction. The analogous substance present in cells corresponds to what was originally called by Bach an 'oxygenase,' because it is by its means that molecular oxygen is activated. This activation is produced by the agency of an enzyme, peroxidase, which is a colloidal hydroxide of iron or manganese, kept active, or protected from precipitation or aggregation, by a stable or hydrophile colloid. This enzyme decomposes the peroxide with the liberation of 'active' oxygen, with its powerful oxidizing properties. We are still in ignorance as to what this form of oxygen is, but there seems to be much evidence to show that electrical forces play an important part."—W. M. Bayliss: *The Nature of Enzyme Action*. Fourth edition, 1919, p. 158.

This quotation reminds us that both manganese and iron must be present in the tissues before complete oxidation processes can take place. Now, if some people are not obtaining a sufficiency of either of these two essentials to oxidation, the fact that the consumption of meat has increased so much among Western peoples in the last few years would suggest that it is not the iron content of the food which is at fault.

Potassium.

Mathews of Chicago, who established the prime importance of potassium in the chemistry of the cell, writes:

"Another substance which is in some way concerned in the respiration of the cell is the element potassium. It has been found that in the presence of potassium hydrate, phloroglucin and similar substances undergo oxidation better than with the equivalent amount of sodium hydrate. This indicates that the potassium salt is more easily oxidized than the sodium. There must be some reason for the preference cells have for potassium over sodium. A general richness of potassium in cells of widely different character indicates that this element must be concerned with some fundamental process or condition in the cell, and it is possible that that process is respiration. But just why it is favourable, or what its real function is, it is impossible to state."—A. P. Mathews: *General Cytology*, Chicago, 1924, p. 55.

This, then, may account for the moderate success with the sodium salt.

Manganese.

Manganese has been shown by Bayliss to be essential to oxidative processes; it is found in the ash of all enzymes, and is a most potent catalytic agent. It increases antibody formation.¹ It is essential to growth, not only to the body generally, but to the thyroid gland.² Manganese exists in deficient quantity in white flour; wholemeal flour is rich in it.³ Artificial enzymes have been produced with alkaline solutions of manganese in gum arabic (Bertrand's "laccase").⁴

For some time it was hard to conceive how manganese entered the blood stream, seeing that it exists after reduction as the insoluble salt Mn_2O_3 . But it has been suggested that it probably forms a laccase with the mucus of the bowel or stomach, taking the place of Bertrand's gum arabic, and enters the blood stream as colloids do.

The Thyroid Gland.

The following appear to be the chief points in the later knowledge of this gland which are applicable to this subject.

1. The thyroid gland is generally considered to be the chief activator of oxidative processes, and is now given pride of place in the control of the endocrine system.

2. It is intimately concerned in regulating growth, and is profoundly affected by insufficiency in the food of growth-promoting factors of all kinds.

3. It is susceptible to toxic action even when the diet is not at fault, but more so when the diet is wrongly balanced.

4. When affected by toxins or ill-balanced diets it is so handicapped in its work as to be rendered physiologically subnormal and still more susceptible to toxic action.

5. It accordingly attempts a compensatory hypertrophy, sometimes succeeding, more often failing, and suffers from the usual effects of overstrain (that is, diffuse fibrosis), which induces a state of hypothyroidism (probably unregistrable by basal metabolic rate methods), of which the clinical signs may be so slight that they are only rendered apparent by the effects of thyroid therapy.

The last four paragraphs contain the gist of McCarrison's important paper (*Lancet*, April 30th, 1927, p. 916), in which he announced the origination of a new type of goitre in rats; by giving them diets resembling the average diet of Western nations he has caused serious changes to occur in their thyroid glands. Such an achievement may have widespread consequences, for human beings may be similarly affected. Need we wonder at so many women having unsymmetrical necks, and need we be surprised if thyroid therapy is found useful to a far-reaching degree?

This completes a short setting out of the basic principles which seem to underly this treatment. We have oxygen present in a form which may have great potentialities. Potassium is there to help the cellular interchange with a quickening of oxidative processes. Manganese, which is deficient in the modern human body, and which is indispensable for oxidation, is supplied in quantity and carries with it other necessary properties. Lastly, we have thyroid substance with all that it stands for to a generation which apparently is suffering from incipient hypothyroidism.

Here, then, is a formidable array of oxidizing agents, the combined action of which is likely to result in reinforced oxidative processes. This action might be compared to the fire on the domestic hearth, by substituting potassium for the paper, manganese for the wood, metabolites for the coal, and thyroid substance for the bellows. The potassium, or paper, sets up combustion; the manganese, or wood,

conveys the heat to the metabolites, or coal; if insufficient oxygen is present, or if the coal is of the foreign or "out-crop" quality familiar to all of us during the coal strike, the bellows, or thyroid substance, may be required before the whole of the material is consumed.

CONCLUSION.

It is the opinion of many who have used this treatment extensively—heightedened in the case of those who know what has been effected during the last two years in certain farms where domestic animals suffering from sepsis have been treated on like principles—that when these drugs are combined in their action oxidative processes are greatly increased, with the following effects, though not necessarily always in this sequence.

1. Rapid reduction of infective agents and of toxic products in the blood and tissues.

2. This is followed by a lessening of the load on the thyroid gland, which quickly, or slowly, returns to its normal action according to the degree of injury it may have received.

3. The continued administration of thyroid substance provides for any deficiency of supply of thyroxine when the gland is seriously injured.

4. A purer blood supply also eases the strain on the other internal secreting organs, and more normal secretions are prepared. Where hormones of good quality have been secreted, or where vitamins in good quantity have been ingested, only to be neutralized by toxic products in the blood, this waste is quickly lessened or ceases altogether.

5. By the removal of the condition of nerve-block, conceivably brought about by organic compounds just as selective in their action, perhaps, as are the drugs atropine, digitalis, and aconite, the delicate balance of the autonomic nervous system is restored in so far as degenerative changes will allow.

REFERENCES.

¹ Walbum and Morsh: *Ann. de l'Inst. Pasteur*, 37, 1923, 396. ² R. McCarrison: *Indian Med. Research Journ.*, January, 1927. ³ *Idem*: *Lancet*, April 30th, 1927, p. 916. ⁴ Bayliss: *Principles of General Physiology*, 1924, p. 585.

Memoranda :

MEDICAL, SURGICAL, OBSTETRICAL.

PLACENTA PRAEVIA IN FOUR SUCCESSIVE PREGNANCIES.

THE following notes seem to be of sufficient interest to warrant publication.

In March, 1920, I was urgently summoned to an agricultural labourer's wife. I found that she was six months pregnant and had lost a large quantity of blood. She was very blanched, with a small, rapid pulse, and was very faint, but quite conscious. Examination showed a central placenta praevia with a lax os the size of a five-shilling piece. A little chloroform was at once administered, the hand put into the vagina, and in a couple of minutes the os was digitally expanded sufficiently to allow the edge of the placenta to be reached. The placenta was pushed aside, forceps put on, and the child delivered.

The next time I saw this woman was in response to an urgent summons in May, 1922. I found exactly the same state of affairs save that she was now seven months pregnant. I acted exactly as before, with the same result.

In April, 1925, there came another urgent summons; everything was as on the previous occasions save that she was eight months pregnant. I acted as before, and with the same result.

I next heard from her about the middle of November, 1927. She told me that she was expecting to be confined early in December, and asked if I would attend her; she had never troubled to engage me on the other occasions. I wrote saying that I would much prefer not to, and heard no more. On December 9th, however, I got an urgent message at 8 a.m. that she had lost a lot of blood, more than ever before. I went at once, and found her blanched and fainting; the os was about the size of half a crown. I plugged the vagina very tightly, put on a tight binder, and took her in my car to the maternity ward of the local hospital. She appeared to lose little after this, and two hours after admission I was able to do as on the three previous occasions.

The woman is now 40 years of age; her first placenta praevia was her tenth pregnancy. The durations of the

pregnancies were six, seven, eight, and nine months respectively; all the infants were dead. The question of performing a Caesarean section never really arose save perhaps in the last pregnancy, but if she again becomes pregnant, and information is given in time, section with removal of the uterus would probably be the best plan. I am aware that the classical method is to turn and bring down a leg, but in this woman's case, and in that of two other multiparae with central placenta praevia who were unconscious when first seen, and in whom the placenta was torn through centrally and the forceps applied, it appeared to me that less blood would be lost by the method used than if turning were performed.

I realize that I have been very fortunate in not having to deal with severe haemorrhage with an undilated os.

T. MACCARTHY, M.R.C.S., L.R.C.P.

Sherborne, Dorset.

THE FORGOTTEN SWAB.

THE subject of this note is the first surgeon's swab as the second surgeon sees it. Case I gives an excellent example of the powers of the body in walling off a foreign body in the abdomen. Case II is perhaps unique, and I should be interested to hear if a similar case has been reported.

CASE I.

A married woman, aged 34, was admitted to hospital in May, 1926. Three days before admission she was seized with sudden acute pain in the lower part of the back, and a diagnosis of acute lumbago was made. She had been operated on in Calcutta a year previously for a ruptured ectopic gestation. There was no other history of note. On examination there was found a fixed, fluctuant, tender swelling arising out of the pelvis; per vaginam the same swelling could be felt and its fixity and fluid nature could more easily be demonstrated.

Operation.—The abdomen was opened by a right paramedian incision below the umbilicus, and a cyst the size of a grape fruit was found fixed in the pelvis; it extended upwards and was