who still have symptoms have been operated upon at a relatively short period, and will almost certainly continue to improve until they are completely cured. Three of the cases have developed a considerable hypertrophy of the remaining portion of the thyroid, and have had little or no improvement of their symptoms. It is probable that they may yet be improved by further operation, which is now contemplated. In addition to these, 3 others, who had apparently passed to a state of complete cure, relapsed owing to some severe mental stress, 2 of them as a result of having been exposed to air raids. These have since improved so much under medical treatment that no operation has been contemplated, although they still have some evidence of the disease; 8 have died since operation—one from pneumonia; one during a confinement; 2 six and four years respectively after the operation, apparently from conditions not connected with the thyroid; one developed carcinoma of the skin owing to an x-ray burn and has since died from this; 2 were said to have died from heart disease, which must be regarded as being dependent upon their Graves's disease; one developed diabetes and died, and although all symptoms of Graves's had disappeared there is so close a relationship between these two conditions that this fatality must be included as a post-operative possibility. In addition to the patient who died from diabetes, one other has diabetes, which is controlled by medical treatment, and she has no symptoms of exophthalmic goitre. She has therefore been included among the group who are relieved. Nineteen have been lost sight of.

These results may be tabulated thus:

Quite cured	...	...	...	...	...	94
Greatly relieved	...	...	...	...	...	46
No better	...	...	...	...	...	3
Died after operation	...	...	...	...	...	9
Late deaths	...	...	...	...	...	8
Lost	...	...	...	...	...	19
Relapsed	...	...	...	...	...	3

182

It is justifiable to claim, therefore, that as a result of operative treatment there will be an immediate mortality of about 5 per cent., a late mortality of under 2 per cent., a certainty of complete cure in about 55 per cent., and of sufficient relief for the patients to earn their own living in comfort in 81 per cent.—results which are unlikely to be obtained by medical treatment alone.

It is important, however, to realize that there is not an immediate cure as a result of operation. I have on several occasions pointed out that the post-operative progress passes through several stages, which may be classified as follows:

1. *The Stage of Reaction.*—This is the dangerous post-operative period, and it lasts from three to four days after operation, the symptoms in the majority of cases being due to a definite post-operative hyperthyroidism. With modern methods of operation it is much less in evidence. If the symptoms become serious they can generally be controlled by the administration of Lugol's solution, of large quantities of fluid, and of morphine.

2. *The Stage of Primary Improvement.*—Within a fortnight from operation a great improvement is nearly always manifest. Many patients before leaving hospital have a normal pulse with no tremor; they are eating and sleeping well, the exophthalmos may be much reduced, and the improvement from their condition on admission is often startling.

3. *The Stage of Primary Relapse.*—Nearly always on returning to home conditions there is a disappointing return of the symptoms. The relapse is slight in degree, but it is because of its presence that a rather prolonged convalescence is required, and that the patients must be carefully watched and treated for a considerable period. As a general rule, it does not last longer than four to six weeks, but if the case has been so severe that it has only been possible to remove a relatively small portion of the thyroid, the patient may permanently remain in this stage, and although showing improvement will not pass to a stage of complete cure. This is not uncommonly a difficulty with such cases. It is evident that further operation is required to bring about a cure, but the patient may be so much relieved that she refuses to go through the discomfort of a second operation.

4. *The Stage of Apparent Cure.*—After a few months these patients lose all their symptoms. There is no enlargement of the thyroid, all nervousness disappears, the pulse is normal, and the exophthalmos slowly disappears. The patients appear to be absolutely cured, and if everything is satisfactory they will pass on to a state where they are absolutely cured. It is only by having watched many cases that one realizes that during this period any severe mental shock or upset may be followed by an enlargement of the remaining portion of the thyroid and by recurrence of the symptoms. It is very difficult to estimate how long this period lasts, but it would appear to persist for one or two years. At the end of this time the cure is apparently complete, for not only are the patients free from all symptoms, but they seem to be able to face all normal disturbances and upsets with no fear of a recurrence of the symptoms.

LITERATURE.

1. de Quervain, F.: *Goitre*, London, 1924.
2. Fraser, F. R.: Iodine in Exophthalmic Goitre, *BRITISH MEDICAL JOURNAL*, 1925, i, p. 1.
3. Jackson, A. S.: Iodine as an Aid to Surgery in the Treatment of Exophthalmic Goitre, *Lancet*, April 11th, 1925.
4. Murray, G.: Discussion on Exophthalmic Goitre, *BRITISH MEDICAL JOURNAL*, 1922, ii, p. 908.
5. Plummer, H. S., and Boothby, W. M.: *Journ. Iowa State Med. Soc.*, 1924, p. 1466.
6. Pemberton, J.: Mortality in the Surgery of Exophthalmic Goitre, *Surg. Gynecol. and Obstet.*, April, 1923, p. 458.
7. Walton, A. J.: The Effect of Heat upon Operations for Exophthalmic Goitre, *BRITISH MEDICAL JOURNAL*, 1923, i, p. 1045. A Consideration of Some Cases of Exophthalmic Goitre Treated by Operation, *Lancet*, 1914, i, p. 1387. The Operative Treatment of Exophthalmic Goitre, *Practitioner*, October, 1914. Fifty Cases of Exophthalmic Goitre Treated by Operation, *Practitioner*, October, 1917. Discussion on the Medical and Surgical Treatment of Graves's Disease, *Proc. Roy. Soc. Med.*, July 21st, 1921, p. 42. Surgery of the Thyroid Gland, *Lancet*, July 14th, 1923, p. 53, and August 11th, 1923, p. 267.

A Lecture

ON

THE USE OF HYPERTONIC SOLUTIONS IN THE TREATMENT OF INCREASED INTRACRANIAL PRESSURE.*

BY

W. RUSSELL BRAIN, M.D., M.R.C.P.,

ASSISTANT PHYSICIAN TO THE LONDON HOSPITAL AND TO THE HOSPITAL FOR EPILEPSY AND PARALYSIS, MAIDA VALE.

Two cardinal facts about the skull and its contents form the basis of a true understanding of the pathology of the intracranial pressure. The skull in the adult is a rigid case which is incapable of expansion, and the total volume of the intracranial contents is therefore constant. These contents may for practical purposes be regarded as three—the brain, the cerebro-spinal fluid, and the blood in the blood vessels. Since their total volume is constant, an increase in the volume of any one of them can only occur at the expense of the volumes of one or both of the others. For example, if the brain becomes enlarged by the growth of a tumour there must necessarily be a corresponding decrease in the volume of the blood or of the cerebro-spinal fluid, or of both. Such readjustments can only occur to a very limited extent without raising the intracranial pressure.

The most obvious way to reduce raised intracranial pressure is to abolish in part the rigidity of the skull—in other words, to perform a decompressive operation. This was till recently the only method available, and it is still the method of election in certain circumstances. There are, however, a number of conditions associated with increased intracranial pressure in the treatment of which such a serious operation is unjustifiable, and others again in which it is to be avoided if possible. My present object is to describe another method of lowering the intracranial pressure which has many therapeutic applications, and which can be used sometimes as a substitute for, and sometimes as an adjuvant to, craniectomy.

The Physiological Action of Hypertonic Solutions.

We owe to Weed the discovery that hypertonic solutions can be used to lower the intracranial pressure. Weed

* Delivered to the South Middlesex Division of the British Medical Association.

injected intravenously into animals a concentrated solution of sodium chloride, and showed that a pronounced fall in the pressure of the spinal fluid occurred, and that if part of the skull were first removed shrinkage of the brain could be observed to follow the injection. Further investigations demonstrated the means by which these results were brought about. The intravenous injection of hypertonic solutions raises the osmotic tension of the blood, and so leads to the passage of water from the brain into the blood stream. There is an actual resorption of cerebro-spinal fluid, which passes up the perivascular spaces into the cerebral capillaries, and, according to some authorities, also passes back into the blood through the choroid plexuses. The result is a marked fall in the intracranial pressure. It was not long before other workers demonstrated that the same results could be achieved less directly by administering hypertonic solutions by way of the alimentary canal. These raise the osmotic tension of the blood by withdrawing water into the intestine.

Modes of Administration.

1. *Intravenous Injection.*—Intravenous injection is used when it is desired to lower the intracranial pressure as rapidly as possible. The most convenient hypertonic solution for intravenous use is a sterile solution of sodium chloride in distilled water. The maximal dose is 100 c.cm. of a 30 per cent. solution, and I have given this amount in certain conditions without ill effects. For most purposes, however, this is more than is necessary, and it is sufficient to give 70 to 100 c.cm. of a 15 per cent. solution, or half these amounts of a solution double the strength. Some workers employ concentrated solutions of glucose, for which certain advantages are claimed. It is said that after the intravenous injection of glucose the fall of intracranial pressure is slower and more sustained, and is less likely to be followed by a reactionary rise than after the use of sodium chloride. Moreover, glucose possesses nutritive value, and is of help in combating shock and acidosis. The maximal dose is 100 c.cm. of a 50 per cent. solution in normal saline. Whether sodium chloride or glucose solution is employed, it should be administered very slowly at a rate not exceeding 3 c.cm. a minute.

2. *Administration by the Mouth.*—To obtain comparable results by oral administration it is necessary to give large amounts either of sodium chloride or of magnesium sulphate. The dose of the former is 16 grams, given in 2-gram capsules with 80 c.cm. of water, and of the latter 3 ounces of a 50 per cent. solution. Oral administration possesses certain disadvantages, and I have not employed it to obtain a rapid reduction of intracranial pressure. Repeated doses of a half to one drachm of magnesium sulphate, however, may be given by the mouth when a mild continuous action is desirable.

3. *Rectal Administration.*—This is the most generally useful way of giving hypertonic solutions. The dose is 6 ounces of a 50 per cent. solution of magnesium sulphate, which should be run slowly into the rectum at body temperature. Unless it is retained for half an hour it will not produce its full effect and should be repeated an hour later.

Some points of importance in the use of hypertonic solutions must be mentioned. Since the method operates by dehydration the patient's intake of water must be restricted if the full effect is to be obtained. Solutions of magnesium sulphate should on no account be given intravenously, since, when administered by this route, they produce general anaesthesia and respiratory paralysis. The rectal injections, if repeated too frequently, may produce irritation of the rectum.

Persistent Cerebral Contusion.

We now come to the consideration of conditions which are amenable to treatment by means of hypertonic solutions. We owe to Trotter the recognition as a clinical entity of a condition which is perhaps best described as "persistent cerebral contusion." This is the result of a minor head injury—either a fall on the head which at the time gives rise to simple concussion or a localized injury not associated with loss of consciousness and produced by a

fall or a blow. Following such an accident a patient may develop a train of symptoms which are only too familiar, and which, since they are often extremely disabling, are of considerable medico-legal importance. The principal symptom is headache, which may not appear until the patient gets up, if he has been confined to bed following his injury. The headache is paroxysmal and is sometimes extremely severe. It is especially liable to be brought on by mental excitement, muscular exertion, and stooping—activities which often lead also to giddiness. The patient is usually nervous and irritable, and may develop a typical anxiety neurosis. There are as a rule no physical signs of a gross injury to the brain or of raised intracranial pressure, but in a considerable number of cases bradycardia is present. It is unusual for radiograms of the skull to show any abnormality.

The persistence of the symptoms, which may endure for months or years after the injury, is explained by reference to the pathology of the condition. A blow which does not fracture the skull may, owing to the elasticity of the calvarium, nevertheless produce a localized contusion of the brain. Such a contusion is characterized by the extravasation of blood and plasma from damaged vessels. Owing to the rigidity of the skull, the brain, as we have seen, can only swell at the expense of the cerebro-spinal fluid and the circulating blood. The localized area of oedema due to the contusion thus becomes surrounded by an area of impaired circulation, which prevents the removal of the exudate, a vicious circle being established which it is the object of treatment to break.

The prevention of persistent cerebral contusion is to be achieved only by the adequate treatment of the case in the early stages; but the method of treatment is the same whether the case is seen early or late. Rest is the first essential, and the patient should be absolutely confined to bed and not allowed to get up for any purpose. Hypertonic solutions are employed to lower the intracranial pressure and assist in the absorption of the exudate. In mild cases it may be sufficient to give half to one drachm doses of magnesium sulphate thrice daily by the mouth, and this may be usefully combined with 10 grains of potassium bromide. Intravenous injections of hypertonic saline, however, are more effective and should, if possible, be used in all cases. The usual dose is 100 c.cm. of 15 per cent. sodium chloride in distilled water, and this may be repeated in four or five days if necessary. In some cases one such injection is sufficient to free the patient from headache, and, since the treatment is curative, he may then be allowed to get up. If medical measures fail, a subtemporal decompression can usually be relied upon to relieve the symptoms, but it is hardly ever likely to be necessary if hypertonic solutions are given a fair trial. Early treatment of cerebral contusion is of great importance, since if the patient is given time to develop, in addition, an anxiety neurosis, he presents a much more difficult problem.

Head Injuries in the Acute Stage.

Cases of head injury in the acute stage present many difficulties, both of diagnosis and treatment, which can only be successfully met by the close co-operation of surgeon and neurologist. Hypertonic solutions are of definite value in treatment, whether or not operative intervention is undertaken. They can be used as an adjuvant to decompression to aid in the reduction of cerebral oedema. They may tide the patient over the difficult period when operation hangs in the balance, and so render decompression unnecessary; and they undoubtedly promote recovery and diminish the risk of sequelae in the milder cases of contusion and concussion in which operation is not indicated.

In all cases of severe injury the intravenous mode of administration should be used, and the glucose solution would seem to possess the advantage of combating acidosis and shock. Hypertonic solutions, however, should not be employed in any form if shock is profound, as indicated by low temperature and blood pressure, and a rapid or rising pulse rate. The treatment of head injuries by hypertonic solutions is the rational use of methods which in the form of free purgation with calomel and salines have been used empirically for many years.

Intracranial Tumour.

To explain how relief can be obtained by means of hypertonic solutions in cases of intracranial tumour it is necessary to give a brief account of the way in which a rise of intracranial pressure is brought about in this condition. We may recognize three factors which contribute to this result—the direct pressure of the tumour, the effect of the tumour upon the intracranial circulation, and the combination of both of these factors to produce internal hydrocephalus. It is easy to understand that the growth of a tumour within the brain raises the intracranial pressure in its immediate neighbourhood. In addition, by pressure upon surrounding veins, it raises the venous pressure locally, and in many cases throughout the whole cranial cavity. The local venous congestion leads to oedema of areas of brain surrounding the tumour, and so intensifies the local rise of pressure. The main causes of internal hydrocephalus in cases of intracranial tumour are obstruction to the free passage of fluid from the ventricles by the tumour, and the general rise of intracranial venous pressure, which leads to both increased formation and impaired absorption of cerebro-spinal fluid. Thus an intracranial tumour sets up a series of vicious circles whereby the intracranial pressure, once raised, tends to rise more and more.

The value of hypertonic solutions in cases of cerebral tumour lies in the fact that they break these vicious circles and permit a readjustment of the volumes of the intracranial contents which may last for a considerable time. This they do mainly by reducing the formation and increasing the absorption of cerebro-spinal fluid, and so temporarily relieving the hydrocephalus. The following are the chief indications for employing them.

1. *As an Aid to Diagnosis.*—Not uncommonly a patient with an intracranial tumour is in a semi-comatose condition when first seen, and is quite unable to co-operate in the examination of sensibility or the visual fields. In such cases two or three rectal injections of magnesium sulphate solution may be sufficient to restore the patient to consciousness and render a full examination possible.

2. *In Emergencies.*—If a patient with an intracranial tumour suddenly becomes comatose, as may occasionally happen after ventriculography or in other circumstances, an intravenous injection of hypertonic saline may reduce the intracranial pressure sufficiently long to permit operation to be performed.

3. *As Palliative Treatment.*—In inoperable cases or while a patient is awaiting operation it is often possible to relieve headaches and vomiting by giving magnesium sulphate either by the mouth or by the rectum.

Cerebral Haemorrhage.

Both Leonard Hill and Cushing showed a good many years ago that the injection of fluids under pressure into the subarachnoid space led to a rise of blood pressure. This response of the vasomotor centre to a rapid rise of intracranial pressure is responsible for yet another vicious circle in intracranial pathology. The majority of cerebral haemorrhages occur in patients whose blood pressure is already high, but the haemorrhage, by raising the intracranial pressure, tends to cause the blood pressure to rise still further, and this in turn favours the continuance of the bleeding, or may even result in further haemorrhages elsewhere. Thus a rising blood pressure in a patient with a cerebral haemorrhage may be taken to indicate that the bleeding is still continuing. On the strength of these facts it has been urged that venesection is contraindicated in cerebral haemorrhage, because the medulla is already threatened with anaemia, and to withdraw blood from circulation is to add to its difficulties. Those who hold this view advocate lumbar puncture as the correct treatment, since by this means the intracranial pressure can be directly reduced without restricting the blood supply to the medulla. When the intracranial pressure is lowered, it is said, the blood pressure will fall, and the bleeding is then likely to stop.

These arguments against venesection and in favour of lumbar puncture do not take account of all the circumstances. I have never seen benefit result from lumbar

puncture in a patient suffering from cerebral haemorrhage, and I believe it to be definitely contraindicated in such cases. There are two serious risks attaching to it. The fall of pressure in the spinal theca may lead either to the rupture of the haemorrhage into one of the ventricles or to the downward displacement of the contents of the posterior fossa into the foramen magnum, in either case with rapidly fatal results. Venesection, on the other hand, could hardly have enjoyed so enduring a reputation in the past as a mode of treatment of cerebral haemorrhage if its results were uniformly harmful. If it merely reduced the blood pressure without at the same time lowering the intracranial pressure it would clearly be a dangerous procedure. Actually, however, since the intracranial pressure is ultimately dependent on the blood pressure, a fall in the blood pressure leads to a reduction of the intracranial pressure. Moreover, a diminution in the volume of the circulating blood also leads to a fall of intracranial pressure. Hence the theoretical risk of increasing the medullary anaemia is compensated by the relief afforded by a lowered intracranial pressure. Further, we must place to the credit side of the transaction the tendency of the lowering of the blood pressure to check the haemorrhage, its relief of the strain upon the heart, and its influence in increasing the coagulability of the blood. Venesection is therefore a rational form of treatment for cerebral haemorrhage.

Hypertonic solutions have only a limited value in this condition. Intravenous injections are contraindicated as tending to raise the blood pressure, and it is doubtful if the rectal mode of administration can influence a haemorrhage which is still in progress. The rectal injections, however, may be of value as a means of reducing the intracranial pressure in patients in whom there is reason to believe that the haemorrhage has stopped but who show no signs of recovering consciousness. Free purgation reduces the intracranial pressure in exactly the same way as hypertonic solutions administered by the alimentary canal, and possesses the same physiological justification.

Other Conditions.

There are other conditions associated with increased intracranial pressure in the treatment of which hypertonic solutions are useful. In the post-operative treatment of cerebral abscess they may be used to reduce oedema of the brain around the abscess cavity, and so diminish a tendency to herniation and promote drainage. They are also of value for the relief of headache in epidemic encephalitis and meningitis. Other applications will suggest themselves, but enough has been said to show that Weed's discovery has rendered available therapeutic methods of the greatest value in diseases of the nervous system.

EXPERIMENTS IN MALNUTRITION.*

BY

V. H. MOTTRAM, M.A.CAMB.,

PROFESSOR OF PHYSIOLOGY IN THE UNIVERSITY OF LONDON, AND HEAD
OF THE PHYSIOLOGY LABORATORY, KING'S COLLEGE FOR WOMEN
(HOUSEHOLD AND SOCIAL SCIENCE DEPARTMENT).

THE lines along which a diet may be inadequate so as to entail malnutrition are many. It may be insufficient in quantity or in quality. It may yield too little building material (protein mainly), too little fuel material (fats, carbohydrate, and surplus protein), or too little in the way of catalysts (mineral matter—for example, iron, iodine, calcium—and vitamins). The quality of the proteins may be poor—cereal proteins not having the biological value of meat or milk proteins. The diet may be inadequate because it has too little (or even perhaps too much) roughage, because the proportions between the different constituents are wrong, or because the way in which the food is presented is wrong.

It is clear from this that it is extremely difficult to work out experimentally the different possibilities of

* A lecture delivered to the St. Pancras Division of the British Medical Association, November, 1927.