power. This seems to me to be one of those cases of encephalitis so well described by Symons.

Furuncle is mentioned, not as being a complication of acute mastoid disease, but for the opportunity of making this assertion—that in all cases of long-standing otorrhoea in which a furuncle large enough to close or partially close the meatus occurs a radical mastoid operation should be done, especially if the furuncle be on the posterior wall of the meatus. In each of three recent cases I was fully justified in this line of action. In many of these cases there was a cholesteatoma, and this might mean erosion of bone; in one case the pus tracked backwards between the skull and the dura to make its exit by a hole in the skull in the sub-occipital region, burrowing its way beneath the pericranium into the soft tissues of the neck, where I ran it to ground, after freely removing the bone from the mastoid to the secondary hole in the skull. The dura was covered with granulations, as was the lateral sinus, and at the completion of the operation the lateral sinus burst. This man has so far made slow progress. His age is 47, and he has well marked emphysema of both lungs. The lateral sinus was never opened and the jugular never tied unless there were definite signs of phlebitis, when the jugular was tied high up in the neck and the lateral sinus exposed; if thrombosis was present it was dealt with in the usual way. If there had been signs of sinus infection, such as rigors without definite signs of phlebitis, the sinus was not opened, but if there were definite signs of phlebitis with or without thrombosis the sinus was opened, allowed to bleed, plugged, and saline given. In a boy with marked lateral sinus thrombosis, necessitating wide excision of bone for removal of a septic clot which extended very near to the torcular, a complete recovery resulted with good hearing power. In all the acute cases, primary or secondary uncomplicated, there were no deaths. Skin grafting was never done.

THE ETIOLOGY OF LEUKAEMIA.

BY

H. M. WOODMAN, M.B., B.Ch.

THE key to the etiology and nature of leukaemia is presumably to be found in the study of those factors which control the leucocyte content of the blood. While the line of investigation necessary is at present rather obscure, the constant recording of cases, especially those which do not strictly fall into any standard classification, may throw some light on the main causative factors. For this reason I have analysed the histories of twenty-five cases of leukaemia observed during the last six years.

The series consisted of 11 patients with myeloid leukaemia, of whom 7 were males and 4 females, and 8 cases of acute and chronic lymphatic leukaemia, 5 in males and 3 in females. The remaining 6 cases were atypical. The red blood count was below 1,000,000 in 37 per cent. of the series.

Myelogenous Series.

In the myelogenous series the total number of leucocytes ranged from 32,000 to 954,000 per c.mm. and the lymphocyte proportion from 0.8 to 40 per cent. In the case with the most marked lymphocytosis the polymorphonuclears and myelocytes were both 29.6 per cent. and the lymphocytes later fell to 1 per cent. The polymorphonuclear count was never lower than 29 per cent. and was usually above 40 per cent. This latter point may have something to do with the better prognosis of myelogenous leukaemia as compared with the non-granular form, where the polymorphonuclear count was never higher than 18 per cent. and ranged from 3.7 to 18 per cent.

Bad teeth or pyorrhoea occurred in 7 cases (64 per cent.), night sweats in 3 (27 per cent.), and a chronic infective septic focus of some kind was present in 82 per cent. of the cases.

There was nothing in any of the family histories to support the possibility of contagion from other patients. The duration of the disease varied from two to eight years.

The spleen was enlarged in all cases, and other lymphatic glands in 2 cases. The temperature usually ranged between the normal and 100°; in 3 cases there was no pyrexia and in one case the temperature reached 102°. Out of the 11 cases 7 were discharged as "better."

Non-Granular Series.

In the non-granular series, out of 9 cases 4 were chronic, 5 acute. The total leucocytes ranged from 4,800 to 928,000, but in one atypical case the figure was as low as 600. The lymphocytic percentage varied from 74 to 96. In all cases except one it was 80 per cent. or over.

Pyorrhoea or some chronic primary focus of infection was present in the histories of 6 cases (66 per cent.). Night sweats were recorded only in one case. The length of history was from one to seven months in the acute and five to eighteen months in the chronic forms. There was nothing in any family history to support contagion with any other case. The spleen and other lymphoid glands were enlarged in all cases except two; in one of these there was no splenic enlargement and in the other the spleen only was enlarged. There was usually no pyrexia in the chronic cases; the temperature rose as high as 105° in the acute. Two patients were discharged—these were of the chronic type; the remainder died.

Etiological Theories.

Of theories as to the causation of this disease only two need be considered in the light of present knowledge.

1. *The Tumour Theory.*

This theory is the most popular one at the present time. It is based mainly on certain changes which appear to be analogous to tumour formation, such as the excessive and apparently purposeless proliferation of leucocytes, and their tendency to metastasize. Several factors, however, indicate that if the nature of the agency instigating the condition is similar to that of tumour formation, it behaves quite differently from any other kind of malignant lesion. So far as is known, in a necropsy of a case of leukaemia, no site can be indicated as the possible primary focus.

The changes in the bone marrow and reticulo-endothelial system are more or less uniform, but certain areas may tend to be more dense than others. Patches of leucocytic deposit or gross infiltration will be relatively more marked in places, but these differ microscopically from centres of malignant proliferation. "The newly formed cells are not in possession of resolving power (of fermentative nature?) which proper tumour cells possess, and by which they entirely destroy the organs attached."¹ This is a very essential difference; it practically amounts to the difference between a benign leucocytic infiltration and an invasion by malignant cells.

Patients, moreover, have been known to survive for ten years or longer, and the condition may become entirely arrested for as long as seven years (Price). These are points hardly compatible with our knowledge of carcinomatosis and sarcomatosis.

2. *The Infective Theory.*

In the discussion on leukaemia and allied conditions at the Annual Meeting of the British Medical Association at Nottingham, Dr. R. Donaldson, after surveying some of the arguments against the sarcomatosis theory, concluded that there was more to be said for a possible virus or irritant, using the latter in its broad sense.² This certainly appears to be the most plausible etiological theory, and a survey of the present series of cases would seem to bring out points in its favour. Early investigators such as Virchow held this view and maintained that leukaemia was the result of specific infection. Others believe that it is non-specific in origin. In the histories of the above cases, 17 showed varying degrees of irregular temperature; in all the acute and atypical varieties the temperature was 103° or higher, but the chronic lymphatic patients had no rise at all.

There was a history of pyorrhoea, tonsillitis, sore throat, bronchitis, or some chronic infective focus in 82 per cent. of the classified cases and in all the atypical ones. The

occurrence of night sweats, to which Donaldson drew special attention as being a common feature, was recorded in 20 per cent. of cases. The length of history in chronic cases varied from two to eight years.

In rare instances cases have been recorded where contact with a leukaemic patient has apparently led to transmission of the condition, and Weiss and others have reported instances where several members of the same family were victims of the disease. Such incidents are, however, so rare that they can hardly be employed as evidence of the infective nature of leukaemia.

Animals grafted with leukaemic exudate from human beings have only in one questionable instance (that of Wiczowsky's experiment) developed leukaemia. Three patients dying from carcinoma were inoculated by Schuffer with leukaemic blood from another patient; none of them developed leukaemia.

It could be argued that if the disease is infective in nature it is hardly conceivable that the latter experiment could have failed. Nevertheless it may reasonably be supposed that a patient suffering from malignant disease is not to the same extent susceptible to the virus. In any case this experiment would have to be repeated by other observers to provide confirmatory evidence.

Ellerman has succeeded several times in inoculating healthy fowls with a filtrate from leukaemic fowls. The filtrate used consisted of an emulsion of organs passed through the Berkefeld filter and, therefore, cell free and microbe free for organisms up to the size of bacteria. The condition could thence be transferred from one bird to another indefinitely. These experiments go to show that at least, so far as the leukaemia of fowls is concerned, it is a definitely infective condition. Whether the disease in human beings is of the same nature is still open to question, but it would be difficult to imagine that leukaemia in human beings with its fundamental resemblances to the disease in fowls could be induced by an entirely different process. The following clinical details are of interest in this connexion.

A man, aged 66, had a two months' history of pain on the right side of the abdomen spreading up towards the thorax. The liver was large with a smooth, round, hard, and sharp edge, and there was a large palpable tumour on the left side with some of the characters of the spleen. This diminished under x-ray treatment. The red cells and colour index were normal. The leucocytes numbered 32,000 per c.mm., and the differential count was polymorphonuclears 64 per cent., lymphocytes 27 per cent., large mononuclears 4.4 per cent., mast cells 0.2 per cent., and myelocytes 4.4 per cent. Megaloblasts and normoblasts were present. There was very grave oral sepsis, indicating total extraction. The temperature ranged from 98° to 100°, but after extraction of the teeth it was above 100° for a few days. The total leucocytes varied from 8,000 to 32,000 during the month he was in hospital. No further symptoms developed, and he was discharged feeling well.

This case was diagnosed as chronic lymphatic leukaemia, though it was recognized as not being a typical case.

Many chronic infections—tuberculosis, for example—show a relative lymphocytosis, though admittedly the total leucocyte count would not be expected to reach 32,000. The lymphocytes varied from 24 to 38 per cent. There was certainly as much a parallel between the course of this case and that of a chronic infection (there was a very definite chronic septic focus), as between the course of this case and that of a case of "useless white cell proliferation," which leukaemia is supposed to be.

We might conclude, then, from the above analysis that leukaemia is a condition which runs a course in many striking respects parallel with that of an acute or chronic infective process, and that while there are clearly some unexplained resemblances to tumour formation (particularly if myelomas and chloromas are included under the heading of leukaemia) we must await further light on the whole problem of the etiology of tumours; this in turn will reveal the relation between the two conditions.

It may yet be proved that Gye's recent dictum is true—that the "agent of malignant disease is a living filterable microbe."

REFERENCES.

¹ *Leukosis in Fowls*, Ellerman, p. 12. ² BRITISH MEDICAL JOURNAL, October 2nd, 1926, p. 595. ³ *Ibid.*, November 13th, 1926, p. 870.

Memoranda:

MEDICAL, SURGICAL, OBSTETRICAL.

FOREIGN BODY IN THE BLADDER.

DR. HUNTER WOODS reports an interesting case of foreign body in the bladder (BRITISH MEDICAL JOURNAL, December 17th, 1927, p. 1140). May I mention two cases which came under my care at Guy's Hospital during the summer of 1926. Both were young females aged about 20.

The first case, when examined, showed a metal bar lying transversely in the bladder. The gentle efforts which were made to dislodge it through the urethra failed to move it, so I at once, fearing damage to the large pelvic blood vessels, cut down through the suprapubic route, and opened the bladder. With some difficulty I removed the foreign body, which was embedded in the right wall of the bladder and touched the left wall. The foreign body was a metal thermometer case, minus the lid, about three inches long. The patient afterwards developed left iliac thrombosis, so my fears as regards damage to important blood vessels were justified.

The second patient had caused much trouble at a country hospital, and was considered to have produced an artificial haematuria. She was cystoscoped during my holiday, and it was thought that she had a growth on the left wall of the bladder. When I cystoscoped her I felt certain it was not a growth, as I saw a similar mass at the base. I removed, through the urethra, two fruit stones. I set them in earth, and one produced a seedling tree. Unfortunately it suffered from being covered up and became mouldy, but I think it was some form of plum, cultivated, I suppose, as the stones were about an inch long—that is, much longer than the stone of any sloe, bullace, or wild plum that I have seen.

I think that, in the second case, the stones were introduced deliberately. But in the first case the thermometer case may have been introduced into the urethra during the passage of urine in order to allay spasm and pain.

In this connexion I may mention three other cases.

The first, a young child, had a vesico-vaginal fistula and a phosphatic stone in the bladder, which was found to have formed round a bone collar-stud about seven-eighths of an inch long. The second was that of an old man who had been passing bougies on himself for many years. I removed, suprapubically, three stones, all surrounding broken fragments of stiff bougies, but they were not of metal. The stones were phosphatic.

The third case was that of an elderly man who had been operated upon by the late Mr. Dunn for strangulated hernia. He suffered from retention afterwards and a coude catheter was passed. When this was removed it occurred to the patient that the catheter did not present quite the same appearance that it showed before introduction. When I examined him I found a firm mixed phosphate and urate stone, but in view of the history decided to perform suprapubic lithotomy rather than lithotripsy, for which the conditions otherwise appeared favourable, and a stone 1½ inches long, oval in shape, was removed from the bladder. The stone was sawn in half, and it was found that the upturned end of the coude catheter formed the nucleus of the stone. In my opinion the mere suspicion of a foreign body in a stone precludes lithotripsy in such cases.

A. RALPH THOMPSON, Ch.M., F.R.C.S.,
Surgeon, Genito-Urinary Department, Guy's Hospital.

INTESTINAL OBSTRUCTION IN AN INFANT.

A FEMALE infant, aged 3 days, was admitted to Charing Cross Hospital on November 27th, 1926, on account of vomiting. The pregnancy and birth had been normal and the child appeared well made. She took the breast well when put to it for the first time, but vomited soon after. Vomiting continued at frequent intervals (on one occasion continuously for an hour), though the child did not again take the breast. Meconium unusually light in colour was passed.

On admission the child seemed well and cried lustily. Some distended veins were visible in the abdominal wall, otherwise nothing abnormal could be detected. Some bile, but no blood, was seen in the vomit; the stools were small, dry, and putty coloured. An occult blood test was positive. A diagnosis of duodenal atresia was made. Saline and glucose were given subcutaneously, but the child continued to vomit and rapidly went downhill.

Laparotomy was performed by Mr. N. C. Lake on November 28th. A mid-line incision was made. On opening the abdomen a blind-ended distended piece of small intestine presented, and proved to be the upper few inches of the jejunum. Near to this blind end was found the proximal end of the rest of the small intestine, also blind, and of very small foetal type. The colon was undeveloped, the calibre of a lead pencil, whereas the distended small intestine was the size of a normal adult jejunum. This discrepancy in size made it impossible to perform an anastomosis, even had the condition of the child been good enough to justify such a procedure.