solution and swallowed may be caught up in the same way (H. K. Wilson). Bone buttons can frequently be detected after the swallowing of a bismuth solution.

Whether the foreign body be located in the larynx, the trachea, the bronchi, or the oesophagus, an antero-posterior and a lateral or lateral oblique x-ray examination should always be made. A comparison of the results thus obtained may be of the utmost value to the endoscopist.

Radiography is indeed the handmaiden of endoscopy, and her aid has often been and will continue to be of indispensable assistance in this branch of surgery.

In conclusion I make no apology for omitting any description or illustrations of the many types of endoscopes

and instruments used with them; they can be seen in the surgical instrument catalogues, their number is legion, and they increase more rapidly than the foreign bodies which they are devised to deal with. Still less would it be worth while to dwell on the details of manipulative technique, because these can only be acquired by patient and pains-taking practice.

My object will be attained if those who have glanced through the histories of the cases which I have reported will learn from them some of the lessons which it has seemed worth while to emphasize.

REFERENCE.

¹ BRITISH MEDICAL JOURNAL, October 17th, 1925, p. 686.

NORMAL CHOLECYSTOGRAPHY.*

BY

FRANCIS DAVIES, M.D.LOND.

(From the Anatomy Department, University College, London.)

(With Special Plate.)

ON examining a number of cholecystograms it at once becomes evident that the shadow of the gall bladder varies in different individuals, both in size, shape, position, and density. Dr. Evarts Graham,¹ the pioneer of cholecystography, predicted in an earlier paper that the shadows of gall bladders would probably be found to vary in position with the different types of bodily habitus. This preliminary report is concerned with the cholecystographic investigation of 100 medical students at University College, London, after administration by the mouth of sodium tetra-iodo-phenol-phthalein, in the endeavour to determine primarily whether the characters of the shadows of gall bladders in normal individuals vary with the different types of bodily habitus. Comment is also made on the oral administration of the dye, and on the details of the radiological technique employed.

1. Relation of Gall-Bladder Shadows to Bodily Habitus.

The students, whose ages varied from 18 to 30 years, comprised 80 males and 20 females, and they were classified after the manner of Mills into the following types of bodily habitus. Mills² gives the chief characteristics of each of these types as follows (Fig. 1):

1. *Hypersthenic*.—Massive physique, great body weight, heavy bony framework, short wide deep thorax, relatively long abdomen of greatest capacity in upper zones, subcostal angle very obtuse, lungs wide at bases and contract sharply to apices which project but little above clavicles, long axis of heart nearly transverse, alimentary tract high in position, gastric motility fastest and tonus most marked of any type (stomach emptying time about 4 hours 22 minutes), colon high and short, caecum well out of iliac basin (even standing), transverse colon truly transverse, marked colonic tonus, and motility rapid. This group comprises about 5 per cent. of individuals.

2. *Sthenic*.—Resemble hypersthenic but characters less marked; considerable body weight, rather heavy bony framework, thorax somewhat short and deep, lungs widest at bases, heart moderately transverse, subcostal angle about 90 degrees, digestive tract high with good tonus, and motility fairly rapid (stomach emptying time about 4 hours 46 minutes). This group comprises about 48 per cent. of individuals.

* This work was made possible by a grant from the Medical Research Council.

3. *Hyposthenic*.—Frailty of build approaches that of asthenic type, visceral characteristics approach those of asthenics, but lack the extremely low position and peculiar form of such; tonus rather poor and alimentary motility somewhat slow (stomach emptying time about 5 hours 26 minutes); lungs intermediate in proportions, heart pendent. This group comprises about 35 per cent. of individuals.

4. *Asthenic*.—Frail slender physique, light body weight, delicate bony structure, long narrow thorax, short abdomen of least capacity in upper zones, marked disproportion between great pelvic capacity and capacity of upper abdomen (pelvis may be as wide and capacious as that of a hypersthenic of twice the body weight); subcostal angle narrow, lungs widest in upper zones and apices extend well above clavicles, heart pendent ("drop-heart"); gastro-intestinal tract low, stomach often largely pelvic when standing, colon long, caecum capacious and low in pelvis, poor tonus and motility slowest (stomach emptying time about 5 hours 43 minutes, and often not empty after 6 hours). This group comprises about 12 per cent. of individuals.

In the investigation here described radiograms of the gall bladder were taken, after administration by the mouth of sodium tetra-iodo-phenol-phthalein, with the radiological technique described below. The data obtained from the cholecystograms are given in the table. From these data the following deductions may be made.

1. The length of the gall-bladder shadow is not definitely related to the type of bodily habitus.

2. The breadth of the gall-bladder shadow is greater in the "better" types of bodily habitus (that is, in the hypersthenic and sthenic types).

3. The neck of the gall-bladder shadow lies at a higher level and at a greater distance from the mid-line of the vertebral column in the "better" types of bodily habitus.

4. The fundus of the gall-bladder shadow lies at a higher level and at a greater distance from the mid-line in the "better" types of bodily habitus.

5. The rate of emptying of the gall bladder (as indicated by the extent of reduction in the size of the shadow at 1½ hours—that is, two hours after taking a fatty meal) is quicker in the "better" types of bodily habitus.

6. The respiratory excursion of the gall-bladder shadow is greater in the "better" types of bodily habitus.

It must be noted that, though this table indicates that the length of the gall-bladder shadow is not definitely related to the type of bodily habitus, the *actual* length of the viscus may vary with the type of bodily habitus, because in the radiograms we have a more foreshortened view of the gall bladder in the "better" types of bodily habitus.

Table of Results in Various Types and at the Intervals Specified after Administration by the Mouth.

Type of Bodily Habitus.	No. of Cases.	Length of Gall-Bladder Shadow (Inspiration).	Maximum Breadth of Gall-Bladder Shadow (Inspiration).	Position of Neck of Gall-Bladder at 1½ Hours (Inspiration).	Position of Fundus of Gall-Bladder at 1½ Hours (Inspiration).	Respiratory Excursion.
Hypersthenic	6	14½ hrs.: 4.9-9.8 cm. 16½ hrs.: 2.6-5.3 cm.	1½ hrs.: 2.8-3.7 cm. 16½ hrs.: 1.2-1.9 cm.	Upper part D12—lower border L1 3.5-5.8 cm.	Lower part L1—lower border L3 4.5-8.5 cm.	2.5-4.9 cm.
Sthenic	44	14½ hrs.: 4.9-8.5 cm. 16½ hrs.: 2.8-6.1 cm.	1½ hrs.: 2.6-3.7 cm. 16½ hrs.: 1.2-2.8 cm.	Lower border D12—middle L2 2.2-5.6 cm.	Upper part L2—upper border L4 3.8-6.9 cm.	2.5-4.2 cm.
Hyposthenic	17	14½ hrs.: 5.4-8.0 cm. 16½ hrs.: 3.5-5.9 cm.	1½ hrs.: 2.5-3.5 cm. 16½ hrs.: 1.4-2.7 cm.	Middle L1—lower part L2 2.7-4.8 cm.	Lower part L2—middle L4 3.5-5.0 cm.	1.9-2.7 cm.
Asthenic	5	14½ hrs.: 6.0-9.8 cm. 16½ hrs.: 4.9-7.8 cm.	1½ hrs.: 2.0-2.6 cm. 1½ hrs.: 1.1-.9 cm.	Upper border L2—lower border L2 2.7-4.0 cm.	Middle L1—lower border L4 3.4-4.5 cm.	1.5 cm.
	72					

J. S. GOODALL AND L. ROGERS: THYROTOXIC MYOCARDITIS.

Microscopic appearance of a section of the wall of the left auricle of a woman, aged 21, who died from acute Graves's disease.

E. D. TELFORD AND J. S. B. STOPFORD: THROMBO-ANGIITIS OBLITERANS.

F. DAVIES: NORMAL CHOLECYSTOGRAPHY.

FIG. 1.

Hypersthenic.

Sthenic.

Hyposthenic.

Asthenic.

Fig. 2 shows the areas including the varying positions of the neck and fundus of gall-bladder shadows in each type of bodily habitus—for example, the neck and fundus of the gall-bladder shadow of a hypersthenic individual will normally be found to lie within the area 1. Fig. 3 shows an average gall-bladder shadow for each type of bodily habitus, the measurements being obtained from the averages of those in the table.

Mills showed that gastric motility and tonus varied with bodily habitus, the motility being fastest in the "better" types. The rate of emptying of the gall bladder is shown above to be quicker in the "better" types of bodily habitus. A definite correlation is thus indicated between gastric motility and rate of emptying of the gall bladder.

2. Method of Administration.

A dose of 5 grams of sodium tetra-iodo-phenol-phthalein was given to each student. It was made up in five gelatin capsules, each of which was coated a few hours before administration with stearic acid, the coating being thin and uniform. The capsules were swallowed at intervals during the course of the 7 o'clock evening meal, a fair quantity of water being drunk with the meal. To minimize evil effects, the student was advised to remain comparatively quiet during the evening. The following morning, without breakfast, the student presented himself for x-ray investigation at 9.30 o'clock (that is, 14½ hours after taking the dye). Then he took a fatty meal consisting of eggs, bacon, milk, cream, butter, etc., and at 11.30 presented himself for the second x-ray investigation (that is, 16½ hours after taking the dye, or 2 hours after the fatty meal). He then resumed his normal life. A dose of 5 grams is quite sufficient, as it was noted in a number of cases which showed a good gall-bladder shadow that some of the dye was still residing unabsorbed in the large intestine. Moreover, the more serious of the toxic symptoms that have been reported have generally been observed after larger doses. In this series of cases, with the above dosage and method of administration, toxic symptoms resulted in very few instances, and in these they were very mild, not sufficient to interfere with the work of the student or to cause any student to refuse to take the dye a second time if required. The symptoms comprised slight nausea shortly after swallowing the capsules (in 8 per cent.), slight vomiting soon after taking the dye (in one case), slight epigastric discomfort during the

night (in 7 per cent.), slight headache in the morning relieved by breakfast (in 10 per cent.), or a nasty taste in the mouth in the morning variously described as "iodoform" and as "metallic" (in 10 per cent.). The dye is a natural purgative, and slight laxity of the bowels (not actually amounting to clinical diarrhoea) was exhibited in 20 per cent. of the cases. Recently I have been using a form of sodium tetra-iodo-phenol-phthalein which is almost free from toxic effect when administered as above described. Slight headache may occur, but is quickly dispelled by 10 grains of aspirin.

3. Radiological Technique.

The following points are emphasized in the standardized technique which we have gradually come to employ. For this credit is due to Mr. Frederick Melville, radiographer at University College, London.

1. The tube should be centred over a point about a hand's breadth to the right of the spine of the second lumbar vertebra.
2. The radiogram is taken postero-anteriorly (P.A.)—that is, abdomen towards film.
3. The radiogram is taken at the end of inspiration.
4. The use of the immobilizing band is no definite advantage.
5. The Potter-Bucky diaphragm is always used.
6. Four types of tubes were used with no material difference in the results.
7. Tube condition: 30 milliamperes, 5-inch spark gap, tube distance 36 inches, exposure 4 to 6 seconds (depending on the size of the individual).

It will be noted that the tube distance is great (36 inches), the result of which is to reduce distortion. It was observed that the best cholecystograms were obtained in those films which did not show first-class bone definition—that is, those in which a smaller tube penetration was used as a result of the small spark gap (5 inches).

Comment.

In this investigation a positive "good" shadow of the gall bladder was obtained in 72 per cent. of the cases. This figure includes only those shadows from which definite measurements could be taken. Ill defined or doubtful shadows are not included. The number of positive results in the earlier cases was affected by the fact that the technique was somewhat experimental. It was necessary to determine by experience the best method of administering the dye and the most suitable radiological technique. The later cases showed a larger number of positive results. When the improved (and now constant) technique was employed it was found that of fifteen students who failed to give a positive result on the first investigation, nine gave

FIG. 2.—Areas including the varying positions of neck and fundus of gall-bladder shadows for each type of bodily habitus. 1, Hypersthenic; 2, sthenic; 3, hyposthenic; 4, asthenic.

FIG. 3.—Average type of gall-bladder shadow for each type of bodily habitus (based on average position of neck and fundus and average breadth). 1, Hypersthenic; 2, sthenic; 3, hyposthenic; 4, asthenic.

a good gall-bladder shadow when the investigation was repeated six to twelve weeks later, using precisely the same technique. This is a matter of some importance, since many investigators have declared that the absence of a positive result with a reliable technique is the most certain evidence of disease of the biliary tract. The above evidence shows that a certain number of normal individuals may fail to give a positive cholecystogram on the first investigation, but give a good gall-bladder shadow on repeating the investigation some time later. Also in this series it was noted that two students who had previously given positive cholecystograms failed to give a shadow when the investigation was repeated a few weeks later, using the same technique. Evidently, therefore, the factors concerned in the production of a gall-bladder shadow must be liable to physiological variation. The following physiological factors are essential for the production of a positive cholecystogram after the oral administration of sodium tetra-iodo-phenol-phthalein:

1. A definite rate of absorption of the dye from the intestine.
2. A definite rate of specific excretion of the dye by the liver.
3. Entrance of the dye into the gall bladder in sufficient amount.
4. A definite degree of concentration of the dye-containing bile by the gall bladder.

It is possible that any of these factors may be subject to physiological variation which may affect the production of a positive cholecystogram. Have we any means of determining the adequacy of each of these factors and thus of observing any possible variations that may occur in the normal individual? It might be possible to obtain an index of the rate of absorption of the dye from the intestine by observing the rate of elimination in the urine (a small fraction of the dye being eliminated by this channel). The dye has a slight laxative action on the bowel and the greater part of the dye is eliminated in the faeces. No definite correlation between the degree of laxative action of the dye and the density of the shadow of the gall bladder was noted in this series. The rate of excretion of the dye by the liver can be checked by Rowntree's method of estimating the "global capacity" of the liver to eliminate similar dyes—for example, phenoltetra-chlorophthalein. This is done by injecting intravenously a known quantity of a solution of this dye and estimating its rate of disappearance from the blood stream, or its rate of appearance in the contents of the duodenum aspirated through an Einhorn's duodenal tube. The filling of the gall bladder with dye-laden bile from the liver is brought about by a mechanism at the termination of the common bile duct which resists the passage of bile direct to the duodenum. It is possible that this resistance mechanism may be liable to physiological variation. As Mann³ states, it is not known what percentage of bile normally enters the duodenum without previously entering the gall bladder. The dye in the bile leaving the liver is not sufficiently concentrated to produce a shadow. The concentration takes place in the gall bladder by the absorption of water by the wall of the viscus. For a sufficient degree of concentration to take place the wall of the gall bladder must be healthy, but the process may be liable to physiological variation. The degree of absorption of water by the gall bladder may be affected by the water content of the tissues at the time. There may be other factors, besides those mentioned, which are necessary for the production of a gall-bladder shadow, and physiological variations in these may likewise modify the results of an x-ray investigation of the biliary tract.

The complexity of the physiological processes underlying the production of a gall-bladder shadow is such that even on *prima facie* grounds the statement that absence of a shadow (*per se*) always indicates a pathological state seems unwarranted. Moreover, the above observation that a normal individual may fail to give a shadow on the first investigation but give a positive result on repeating the investigation is in direct conflict with such a statement. It thus appears that a negative cholecystogram *per se* (that is, in the absence of clinical evidence of disease of the biliary tract) is of less diagnostic value and significance

than positive cholecystographic evidence of disease. Likewise, if a shadow be produced, the above factors will undoubtedly have an influence upon the density of the shadow, so that too much weight must not be attached to diminished density of the shadow as an indication of a pathological state.

Summary and Conclusions.

1. The position of the gall-bladder shadow in cholecystograms varies with the type of bodily habitus. (See Fig. 2.)
2. The rate of emptying of the gall bladder varies with the type of bodily habitus. A correlation is established between bodily habitus, gastric motility, and the rate of emptying of the gall bladder.
3. The oral method of administration of sodium tetra-iodo-phenol-phthalein, using the technique described, is reliable, safe, and comparatively free from toxic effects.
4. The radiological technique described is reliable.
5. A certain number of normal individuals may fail to give a shadow on the first investigation, but on repeating the investigation with precisely the same technique many of these may give a positive result.
6. A negative cholecystogram, in the absence of clinical evidence, is less reliable than positive cholecystographic evidence of disease.

I desire to express my thanks to my chief, Professor G. Elliot Smith, at whose suggestion this work has been carried out, and to Mr. H. A. Harris for valuable advice and help in the investigation. I also wish to express my appreciation of the medical students of University College who offered themselves as subjects for this investigation.

REFERENCES.

- ¹ Graham, E. A.: BRITISH MEDICAL JOURNAL, October 16th, 1926, p. 671. (In this paper an extensive bibliography is given.) ² Mills, R. W.: *Amer. Journ. Roentgen*, 1917, 4, 155-169; and 1922, n.s., 9, 731-734. ³ Mann, F. C., and Kirklin, B. R.: *Proc. Staff Meet. (Mayo Clinic)*, January, 1927.

TWO CASES OF THROMBO-ANGITIS OBLITERANS IN WOMEN.

BY

E. D. TELFORD, M.A., F.R.C.S.,

PROFESSOR OF SYSTEMATIC SURGERY, UNIVERSITY OF MANCHESTER;

AND

J. S. B. STOPFORD, M.D., F.R.S.,

PROFESSOR OF ANATOMY, UNIVERSITY OF MANCHESTER.

(With Special Plate.)

THE symptoms of thrombo-angiitis obliterans are already becoming widely known, thanks to the reporting of cases by several observers, and it is not likely that much more is to be gained by the bare recording of further instances. The rarity of the lesion in women is, however, one of the extraordinary features of the disease. Buerger,¹ to whom much of our knowledge of the subject is due, states: "So rare is the disease in females that doubt has been cast on the possibility of its occurrence outside of the male sex." He further states that he has observed only three cases.

In the course of further investigation of thrombo-angiitis obliterans we have encountered two instances of the disease in women. These we think worthy of being recorded, not only by reason of their rarity in that sex, but because each presents certain very unusual features.

Dr. Williams of Salford drew our attention to the case of a woman patient who had continued for no less than twelve months with the left hand, wrist, and part of forearm in a condition of total mummification. By the kindness of Dr. Williams we were able to make an examination of the case and to obtain a photograph. Short details of the case are as follows:

A married woman, aged 52, British born, stated that her trouble began, some fifteen months ago, with severe aching pain in the index and middle fingers of the left hand; she noticed that they were "slightly blue." Within three weeks the whole hand was aching very severely; it was "cold and dark red." She continued in this way for three months, and at the end of this time the hand, wrist, and lower part of the left forearm were in a condition of black-dry gangrene. The pains then ceased, and when we saw her the condition of the limb had shown no change