and of the concomitant fall of sphygmometer pressure varying from 30 to 75 or more millimetres. To me the judgement to be pronounced is clear, plain, and inevitable.

CAUSATION.

As to causation, let me refer for a moment to the term "hyperpiesis," introduced to define cases of raised sphygmometer readings in persons without arterio-sclerosis and with no evidence of kidney lesion. They were therefore inexplicable, and were labelled with this name. I have seen many cases of this kind, but I never saw one without palpable constriction of the brachial artery. But the users of this word, and the exponents of what is called "primary and essential hypertension"—hypertension meaning sphygmometer readings—do not recognize that a constricted normal artery, with its living and thickened wall, and lessened lumen, materially affects sphygmometer pressures. I have stated that Janeway and Park gave over 60 mm. for hypertonic contraction in an unthickened artery; the first three cases recorded here gave relaxation readings of 25, 30, and 25 mm.—and that figure after the left ventricle had improved. In Case IV, after the initial fall of 45 mm., the extreme sensitiveness of the arterial wall gave sphygmometer readings which varied from day to day between 120 or 130 and 190 mm. If heart power was being recorded in this case the readings would have indicated an erratic variation of heart power of from 90 to 100 mm. without any indication that the left ventricle was responsible.

As to the influences invoked to account for high pressures, Professor MacWilliam² in a recent lecture says:

"While it is obvious that some powerful agency is at work, the search for definite pressor bodies (from any source) in the circulation has proved to be singularly inconclusive. Clifford Allbutt said: 'Such catch phrases as intestinal toxins, kidney inadequacy, endocrine disorder, and the like are but dust in our eyes.'"

As a clinician who for many years has watched and measured the constriction and relaxation of the arteries examined for clinical purposes, I have learnt the great sensitiveness of this arterial function. The physiological conception that it is mainly, if not wholly, to maintain or regulate blood pressure is much too restricted a view. The arteries respond not only to cardiac output and to nervous influences, but to influences from the alimentary tract, including excess of protein food, wine, tobacco, insufficient removal of hepatic excretion, constipation, and so on. The subtlety of the chemico-vital composition of the blood in different individuals is a fact of which the observant physician has abundant proof, although the laboratory may not be able to explain it or define it by analysis. Biochemistry has done much; but can chemical analysis of the blood determine minor degrees of what is known as acidosis? Yet the clinician removes symptoms, and relaxes hypertonus, by the administration of sodium bicarbonate. So also with treatment directed to biliary secretion and suitable colon emptying; in other cases to the reduction of protein food; in others abstention from wines, and so forth. In all these classes the only feasible explanation is that there is a chemical substance present in the blood which acts directly on the arterial wall. Clinical medicine cannot wait for the advance of biochemical analysis for the confirmation of treatment results. Supposing that large protein feeding acted by increasing the secretion of adrenaline, could biochemistry show its increase in the systemic blood? The same question can be asked with regard to other possible constituents. As a clinician, with some understanding of chemical pathology, I entirely repudiate the doctrine that "such catch phrases as intestinal toxins, kidney inadequacy, endocrine disorder, and the like are but dust in our eyes." Any one of them, or any combination of them, can, and does, cause arterial constriction; and that, as we have seen, means a rise in sphygmometer reading.

The clinical observation which it is most important for us to train ourselves to make is the tightening up of the arteries we examine for clinical purposes. Arterio-sclerosis is the result of more or less continued tightening up in persons whose hearts readily respond to extra strain. It takes years to develop; and it can be checked if tightening

is recognized and means are taken to eliminate the factor which is the prime cause. The heart hypertrophy is the response of the heart to the extra strain of driving blood through constricted channels. Kidney changes develop of an interesting and informative type. The renal artery outside the kidney shows the type of change seen in Figs. 3 and 4, whilst the divisions of the artery inside the organ show an irritative and progressive endarteritis. This goes on to the "gouty kidney" of last century, the arterio-sclerotic kidney of a later time.

REFERENCES.

- ¹ W. Russell: *The Sphygmometer*, Baillière, Tindall and Cox, p. 79.
- ² Professor MacWilliam: *BRITISH MEDICAL JOURNAL*, January 22nd, 1917, p. 125.

THE HISTOLOGY OF THE PERIODONTAL SULCUS.

BY

J. G. TURNER, F.R.C.S.,

DENTAL SURGEON AND LECTURER ON DENTAL SURGERY.

(From the Research Laboratory of the Royal Dental Hospital of London.)

(With Special Plate.)

THE importance of the periodontal sulcus as being the seat of origin of pyorrhoea will be patent to all who have grasped the significance of dental sepsis, yet it is a fact that there are no definite data as to its intimate structure. The chief point at issue has been how is epithelial continuity provided for in the normal condition? Does the gum epithelium attach itself to the tooth, or is there normally a breach of epithelial continuity where the tooth projects? In the latter case the apex of the sulcus would be a serious defect in our defensive armament.

For many years I failed in my efforts to get a section of the normal periodontal sulcus, owing to the dissociation of the tissues which occurs during decalcification and sectioning. A few years ago, however, I hit on the plan of embedding in celloidin before decalcifying, and have been able to get some very useful sections. Having no access to the dead-house I have had to rely on the forceps for my material. Obviously the forceps destroys the lingual and labial parts of the periodontal sulcus, so that my observations relate only to the interstitial part. In many cases a piece of the interstitial gum comes away attached to the tooth, probably more or less dissociated, but possibly intact. I have had a large number of such specimens sectioned, and have been led to the conclusion that interstitially at least there may be, possibly should be, no periodontal sulcus. I have found a considerable number of cases in which the gum is attached directly to the tooth-crown, and the attachment has been firm enough to resist disruption at the moment of extraction. Some of these cases show that there was no possibility of a sulcus (Fig. 1), the gum epithelium being prolonged on to the tooth-crown and fusing with Nasmyth's membrane; others leave a possibility that there was a shallow sulcus, not, however, at the neck of the tooth, but in a position halfway along the crown (Fig. 2). Some sections, again, suggest a potential rather than an actual sulcus (Fig. 3).

The question of epithelial continuity may be definitely answered. There should be no breach until the dentine of the tooth is exposed, or the neck denuded. The gum epithelium may spread over the tooth-crown, much as the skin epithelium over a nail (Fig. 1), becoming continuous with Nasmyth's membrane over a wide area; or it may be continuous with Nasmyth's membrane only at the edge, leaving the fibrous gum to take direct attachment to the tooth-crown (Fig. 5); or for a varying distance it may be split away from Nasmyth's membrane, taking a vital attachment only as the neck of the tooth is approached (Fig. 6); or it may take attachment only to the neck of the tooth—that is, to the cementum (Fig. 7). The first and last two cases leave a possibility of a periodontal sulcus which would develop as the gum epithelium split away from Nasmyth's membrane, possibly as the latter became less vital owing to ageing, or as the gum epithelium peels off

FIG. 1.—From a temporary molar patient, aged 12. *a*, Gum epithelium deeply stained. *b*, Nasmyth's membrane staining badly.

FIG. 2.—From a partly erupted permanent molar. N.M., Nasmyth's membrane broken in sectioning. The apex of the sulcus was at *A*, well up the crown of the tooth. *K*, Epithelial prolongation along root. *J*, Junction of gum epithelium with remains of outer portion of enamel organ and Nasmyth's membrane.

FIG. 3.—Temporary canine patient, aged 7. The gum epithelium is attached to the tooth, but between it and Nasmyth's membrane is a line of demarcation (*b*) along which splitting can readily occur. There is thus a potential sulcus.

FIG. 4.—A higher power of *a* in Fig. 3. It suggests stellate epithelium, and so the outer layer of the enamel organ as its origin.

FIG. 5.—A bicuspid patient, aged 12. The fibrous gum is attached directly to Nasmyth's membrane, and there was epithelial continuity only at the edge.

FIG. 6.—Temporary canine patient, aged 12. Nasmyth's membrane is badly stained (necrotic) and vital attachment of gum epithelium to tooth obtained only towards the neck. The sulcus was rather potential than actual.

FIG. 7.—Partly erupted lower wisdom. The epithelium of the gum was attached to the root from *a* to *b*. The cleft is artefact, the root between *a* and *b* being coated with epithelium seen in thin black patches in the figure. Almost to the last the gum epithelium shows traces of papillary structure.

FIG. 8.—Unerupted incisor in a girl aged 9½. Nasmyth's membrane is well defined. The remains of the outermost layer of the enamel organ are seen as a rather discontinuous epithelial layer with small columns and coils reaching out into the adjacent gum. There is no epithelial prolongation along the root.

FIG. 9.—A higher magnification of *b* in Fig. 8 where the two epithelial layers have been split in sectioning.

the root. The sulcus is thus seen to be a cleft left by the splitting away of gum epithelium from Nasmyth's membrane. When the gum epithelium has split clear away from the whole length of the tooth-crown, or where it never has united with Nasmyth's membrane, it may take a more or less extensive attachment to the root of the tooth (to the cementum).

Thus, though technically there is a breach of epithelial continuity where the attachment of gum to tooth has receded beyond the enamel, practically there is no breach in the defence till the enamel has been worn through to the dentine (by caries or attrition) or till disease (pyorrhoea) has destroyed the epithelial attachment to the tooth.

The nature of the epithelial prolongation which provides for epithelial attachment to the tooth-root is worth further notice.

It is not always present (Fig. 8), and in such cases, if it be due to develop later, it must be provided by the gum epithelium, as was probably the case in Fig. 7, where the gum epithelium can be seen gradually losing its papillary character as it passes along the root. In many cases, however, it seems to have been provided for from the beginning. Most of my sections of this point are taken from temporary teeth, and it is worthy of note that there is no difference in extent of prolongation between teeth about to be shed and those of earlier age (Figs. 3 and 6). There is thus no evidence in the temporary teeth of "continuous eruption." Fig. 7 shows epithelium whose gum origin is indicated by the gradual fading away of the papillary structure; but Figs. 3 and 6 from temporary teeth, and Fig. 2 from a permanent molar, suggest a different origin. A higher magnification (Fig. 4) of the prolongation strongly suggests stellate epithelium—a form of epithelium more likely to arise from epithelium originating from the enamel organ than from the gum. As Figs. 8 and 9 show, there is at the time of eruption an epithelial lining to the tooth-crypt derived from the outermost epithelium of the enamel-forming organ. To allow gum epithelium to take attachment to the tooth this epithelial lining must either atrophy or blend with the gum. I have not been able to decide which, but am inclined to think it generally atrophies, except at its periphery, where it remains capable at need of further growth.

THE HEALING OF GASTRIC ULCERS: RADIOLOGICAL OBSERVATIONS.

BY

F. G. NICHOLAS, M.B., B.S.LOND.,
D.M.R. AND E.CAMB.,

ASSISTANT RADIOLOGIST TO THE MIDDLESEX HOSPITAL; RADIOLOGIST
TO ST. LUKE'S HOSPITAL, CHELSEA;

AND

ALAN MONCRIEFF, M.D., M.R.C.P.LOND.,
MEDICAL REGISTRAR, MIDDLESEX HOSPITAL.

(With Special Plate.)

THE radiological diagnosis of ulcers of the stomach dates from the classical description by Haudek¹ of the characteristic "niche" seen in the outline of the stomach after an opaque meal. Much interest has been attached to the disappearance of this "niche" under treatment, and many papers have recently appeared on the subject of the "healing" of gastric ulcers. The terms "healing" and "healed" with reference to gastric ulcer are open to a certain amount of criticism; but it must be accepted that disappearance of a "niche" associated with clinical improvement, disappearance of pain and tenderness, disappearance of occult blood from the stools and vomit, gain in weight, must certainly indicate a healing process, as Hamburger² has pointed out. If the healing process continues and no recurrence takes place over a period of, say, two years, many observers would consider such cases as "healed." There is certain histological evidence to support this view, and the matter is more fully discussed later.

During the past two years we have investigated a group

of cases by means of repeated opaque meals, and in this paper we deal with the results. Seventeen cases were selected in which x-ray examination disclosed a definite "crater" on the lesser curvature of the stomach. Certain cases occurring at the time when this investigation was commenced, in which an ulcer was present on the anterior or posterior wall of the stomach, have not been included, since it was found that unless an ulcer could be shown in profile in a skiagraph it was very difficult to estimate any change in size at subsequent examinations. The cases included in-patients and out-patients, but in many instances the attendance of the latter for treatment was very irregular, and no deductions as to the success or failure of medical treatment are made in this paper, nor should such deductions be legitimately made on the evidence put forward.

The "double meal" method has been employed. Two nights before the examination an ounce of castor oil is given, and the evening before a light meal is allowed. Six hours previous to the time arranged for radiography three ounces of barium sulphate are given well mixed with bread-and-milk.

At the screen examination notice is taken of the position of the head of the meal and whether a gastric residue is present or not. Occasionally there has been a residue of barium in an ulcer crater. Next a mouthful of thick barium emulsion is administered, and this is used to "whitewash" the whole of the stomach according to the method described by Scott.³ After a careful examination of the whole of the stomach by this method the patient continues to take barium emulsion until the stomach is filled. The patient is then rotated, and the stomach examined from all angles. A skiagram is taken at that angle which shows the maximum size of the crater with the minimum of distortion. At the re-examination great care is taken that the distance between target and plate, and the position of the patient, are the same.

Results.

Of the 17 cases in this series all were examined on two occasions, and 15 on three occasions, the intervals between the examinations varying from one week to one year.

TABLE I.—Results of First Re-examination (17 cases).

(1) "Healed." No evidence of ulceration ...	11
(2) "Healing." Definite decrease in size of ulcer...	4
(3) "Stationary." No alteration in size ...	2

Average period elapsed since primary examination, six months.

The 11 cases in which no ulcer crater could be seen were all clinically free from symptoms, and in good health. The 4 cases in which the ulcer crater was smaller in size were all improving, although the patients were not altogether free from pain. In one of these cases, although the "niche" on the lesser curvature had disappeared, there was a persistent incisura opposite to the original site of the ulcer and symptoms were still present suggesting that healing was not complete.

TABLE II.—Results of Second Re-examination.

(1) No x-ray evidence of presence of ulcer in 11 cases, made up as follows:	
"Healed" cases without recurrence ...	6
"Healing" cases now "healed" ...	4
"Stationary" case now "healed" ...	1
(2) Ulcer again present ("healed" cases now relapsed) ...	4

Two cases did not attend; one shown as "healed" and the other as "stationary" in Table I.

Average period between the two re-examinations—three and a half months.

It will be seen from this table that in 4 cases reported as "healed" in the first examination definite relapse had occurred. The clinical symptoms and signs had reappeared in all these patients, although at the time of the first re-examination, when no ulcer crater was seen, all these 4 cases were clinically free from symptoms. In one of these cases the ulcer crater was definitely larger than before. Of the 11 cases reported as "healed" in Table II, 6 represented patients in whom the ulcer had remained healed over the whole period of investigation, and the others those in