may occur—how an individual may hide his head in the sand and refuse to face reality, crying peace when there is no peace; but that he would get beneficial results of a permanent nature by such a line of conduct is another matter. An appeal to narcissism might explain the state of mind of a man who stands in the corner of an asylum ward, taking no notice of anyone, but just immersed in his own fancy and in his own mental aberration; or the state of mind of the hypochondriac, with distribution of libido so disturbed that even his bodily organs cannot function adequately or without pain. But that good, or therapeutic, auto-suggestion is of this nature I cannot believe. Auto-suggestion is based on confidence and the abolition of fear. The individual who uses auto-suggestion knows this, and it is definitely admitted that auto-suggestion works better in a more healthy person; the healthier he is the better effects he can get. Auto-suggestion is a method of will training.

In holding such a view the possibility is not excluded that in hypnosis archaic affective tendencies and beliefs may come to the surface and make themselves felt, and that an individual may produce certain results by virtue of the strength of these primitive beliefs which are not justified by modern scientific knowledge, although such beliefs may make a great difference to the individual himself. In him there may be a desire for the miraculous—a desire to believe in the miraculous—which may be liberated under conditions of hypnosis, or suggestion treatment, and may produce results; not that it will produce a miracle, but it may counteract the excessive scepticism and doubt and despair which are so often characteristics of the modern mind. Consequently, bad, or pathogenic, auto-suggestion is perhaps the best example of auto-suggestion in a literal sense—that is, the working of an idea under the influence of an instinctive tendency, in this case one of fear and apprehension. Freud might say that fear is due to bad conscience, and that bad conscience is due to the working of the Oedipus complex, so bringing in the libido doctrine. Therapeutic auto-suggestion goes further, because it replaces the feeling of doubt and fear by a feeling of confidence. The result suggested can then occur, so far as it does not conflict with any of the fundamental laws of nature, and can occur to a maximum degree. The instinct of self-assertion (supported by its own distinctive form of energy, not by libido) may play a prominent part in bringing about a realization or actualization of the suggestion; but the integrated personality, with its entire reserves of instinctive energy, may, by an act of faith, fulfil more adequately the purposes towards which auto-suggestion is half blindly pointing.

REFERENCE.

¹ *Mind and Personality*, University of London Press, Ltd., 1926, pp. 180, 272.

THE CAUSATION OF "STRIAE ATROPHICAE CUTIS" ("VERGETURES") NOT DUE TO STRETCHING OF THE SKIN.

BY

F. PARKES WEBER, M.A., M.D., F.R.C.P.

In all ordinary cases striae atrophicae or striae cutis distensae can be explained as representing a method of cutaneous accommodation towards distension (physiological or pathological), when the skin is relatively insufficient—that is to say, unable to adjust itself by its normal elastic distensibility. In such cases a relatively too rapid growth or increase in bulk of the parts below the skin, or a cause of chronic distension of some kind—such as a relatively excessive growth of subcutaneous fat, muscle, or long bones (especially connected with puberty), mammary gland, the pregnant uterus, ascites, tumours, or subcutaneous oedema—leads to a splitting or cleavage in the deeper parts of the cutis, manifested on the surface by the appearance of striae atrophicae, which are at first red or purplish, and in time gradually become white and smaller as they heal up.

In the present note I shall refer only to those cases of striae atrophicae not due to stretching of the skin; that is

to say, the cases in which striae atrophicae have developed apparently in the absence of any special distension—in the absence of pregnancy, ascites, tumours, subcutaneous oedema, excessive subcutaneous fat, and other usual causes. In such cases the ordinary growth of the long bones, muscles, and subcutaneous fat seems to be associated with some insufficiency of the cutis (with perhaps deficiency of elasticity) which prevents the skin from adjusting itself to the parts beneath, excepting by the process of multiple cleavage—that is to say, by the formation of striae atrophicae. Striae atrophicae of this mysterious kind occur occasionally in apparently healthy individuals about the period of puberty, the skin failing to keep pace with the normally rapid growth of certain parts of the body, such as the shoulders, pelvis, and thighs. Striae atrophicae, in the absence of ordinary causes, are, however, still better known to occur in connexion with enteric fever (notably the "striae patellares") and certain other microbic infections, although even in such cases the selective distribution of the striae, and whether they are symmetrical or more or less unilateral or not, as I have elsewhere endeavoured to explain, must be due partly to mechanical causes, such as the position of the patient in bed, arrangement of pillows, etc.¹

What is the cause of the relative insufficiency of the cutis in these cases to which I am referring?

Here I must speak of a condition of abnormal relative or absolute redundancy of the skin which may be contrasted with, and in some respects is the opposite of, the relative or absolute cutaneous insufficiency manifested by striae atrophicae. I refer to cutis striata or cutis sulcata—that is to say, ploughed or furrowed skin—"bulldog scalp," and cutis verticis gyrata. In these cases the scalp, especially the skin of the scalp, is locally altogether too big for the skull beneath it, and has therefore to become folded into ridges and furrows, so as to adjust itself to the surface which it covers. A local suboccipital variety of furrowed scalp may be etiologically connected with shortness of neck,² and in certain idiots complete furrowed scalp has been associated with more or less microcephaly,³ though I think the microcephaly was insufficient entirely to account for the scalp condition.

In twelve cases, however, the general redundancy of the skin and subcutaneous tissue of the scalp which gave rise to a condition of cutis verticis gyrata was connected with acromegaly,⁴ and was doubtless a manifestation of the functional (hormonic) over-activity of the anterior lobe of the pituitary gland, which is the essential cause of the main features of acromegaly, including skeletal and visceral overgrowths. Though cutis verticis gyrata occurs in relatively few cases of acromegaly, it must nevertheless, when it is present, be regarded as a manifestation of the acromegalic process, just as enlargement of certain viscera ("splanchnomegaly") must, though likewise not occurring in every case.

I suggest that, as the redundancy of the cutis present in certain cases of acromegaly is obviously due to excessive hormonal activity of the anterior lobe of the pituitary gland, or of an adenomatous tumour (an "endocrine-tumour") arising from it, the insufficiency of the cutis manifested by striae atrophicae cutis, when not caused by special distension, may be due to a defective hormonal activity of the same endocrine gland, for which, perhaps, a depressing action of the toxins of enteric fever and other microbic infections may be sometimes responsible. This suggestion is in accordance with the occurrence of obesity (of a cerebral or pituitary type) occasionally observed as a sequel to enteric fever and other microbic infections.

REFERENCES.

- ¹ On this whole subject cf. F. Parkes Weber: "Unilateral Striae Atrophicae," *Practitioner*, London, 1917, vol. 99, p. 453; also *idem*, *Cutaneous Striae, Brit. Journ. Derm. and Syph.*, London, 1926, vol. 38, pp. 1-19; and in discussions, *Proc. Roy. Soc. Med.*, Derm. Section, 1924, vol. 17, p. 45; and *ibid.*, Section for the Study of Disease in Children, 1927, vol. 20, p. 11.
- ² F. Parkes Weber: Suboccipital Furrowed Scalp (the Suboccipital Type of Cutis Striata or Cutis Sulcata), Royal Society of Medicine, Dermatological Section, Meeting of January 19th, 1928.
- ³ T. W. McDowall: *Journ. Mental Science*, London, 1893, vol. 39, p. 62; J. J. Cowan, *ibid.*, 1893, vol. 39, p. 539. See also, on cutis verticis gyrata of mechanical origin, C. Adrian and A. Forster, *Arch. f. Derm. u. Syph.*, Berlin, 1919-20, vol. 127, pp. 767-790.
- ⁴ F. Parkes Weber: A Note on the Nature of Cutis Verticis Gyrata, particularly in Acromegaly, *Brit. Journ. Derm. and Syph.*, London, 1928, vol. 40, pp. 1-11.