ureteral calculus a few hours before the time fixed for his operation. In another case the renal stone was removed from the pelvis and olive oil injected down the ureter by a glass syringe, the nozzle of which was passed through the pyelotomy incision and fitted snugly into the upper part of the ureter. The ureter became distended and this subsided suddenly. The cystoscope was passed at the end of the operation and a small calculus was seen in the bladder lying in a pool of oil. This was passed a few days later.

I have endeavoured to give, as accurately as possible, the late results of these operations for renal calculus. It is very difficult indeed to obtain absolute accuracy of statement on such a subject unless one has the advantage of a large staff to carry out the necessary inquiries and to make the necessary investigations. There is, however, sufficient evidence to show that the frequency of recurrence after operations for renal calculus is such that it demands very serious investigation.

REFERENCES.

¹ Levy: *Journal d'Urologie*, xviii, No. 1, p. 1. ² Marion: *Ibid.*, xxi, No. 1, p. 61.

PERNICIOUS ANAEMIA AND THE LIVER DIET.

BY

C. F. T. EAST, M.A., M.D. OXON., F.R.C.P. LOND.,

JUNIOR PHYSICIAN, KING'S COLLEGE HOSPITAL; PHYSICIAN, WOOLWICH WAR MEMORIAL HOSPITAL.

THE following is an account of the history of three patients treated by the liver diet introduced by Minot and Murphy¹ in 1926. Recent accounts from all sides are showing the good results obtained in promoting remissions in cases suffering from severe relapses. These three patients have been under treatment for a year or more. The cases are recorded in order to show what may happen afterwards.

CASE I.

A man, aged 45, had an attack of gastro-enteritis in November, 1925, which lasted several weeks. Six months later his tongue became sore, and during the summer his throat and tongue were very sore from time to time. Early in the autumn the nausea and vomiting returned, and when admitted to hospital in October, 1926, he presented the typical picture of pernicious anaemia. His mother had had anaemia for a long time, and eventually died from it; her sister was definitely known to have died of pernicious anaemia.

Investigation showed a characteristic blood count. The blood bilirubin was high. There was complete achlorhydria. A haemolytic streptococcus was isolated from the duodenal contents, but it could not be found in the faeces. From the first he was given arsenic, and hydrochloric acid in large doses, as recommended by Hurst.

After a month in hospital he was given liver; owing to his gastric symptoms he had great difficulty in taking it, and the ingenuity of the nursing staff was fully exercised in making it palatable. In one form or another he managed to take about 3 ounces a day. An autogenous vaccine from the haemolytic streptococcus was also administered.

The improvement in the blood picture began at once, before the liver was given; this continued afterwards, especially as regards the haemoglobin. Shortly after this improvement began the haemolytic process abated and the blood bilirubin content fell, but did not reach normal limits. After leaving hospital he kept fairly well, and for a time was on his diet, and took arsenic and hydrochloric acid. After March he was able to return to work until August, when a relapse began, and the soreness of the tongue and vomiting returned. As soon as the nausea was less liver was again given, together with acid and arsenic. There was soon considerable improvement, again notably in the haemoglobin. The blood bilirubin, estimated by the van den Bergh test, fell to a low level, but the red cells improved less satisfactorily. He then left hospital by his own wish, and at home continued to take a proprietary preparation of liver similar to that which he had had in hospital.

An unfortunate relapse has again occurred, and the outlook seems far from favourable. The red cells and haemoglobin have fallen, and the blood bilirubin has again become high.

CASE II.

A woman, aged 54, had had bouts of diarrhoea for three years, and some anaemia was noticed a year before admission. For the last four months she had had a good deal of vomiting, and had been laid up in bed. On admission to hospital she was very ill. Petechial haemorrhages were present everywhere; the heart was large and feeble, and the spleen was increased in size. There was complete achlorhydria. The teeth were septic. Five blood transfusions were given during the first six weeks, and arsenic and hydrochloric acid were also prescribed. After a little time a steady improvement set in. Before leaving hospital she began the liver diet, and since then she has taken it regularly at least

three times a week. The red cells have never returned to a higher level than $3\frac{1}{2}$ million, but the haemoglobin has kept up to about 80 per cent. There has been a normal level of bilirubin in the blood for nearly a year. Before she went out the heart returned to normal size, and the spleen could no longer be felt. She has now felt very well for fifteen months, and has been able to do her housework.

CASE III.

A woman, aged 33, became ill three years prior to admission, the symptoms being mainly dyspeptic. After a year she began to have attacks of diarrhoea, and vomiting became frequent; at times her tongue was sore. A year later her appendix was removed, but without benefit. The anaemia gradually started about this time; for four months she had noticed jaundice. Lately her ankles had become swollen, and she had suffered much from shortness of breath. Her mother had died of pernicious anaemia.

On admission to hospital she was seriously ill; the tongue was sore, and she was markedly yellow. The heart was very feeble and much enlarged, a gallop rhythm being audible over the mitral area. The electro-cardiogram showed heart-block, with a P-R interval of 0.32 second. There was a good deal of numbness and tingling of the hands and feet. Complete achlorhydria was present, and from the duodenum a haemolytic streptococcus was grown; this was not present in the faeces. A vaccine prepared from this organism was given.

At first a blood transfusion was given and she was put on arsenic and hydrochloric acid. Improvement in the anaemia began at once, and was most noticeable in the haemoglobin. At the same time there was a fall in the blood bilirubin, so that, on leaving hospital, it was well below the normal. Liver was not given at first as she could not take it; later, when her stomach was less troublesome, she started the diet, and has kept rigidly to it ever since—a period of ten months. She takes at least 4 ounces of liver three times a week, but lately has been using an equivalent amount of a proprietary preparation. As soon as the anaemia improved her heart returned to normal, and the heart-block disappeared. She left hospital last March, and has since kept very well. The red cells keep about $3\frac{1}{2}$ million per c.mm., but the haemoglobin is maintained at a high level. Haemolysis has not risen above the normal limit.

During the last ten months she has had hydrochloric acid with her meals, and courses of arsenic from time to time. There has been no increase in the sensory symptoms.

COMMENTS.

The Administration of Liver.

Various methods must be tried if the patient cannot take the liver easily. A paste in a sandwich of dry bread is often satisfactory. Sometimes the taste can be covered by the use of one of the strongly flavoured sauces; it goes fairly well with marmite. Small lightly boiled lumps are taken fairly easily, or it may be eaten with lemon juice or lettuce.

The After-Treatment.

These three cases are not of much value in illustrating the effects of the liver diet in the acute stages. It is unfortunate that there is such difficulty in getting patients to take liver when the gastric symptoms are severe.

As to the value of liver in the after-treatment, when once a remission has been secured, Cases II and III bear out the claims of Minot and Murphy,² who have had patients on liver for much longer periods than eighteen months without relapse. The first patient seems to have relapsed because for some months he neglected to take any precautions. The rather poor success subsequently seems to show that in the later relapses the treatment may be less effective. The after-treatment is unquestionably extremely important: "once pernicious anaemia, always pernicious anaemia," potentially at any rate. If, as Hurst has so often suggested, the achlorhydria is the weak point in the patient's defences, it is most important for the patient to continue taking acid. It is true that difficulties with this arise in the acute stages, when the gastric symptoms are severe and the mouth and pharynx sore; but apart from these it is usually well borne. Dilute phosphoric acid may be less effective, but is often much easier to take. It is, perhaps, too much to suppose that the artificial acid barrier can entirely replace the natural in warding off further intestinal infection. Hence it is further of the greatest importance to keep watch against sepsis arising again in the mouth and nasopharynx. The continuance of the liver diet, whatever its mode of action may be, should help in preventing any tendency to relapse.

Prevention.

The history of the prodromal stages in these cases has been given at some length. It is worth while to emphasize this pre-anaemic phase, during which the patient may

suffer from the symptoms of a gastro-intestinal infection from time to time, over a period of several years. If it were possible to foresee the ultimate development of the anaemic phase much might be done to prevent the later stages of the disease from occurring. The family history may give a clue. In two of these cases other members of the family had had pernicious anaemia. In the case of a fourth patient (put on liver diet too late in the last stages), who died, her mother had also died of the disease, after a remission lasting nine years.

Attacks of gastro-intestinal symptoms, with perhaps glossitis, in anyone whose family history shows pernicious anaemia, are a portent of the gravest significance. Without the family history the triad "glossitis," "gastritis," "enteritis" should warrant the examination of the gastric juice for achlorhydria. If this be found the case should be looked upon as one of potential pernicious anaemia. Liver treatment seems to be proving of real value in the acute phases, and may be of assistance in preventing relapse. But many cases might be recognized in the pre-anaemic phase, and a long and dangerous illness be perhaps prevented.

REFERENCES.

¹ Minot, G. R., and Murphy, W. P.: *Journ. Amer. Med. Assoc.*, 1926, lxxvii, 470. ² Idem: *British Medical Journal*, 1927, ii, 674.

A FORM OF SENILE SEIZURE.

BY

HUGH BARBER, M.D.LOND., M.R.C.P.,
HONORARY PHYSICIAN, DERBYSHIRE ROYAL INFIRMARY.

WITHOUT any warning, an apparently healthy elderly person falls down unconscious. There may be some degree of clonic spasm of the face or limbs, perhaps more pronounced on one side. In a few minutes there follow loud shouting and violent behaviour, necessitating restraint, but possibly aggravated by it. After a variable time the patient regains consciousness and reason, although perhaps a little dazed and incontinent. On full recovery of his faculties he will say that the whole incident is a blank. The whole duration of the attack is perhaps about three-quarters of an hour, but may be considerably longer. This type of emergency is seen after the event usually in hospital practice, or only in the casualty department. It does not appear to be described in medical textbooks. Of six attacks that I have witnessed the notes of the following case describe a typical example.

A railway storekeeper, aged 66, was found unconscious at 8.30 a.m. on January 24th, 1928. He was put into an ambulance, in which he became very violent, with struggling and shouting, sweating profusely. At 9 o'clock in the casualty department of the Derbyshire Royal Infirmary he was very violent, and continued so until 9.45, when he vomited twice and became conscious, but was somewhat dazed. He was taken to the ward, where he went to bed quietly, saying everything was a blank since his last duty performed that morning. Examination showed him to be a very stout plethoric man; his tongue was furred, his teeth were decayed, and pyorrhoea was present. The lungs were emphysematous; the heart was normal, but sounds distant. Temperature normal, pulse 92; artery thick; blood pressure 130 mm. systolic and 90 mm. diastolic (two days later it was 170 and 110). There were no abnormal signs in the nervous system. The urine contained a trace of albumin occasionally; but the blood urea was only 48 mg. per cent., and the urea test gave a reading of 2.4 per cent. His mind was quite clear, but the head felt heavy. The day after admission lumbar puncture revealed normal cerebrospinal fluid. Five days later he could walk about quite well.

Of six cases (five men and one woman) all were between 60 and 75 years of age. There was some evidence of cardio-vascular degeneration; but they were robust, stout, somewhat full-blooded people, engaged in their duties. There had been some degree of hyperpiesis, but no serious renal disease. In one case the disposing cause of anger precipitated the attack, but otherwise there had been no history of effort, nor were their duties arduous. In five instances the patient had come to himself in about an hour; the sixth patient, after the violence had passed off, I saw walking up and down his room with the doctor and butler in attendance. He came round about three hours

after the seizure, without any knowledge of what had been taking place. The attacks have not recurred nor have vertigo or other symptoms developed. The expectation of life has been what one would deduce from the age and cardio-vascular condition, but does not appear to have been changed by the accidental occurrence of the seizure; nor have the activities been restricted.

It would seem that the attack is due to slight haemorrhage on the surface of the brain, perhaps below the pia mater, producing coma and cerebral irritation, analogous to what may be seen occasionally after concussion. The bleeding must be quite distinct from a free subarachnoid haemorrhage: the one lumbar puncture observation revealed normal fluid. That it is a vascular lesion seems probable, because of the sudden onset, and there is the analogy of epistaxis and retinal haemorrhage occurring in this type of patient. Uraemia does not seem probable; the recovery is too complete, and the urea in the blood was not unduly high in the one case tested. The type of patient is not suggestive of epilepsy, although the analogy of a convulsion from loss of cerebral function probably holds good. A temporary oedema may be the cause, although a local aura might be expected. A localized haemorrhage, either venous or arterial, seems the most probable explanation. Embolism or thrombosis is possible, but the situation of the lesion is probably in a region where anastomosis is very free.

A SIMPLIFIED METHOD OF ARM-TO-ARM
BLOOD TRANSFUSION.

BY

E. F. SKINNER, M.D., M.B., F.R.C.P.,
ASSISTANT PHYSICIAN, SHEFFIELD ROYAL HOSPITAL.

THERE are so many methods of carrying out the transfusion of blood, each worker having developed his own technique, that it is almost futile to suggest a new method. However, as one who has done a large number of transfusions of all kinds and by all methods, I feel justified in making a short practical note on the subject.

There can be no question nowadays of the efficacy of transfusion as a therapeutic measure for a variety of conditions, and each clinical condition has its supporters for one or other method of carrying out the operation; but also there can be equally no question that, granted facilities, the transfusion of whole blood direct from donor to recipient is the best procedure. This was the original method, but, owing to various technical difficulties, it has long been discarded in favour of the "citrate" or "defibrinated" method.

The transfusion of whole blood is hindered by the readiness to clot exhibited by blood immediately it leaves the endothelial surroundings of blood vessels, and, of course, if blood is to remain whole—that is, to retain all its physiological and physical qualities—this tendency to coagulation must be retained. It forms one of the most important properties of blood, and in cases of haemorrhage is the one quality most desirable in the blood to be introduced. To render the blood uncoagulable by citrate appears to me a distinct disadvantage when dealing with cases of severe bleeding, and it is in such cases, particularly in obstetrical emergencies, that transfusion offers its greatest and most obvious use.

I make a plea, therefore, for the development of whole-blood transfusion, and venture to assert that with the extremely simple apparatus described below anyone who is accustomed to venipuncture can carry out a whole-blood transfusion with ease and certainty.

The apparatus consists of a syringe of 5 c.cm. capacity, with a piston of special pattern having a groove along one side. The barrel of the syringe is closed at the bottom, but has two openings in the middle opposite to each other. When in action the syringe is so arranged that by a simple turn of the wrist the groove in the piston can be put into apposition with either of the two side openings in the syringe, and either draw in or expel its contents.